



Markov Chain Monte Carlo (MCMC) Methods

PHYS 574

April 5th, 2019

Prepared by: Mario Ivanov

Metropolis-Hastings Algorithm

Most common/Widely used

Metropolis-Hastings Algorithm

Most common/Widely used

Let's explore it with an example!

Metropolis-Hastings Algorithm

Most common/Widely used

Let's explore it with an example!


Python module: "emcee"

Prostate Cancer Test


- Blood test: Prostate-Specific Antigen (PSA)
 - PSA concentration in (ng/mL)

Prostate Cancer Test

- Blood test: Prostate-Specific Antigen (PSA)
 - PSA concentration in (ng/mL)
- Problems:
 - Ethnicity/Age/Gender
 - PSA velocity
 - False positives and biopsy complications



Sigmoid Model



Bayes' Theorem

$$p(B \mid A) = \frac{p(A \mid B) p(B)}{p(A)}$$

Bayes' Theorem

$$p(m, b, f \mid x, y, \sigma) \propto p(m, b, f) p(y \mid x, \sigma, m, b, f)$$

Bayes' Theorem

$$p(m, b, f \mid x, y, \sigma) \propto p(m, b, f) p(y \mid x, \sigma, m, b, f)$$

POSTERIOR



PRIOR

LIKELIHOOD

Metropolis-Hastings Algorithm

Draw a sample state from proposed distribution

x' from $p(m, b, f)$

where $x_t = \{m^0, b^0, f^0\}$


where $x' = \{m', b', f'\}$



Metropolis-Hastings Algorithm

Calculate the acceptance probability ratio

$$\alpha = \frac{p(x')}{p(x_t)}$$




Metropolis-Hastings Algorithm


Accept or Reject:


$$u \in [0, 1] \quad \alpha = \frac{p(x')}{p(x_t)}$$


$$u \leq \alpha, \text{ accept : } x_{t+1} = x'$$


$$u > \alpha, \text{ reject : } x_{t+1} = x_t$$




















Gibbs Algorithm

Draw a sample state from proposed distribution

$$\{m^0, b^0, f^0\} \text{ from } p(m, b, f)$$

Gibbs Algorithm

Draw a sample state from proposed distribution

$$\{m^0, b^0, f^0\} \text{ from } p(m, b, f)$$

for t in $[1 : T]$:

$$m^t \sim p(m \mid b^{t-1}, f^{t-1})$$

$$b^t \sim p(b \mid m^t, f^{t-1})$$

$$f^t \sim p(f \mid m^t, b^t)$$

Gibbs Algorithm

Simplest of MCMC algorithms

Gibbs Algorithm

Simplest of MCMC algorithms

Use if conditional probabilities can be sampled

Gibbs Algorithm

Simplest of MCMC algorithms

Use if conditional probabilities can be sampled

Slow for correlated parameters


Metropolis-adjusted Langevin Rule

Similar to MH algorithm, with a target probability distribution function term

Metropolis-adjusted Langevin Rule

Similar to MH algorithm, with a target probability distribution function term

$$\alpha = \frac{p(x')}{p(x_t)}$$

MH 

Metropolis-adjusted Langevin Rule

Similar to MH algorithm, with a target probability distribution function term

$$\alpha = \frac{p(x') p(y | x')}{p(x_t) p(y | x_t)}$$

Hamiltonian Monte Carlo

MH's and Gibbs' (and a little bit of Langevin's) weakness is their reliance on random steps

Hamiltonian Monte Carlo

MH's and Gibbs' (and a little bit of Langevin's) weakness is their reliance on random steps

Add a momentum term to each variable

Hamiltonian Monte Carlo

MH's and Gibbs' (and a little bit of Langevin's) weakness is their reliance on random steps

Add a momentum term to each variable

Gives the expected distance and direction of jump of variable based on the last few jumps

Hamiltonian Monte Carlo

Draw a sample state from proposed distribution

$$\{m^0, b^0, f^0\} \text{ from } p(m, b, f) p(\mathbf{v})$$

Hamiltonian Monte Carlo

Draw a sample state from proposed distribution

$\{m^0, b^0, f^0\}$ from $p(m, b, f) p(\mathbf{v})$

$$H(m, b, f, \mathbf{v}) = E(m, b, f) + K(\mathbf{v}) = \text{constant}$$

Hamiltonian Monte Carlo

Draw a sample state from proposed distribution

$$\{m^0, b^0, f^0\} \text{ from } p(m, b, f) p(\mathbf{v})$$

$$H(m, b, f, \mathbf{v}) = E(m, b, f) + K(\mathbf{v}) = \text{constant}$$

...followed by same accept/reject steps as MH

Slice Sampling

Sample the region under proposed distribution

Slice Sampling

Sample the region under proposed distribution

1. Sample a random initial value x_0

Slice Sampling

Sample the region under proposed distribution

1. Sample a random initial value x_0
2. At each iteration i
 - a. Sample uniformly (vertical slice) for an auxiliary variable in the region $[0, p(x_{i-1} \mid y)]$
 - b. Sample uniformly (horizontal slice) for x_i

Summary

Metropolis-Hastings:	drunk man walking
Gibbs:	drunk man hopscotch
Langevin:	drunk man with a map
Hamiltonian:	drunk man running
Slice Sampling:	drunk man in a room

Summary

While easy-to-use, they are easy to mess up

There are a lot of resources online, especially for the most common types of algorithms

<https://chi-feng.github.io/mcmc-demo/app.html>

Prostate Cancer Research:

<https://www.cancer.gov/types/prostate/psa-fact-sheet>

<https://www.ncbi.nlm.nih.gov/pubmed/11333995>

<https://www.ncbi.nlm.nih.gov/pubmed/15163773>

<https://www.ncbi.nlm.nih.gov/pubmed/29772218>

<https://www.ices.on.ca/Publications/Atlases-and-Reports/2002/Prostate-specific-antigen-PSA-screening>

<http://www.topalbertadoctors.org/download/276/Prostate%2BCancer%2BGuideline%2BEval%2B%2526%2BReferr.pdf>

MH and Slice Sampling :

https://www.sheffield.ac.uk/polopoly_fs/1.60510!/file/MCMC.pdf

Gibbs Sampling:

<http://www.mit.edu/~ilkery/papers/GibbsSampling.pdf>

Metropolis-adjusted Langevin:

<https://warwick.ac.uk/fac/sci/statistics/crism/visitors/vats/misc/malavsrwm.pdf>

Hamiltonian Monte Carlo Reference:

<https://www.cs.utoronto.ca/~radford/ftp/ham-mcmc.pdf>

EXTRA SLIDES

Bayes' Theorem (once again)

LOG LIKELIHOOD FUNCTION

$$\ln p(y \mid x, \sigma, m, b, f) = -\frac{1}{2} \sum_n \left[\frac{(y_n - \text{logit}(x_n, m, b))^2}{s_n^2} + \ln(2\pi s_n^2) \right]$$

$$\text{logit}(x, m, b) = \frac{1}{1 + e^{-(mx+b)}}$$

$$s_n^2 = \sigma_n^2 + f^2 (mx_n + b)^2$$

Doctor's Visit

Doctor's Visit

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	
	-		

Doctor's Visit

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	
	-	0.1	

Doctor's Visit

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	0.2
	-	0.1	

Doctor's Visit

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	0.2
	-	0.1	0.8

Doctor's Visit

Test is positive!

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	0.2
	-	0.1	0.8

Doctor's Visit

Test is positive!

$$p(\text{cancer} \mid +)?$$

		Prostate Cancer	
		Yes	No
Test Result	+	0.9	0.2
	-	0.1	0.8

Doctor's Visit

$$p(\text{cancer} \mid +) = \frac{p(+ \mid \text{cancer}) p(\text{cancer})}{p(+)}$$

Doctor's Visit

$$p(\text{cancer} \mid +) = \frac{0.9 \cdot p(+ \mid \text{cancer}) \cdot p(\text{cancer})}{p(+)}$$

Doctor's Visit

$$p(\text{cancer} \mid +) = \frac{\overset{0.9}{p(+ \mid \text{cancer})} \overset{0.000001}{p(\text{cancer})}}{p(+)}$$

Doctor's Visit

$$p(+)=p(+\mid \text{cancer})\,p(\text{cancer})+p(+\mid \text{no cancer})\,p(\text{no cancer})$$

Doctor's Visit

0.00001

0.99999

$$p(+) = p(+ \mid \text{cancer}) p(\text{cancer}) + p(+ \mid \text{no cancer}) p(\text{no cancer})$$

0.9

0.2

Doctor's Visit

$$p(\text{cancer} \mid +) = 0.000045$$

Doctor's Visit

$$p(\text{cancer} \mid +) = 0.000045$$



Low chance of
prostate cancer
(0.00001)



High chance of
false positive
(0.2)

Doctor's Visit

What if the prior was larger?

Doctor's Visit

What if the prior was larger?

$$p(\text{cancer}) = 0.25$$

Doctor's Visit

What if the prior was larger?

$$p(\text{cancer}) = 0.25$$

$$p(\text{cancer} \mid +) \approx 0.58$$

