AE 504 Compressible Flow

Table of Contents

Vittorio	Baraldi	 1
Fv 1		- 1

Vittorio Baraldi

Homework 3 clear all close all clc

Ex. 1

```
Part c)
%initial data
Pres=3.79;
                   %reservoir pressure [atm]
Pe=1;
                   %exit pressure [atm]
At = 0.5;
                    %throat area [m^2]
Ae=3.5;
                    %exit area [m^2]
Me=0.31;
                   %exit mach no.
                   %heat air coefficient
q=1.4;
exitratio=Ae/At;
syms asol msol
A=2;
                    %initial shock position quess
                    %increment
options = optimoptions('fsolve','Display','none');
for i=1:1000
    %area-mach no. function
    fM=@(x) ((((2/(q+1))*(1+0.2*x^2))^6)/x^2)-(A/At)^2;
    %solving for Mach no. before shock
    M1=fsolve(fM, 2, options);
    %mach no. after shock
    M2=sqrt(((g-1)*M1^2+2)/(2*g*M1^2-(g-1)));
    %total pressure ratio across shock
    Pratio=((((g+1)*M1^2)/((g-1)*M1^2+2))^(g/(g-1)))*...
        (((g+1)/(2*g*M1^2-(g-1)))^(1/(g-1)));
    %Area ratio based on M2
    fA=@(x) ((((2/(g+1))*(1+0.2*M2^2))^6)/M2^2)-x^2;
    Aratio=fsolve(fA,1.1,options);
    %finding the new area ratio between exit area and the sonic area
    %condition after shock
    Ae_A2t=exitratio*(At/A)*Aratio;
```

```
%solving for the exit Mach no. given the area ratio above
    fM2=@(x) ((((2/(q+1))*(1+0.2*x^2))^6)/x^2)-Ae A2t^2;
    Me=fsolve(fM2,0.9,options);
    %with isentropic relations, find the conditions at the exit
    [Pexit,~,~,~,~,~]=isentropicflow(Me,g);
    %Pfinal is the exit pressure, given the initial guess
    Pfinal=Pexit*Pratio*Pres;
    %difference between computed exit pressure and actual exit
 pressure
    diff=abs(Pfinal-Pe);
    %checking on the difference tolerance (1e-3)
    if diff<1e-3</pre>
        sol=A;
        break
    else
        %if condition is not satisfied, guess is incremented
        A=A+dA;
    end
end
fprintf('Shock occurs at A = %1.2f meters squared', sol)
Shock occurs at A = 2.50 meters squared
```

Published with MATLAB® R2020a