

A Krylov Subspace Approach to Modeling of Wave Propagation in Open Domains

Delft University of Technology, CAS Group

Jörn Zimmerling

Monday 2nd February, 2015

Introduction - Motivation

- Fast and robust modeling of wave-propagation essential to many areas of science

Introduction - Motivation

- Fast and robust modeling of wave-propagation essential to many areas of science
- Acoustic/seismic waves: Earth formations for exploration physics
- Electromagnetic waves: Induction logging of borehole

Introduction - Motivation

- Fast and robust modeling of wave-propagation essential to many areas of science
- Acoustic/seismic waves: Earth formations for exploration physics
- Electromagnetic waves: Induction logging of borehole
- Electromagnetic waves: Light matter interaction on the nano-scale
 - Periodic gratings for process control (ASML)
 - Sensors based on resonance shifts in plasmonic resonators

Introduction - Motivation

- Fast and robust modeling of wave-propagation essential to many areas of science
- Acoustic/seismic waves: Earth formations for exploration physics
- Electromagnetic waves: Induction logging of borehole
- Electromagnetic waves: Light matter interaction on the nano-scale
 - Periodic gratings for process control (ASML)
 - Sensors based on resonance shifts in plasmonic resonators

Summary

For the forward and inverse problem fast approximate solvers for wave equations in open domains are required.

forward problem: fast, low memory inverse problem: low order

Introduction - Basic Idea

- Open wave equations can be expressed as

$$(A + I\partial_t)f = b, \quad (1)$$

with the sparse, system matrix A symmetric as $A^T W = W A$.
Direct evaluation of Eq.1 not possible as A ($\text{Order}(A) > 10^6$)

Introduction - Basic Idea

- Open wave equations can be expressed as

$$(A + I\partial_t)f = b, \quad (1)$$

with the sparse, system matrix A symmetric as $A^T W = W A$.
Direct evaluation of Eq.1 not possible as A ($\text{Order}(A) > 10^6$)

- Krylov approximation $A V_m \approx V_m H_m$ with $\text{Order}(A) \gg m$
- Approximate solution $f_m = V_m q$ via

$$(H_m + I\partial_t)q = V^T b \quad (2)$$

Introduction - Basic Idea

- Open wave equations can be expressed as

$$(A + I\partial_t)f = b, \quad (1)$$

with the sparse, system matrix A symmetric as $A^T W = W A$.
Direct evaluation of Eq.1 not possible as A ($\text{Order}(A) > 10^6$)

- Krylov approximation $A V_m \approx V_m H_m$ with $\text{Order}(A) \gg m$
- Approximate solution $f_m = V_m q$ via

$$(H_m + I\partial_t)q = V^T b \quad (2)$$

Motivation for ROM approach

- Discretization of wave problems are oversampled w.r.t. Nyquist
- Approximation of a few resonances can be sufficient (e.g. violin)
- Frequency independent ABC/PML enables single ROM

Outline

- 1 From wave equation to discrete operator
 - System formulation
 - Approximation of open system by PML
- 2 Krylov methods for reduced order modeling
 - Construction
 - Projected solution
- 3 Examples
 - 2D box Problem TD
 - 3D cylinder Problem FD
- 4 Conclusion

Outline

- 1 From wave equation to discrete operator
 - System formulation
 - Approximation of open system by PML
- 2 Krylov methods for reduced order modeling
 - Construction
 - Projected solution
- 3 Examples
 - 2D box Problem TD
 - 3D cylinder Problem FD
- 4 Conclusion

From wave equation to discrete operator - System formulation 1

- Example of wave equation: the Maxwell equations

$$-\nabla \times \mathbf{H} + \sigma \mathbf{E} + \varepsilon \partial_t \mathbf{E} = -\mathbf{J}^{\text{ext}}, \quad (3)$$

$$\nabla \times \mathbf{E} + \mu \partial_t \mathbf{H} = -\mathbf{K}^{\text{ext}}. \quad (4)$$

Solve for \mathbf{E} and \mathbf{H} and given sources \mathbf{J}^{ext} and \mathbf{K}^{ext} and a spatially varying medium characterized by $\sigma(\mathbf{x})$, $\varepsilon(\mathbf{x})$ and $\mu(\mathbf{x})$.

From wave equation to discrete operator - System formulation 1

- Example of wave equation: the Maxwell equations

$$-\nabla \times \mathbf{H} + \sigma \mathbf{E} + \varepsilon \partial_t \mathbf{E} = -\mathbf{J}^{\text{ext}}, \quad (3)$$

$$\nabla \times \mathbf{E} + \mu \partial_t \mathbf{H} = -\mathbf{K}^{\text{ext}}. \quad (4)$$

Solve for \mathbf{E} and \mathbf{H} and given sources \mathbf{J}^{ext} and \mathbf{K}^{ext} and a spatially varying medium characterized by $\sigma(\mathbf{x})$, $\varepsilon(\mathbf{x})$ and $\mu(\mathbf{x})$.

- General first order system formulation for all wave equations is

$$(\mathcal{D} + \mathcal{S} + \mathcal{M} \partial_t) \mathcal{F} = \mathcal{Q}'. \quad (5)$$

Differentiation & medium matricees and field & source vector as

$$\mathcal{D} = \begin{bmatrix} 0 & -\nabla \times \\ \nabla \times & 0 \end{bmatrix}, \quad \mathcal{S} = \begin{bmatrix} \sigma & 0 \\ 0 & 0 \end{bmatrix}, \quad \mathcal{M} = \begin{bmatrix} \varepsilon & 0 \\ 0 & \mu \end{bmatrix}, \quad (6)$$

$$\mathcal{F} = [\mathbf{E}, \mathbf{H}]^T, \text{ and } \mathcal{Q}' = -[\mathbf{J}^{\text{ext}}, \mathbf{K}^{\text{ext}}]^T. \quad (7)$$

From wave equation to discrete operator - System formulation 2

- After discretization the Matrix equation

$$(D + S + M\partial_t)f = q'. \quad (8)$$

describes the wave equation.

From wave equation to discrete operator - System formulation 2

- After discretization the Matrix equation

$$(D + S + M\partial_t)f = q'. \quad (8)$$

describes the wave equation.

- We define $A = (M^{-1}D + M^{-1}S)$ and $q'(x, t) = q(x)w(t)$,

$$(A + I\partial_t)f = M^{-1}q(x)w(t), \quad (9)$$

From wave equation to discrete operator - System formulation 2

- After discretization the Matrix equation

$$(D + S + M\partial_t)f = q'. \quad (8)$$

describes the wave equation.

- We define $A = (M^{-1}D + M^{-1}S)$ and $q'(x, t) = q(x)w(t)$,

$$(A + I\partial_t)f = M^{-1}q(x)w(t), \quad (9)$$


- Equation 9 is solved by

$$f = w(t) * \eta(t) \exp(-At)M^{-1}q(x), \quad (10)$$



with $\eta(t)$ the Heaviside step function. The Laplace image is defined as

$$\hat{f} = \hat{w}(s)(A + sI)^{-1}M^{-1}q. \quad (11)$$

From wave equation to discrete operator - Approximation of open system by PML






From wave equation to discrete operator - Approximation of open system by PML






- Truncation of computational domain necessary

From wave equation to discrete operator - Approximation of open system by PML



- Truncation of computational domain necessary
- Perfectly matched layers (PML) absorb outgoing waves reflection less

From wave equation to discrete operator - Approximation of open system by PML



- Truncation of computational domain necessary
- Perfectly matched layers (PML) absorb outgoing waves reflection less
- Reformulation of PML to complex coordinate stretching
- Optimal frequency independent PML matched over spectral region of interest

From wave equation to discrete operator - Approximation of open system by PML

- After complex coordinate stretching A approximates outgoing waves and incoming waves

From wave equation to discrete operator - Approximation of open system by PML

- After complex coordinate stretching A approximates outgoing waves and incoming waves
 - $\Re(\lambda) \geq 0$ contributes to the causal solution $t \geq 0$ (outgoing)
 - $\Re(\lambda) \leq 0$ contributes to the anticausal solution $t \leq 0$ (incoming)

From wave equation to discrete operator - Approximation of open system by PML

- After complex coordinate stretching A approximates outgoing waves and incoming waves
 - $\Re(\lambda) \geq 0$ contributes to the causal solution $t \geq 0$ (outgoing)
 - $\Re(\lambda) \leq 0$ contributes to the anticausal solution $t \leq 0$ (incoming)
- Apply stability correction/radiation condition by using $\eta(A)$ as

$$f = w(t) * \eta(t) \Re \left[\eta(A) \exp(-At) M^{-1} q \right], \quad (12)$$

ensures causality.

From wave equation to discrete operator - Approximation of open system by PML

- After complex coordinate stretching A approximates outgoing waves and incoming waves
 - $\Re(\lambda) \geq 0$ contributes to the causal solution $t \geq 0$ (outgoing)
 - $\Re(\lambda) \leq 0$ contributes to the anticausal solution $t \leq 0$ (incoming)
- Apply stability correction/radiation condition by using $\eta(A)$ as

$$f = w(t) * \eta(t) \Re \left[\eta(A) \exp(-At) M^{-1} q \right], \quad (12)$$

ensures causality.

- Numerical Greens function
- Symmetry condition $A^T W = W A$ enables fast computation of Krylov basis

Outline

- 1 From wave equation to discrete operator
 - System formulation
 - Approximation of open system by PML
- 2 Krylov methods for reduced order modeling
 - Construction
 - Projected solution
- 3 Examples
 - 2D box Problem TD
 - 3D cylinder Problem FD
- 4 Conclusion

Krylov methods for ROM - Construction

- Polynomial, Extended and Rational Krylov subspace:

$$\mathcal{K}_m^{\text{PKS}} = \text{span}\{b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (13)$$

$$\mathcal{K}_{n;m}^{\text{EKS}} = \text{span}\{A^{-n+1}b, \dots, A^{-1}b, b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (14)$$

$$\mathcal{K}_m^{\text{RKS}} = \text{span}\{(A + \sigma_1 I)^{-1}b, \dots, \prod_{i=1}^m (A + \sigma_i I)^{-1}b\}. \quad (15)$$

Krylov methods for ROM - Construction

- Polynomial, Extended and Rational Krylov subspace:

$$\mathcal{K}_m^{\text{PKS}} = \text{span}\{b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (13)$$

$$\mathcal{K}_{n;m}^{\text{EKS}} = \text{span}\{A^{-n+1}b, \dots, A^{-1}b, b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (14)$$

$$\mathcal{K}_m^{\text{RKS}} = \text{span}\{(A + \sigma_1 I)^{-1}b, \dots, \prod_{i=1}^m (A + \sigma_i I)^{-1}b\}. \quad (15)$$

- In the case of a symmetry in A
 - PKS can be constructed via 3-term Lanczos algorithm
 - EKS can be constructed via 5-term Lanczos algorithm

Krylov methods for ROM - Construction

- Polynomial, Extended and Rational Krylov subspace:

$$\mathcal{K}_m^{\text{PKS}} = \text{span}\{b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (13)$$

$$\mathcal{K}_{n;m}^{\text{EKS}} = \text{span}\{A^{-n+1}b, \dots, A^{-1}b, b, Ab, A^2b, \dots, A^{m-1}b\}, \quad (14)$$

$$\mathcal{K}_m^{\text{RKS}} = \text{span}\{(A + \sigma_1 I)^{-1}b, \dots, \prod_{i=1}^m (A + \sigma_i I)^{-1}b\}. \quad (15)$$

- In the case of a symmetry in A
 - PKS can be constructed via 3-term Lanczos algorithm
 - EKS can be constructed via 5-term Lanczos algorithm
- RKS can be constructed via rational Arnoldi algorithm
Selection of shifts allow the adaption of the spectrum of H_m towards A

Krylov methods for ROM - Projected solution

- The projected solution of our wave equation is given by

$$f_m(t) = w(t) * \eta(t) \Re e[V_m \eta(H_m) \exp(-H_m t) V_m^T M^{-1} q] \quad (16)$$

with Laplace image

$$\hat{f}_m(s) = \hat{w}(s) [V_m (H_m + sI)^{-1} V_m^T + V_m^\dagger (H_m^\dagger + sI)^{-1} V_m^*] M^{-1} q \quad (17)$$

Krylov methods for ROM - Projected solution

- The projected solution of our wave equation is given by

$$f_m(t) = w(t) * \eta(t) \Re[\mathbf{V}_m \eta(\mathbf{H}_m) \exp(-\mathbf{H}_m t) \mathbf{V}_m^T \mathbf{M}^{-1} \mathbf{q}] \quad (16)$$

with Laplace image

$$\hat{f}_m(s) = \hat{w}(s) [\mathbf{V}_m (\mathbf{H}_m + sI)^{-1} \mathbf{V}_m^T + \mathbf{V}_m^\dagger (\mathbf{H}_m^\dagger + sI)^{-1} \mathbf{V}_m^*] \mathbf{M}^{-1} \mathbf{q} \quad (17)$$

- Let (s_j, λ_j) be an eigenpair of \mathbf{H}_m and satisfy $\Re(\lambda_j) > 0$
- A resonance is given by $f_{res} = \mathbf{V}_m s_j$ with complex frequency λ_j

Krylov methods for ROM - Projected solution

- The projected solution of our wave equation is given by

$$f_m(t) = w(t) * \eta(t) \Re[\mathbf{V}_m \eta(\mathbf{H}_m) \exp(-\mathbf{H}_m t) \mathbf{V}_m^T \mathbf{M}^{-1} \mathbf{q}] \quad (16)$$

with Laplace image

$$\hat{f}_m(s) = \hat{w}(s) [\mathbf{V}_m (\mathbf{H}_m + sI)^{-1} \mathbf{V}_m^T + \mathbf{V}_m^\dagger (\mathbf{H}_m^\dagger + sI)^{-1} \mathbf{V}_m^*] \mathbf{M}^{-1} \mathbf{q} \quad (17)$$

- Let (s_j, λ_j) be an eigenpair of \mathbf{H}_m and satisfy $\Re(\lambda_j) > 0$
- A resonance is given by $f_{res} = \mathbf{V}_m s_j$ with complex frequency λ_j
 - $\Re(\lambda_j)$ associated with losses (material and radiation)
 - $\Im(\lambda_j)$ frequency of resonance

Krylov methods for ROM - Projected solution

- The projected solution of our wave equation is given by

$$f_m(t) = w(t) * \eta(t) \Re[\mathbf{V}_m \eta(\mathbf{H}_m) \exp(-\mathbf{H}_m t) \mathbf{V}_m^T \mathbf{M}^{-1} \mathbf{q}] \quad (16)$$

with Laplace image

$$\hat{f}_m(s) = \hat{w}(s) [\mathbf{V}_m (\mathbf{H}_m + sI)^{-1} \mathbf{V}_m^T + \mathbf{V}_m^\dagger (\mathbf{H}_m^\dagger + sI)^{-1} \mathbf{V}_m^*] \mathbf{M}^{-1} \mathbf{q} \quad (17)$$

- Let (s_j, λ_j) be an eigenpair of \mathbf{H}_m and satisfy $\Re(\lambda_j) > 0$
- A resonance is given by $f_{res} = \mathbf{V}_m s_j$ with complex frequency λ_j
 - $\Re(\lambda_j)$ associated with losses (material and radiation)
 - $\Im(\lambda_j)$ frequency of resonance


Conclusion theoretical part

- Single ROM describes dynamics of A over whole frequency band
- After Krylov decomposition frequency sweeps possible in Eq. 17
- Evaluation of solution comes at negligible cost

Outline

- 1 From wave equation to discrete operator
 - System formulation
 - Approximation of open system by PML
- 2 Krylov methods for reduced order modeling
 - Construction
 - Projected solution
- 3 Examples
 - 2D box Problem TD
 - 3D cylinder Problem FD
- 4 Conclusion



Examples - 2D box Problem



Configuration

- Dielectric box with contrast $c_0/c = 2$
- H-polarized 2D
- Derivative Gaussian pulse with $\lambda_{sc}^{max} = 94L$
- Small system of order $n = 36k$



Examples - 2D box Problem - Results



PKS Result

- Output compared with FDTD (time stepping)
- PKS converges within $m=1500$, RKS with $m=48$
- ROM allows convolution with arbitrary input wavelet



Examples - 2D box Problem - Ritz values



Resonance expansion


- Output can be expanded in just 26 approximate eigenpairs of A
- On time interval $[T_{min}, \infty)$ contributing Ritz values in $\Lambda^{\text{res}} := \{z \in \mathbb{C} : 0 \leq \text{Re}(z) < \sigma_{\max}, \omega_{\min} < -\text{Im}(z) < \omega_{\max}\}$
- Error of approximation $O[\exp -\sigma_{\max} T_{min}]$

Examples - 2D box Problem - Ritz vectors



Examples - 2D box Problem - Resonance


Examples - 3D box nanorod



PKS Results

- Spontaneous decay rate: quantum emitter close to resonator
- Electrical field at the source


Examples - 3D box nanorod



PKS Results

- Spontaneous decay rate: quantum emitter close to resonator
- Electrical field at the source
- Full 3D-Maxwell with second order dispersion relation
- $\beta_3 \partial_t^2 \tilde{\mathbf{P}} + \beta_2 \partial_t \tilde{\mathbf{P}} + \beta_1 \tilde{\mathbf{P}} = \beta_0 \mathbf{E}$


Examples - 3D box nanorod



PKS Results

- Spontaneous decay rate: quantum emitter close to resonator
- Electrical field at the source
- Full 3D-Maxwell with second order dispersion relation
- $\beta_3 \partial_t^2 \tilde{\mathbf{P}} + \beta_2 \partial_t \tilde{\mathbf{P}} + \beta_1 \tilde{\mathbf{P}} = \beta_0 \mathbf{E}$
- After discretization $n = 8.5 \cdot 10^6$


Examples - 3D box nanorod



Configuration

- Reduction to $m = 4500$
- $n/m = 2000$



Examples - 3D box nanorod



Configuration

- Reduction to $m = 4500$
- $n/m = 2000$
- Computation of ROM takes 45 min
- Evaluation of ROM at 100 frequencies takes 10 sec

Examples - 3D box nanorod



Configuration

- Reduction to $m = 4500$
- $n/m = 2000$
- Computation of ROM takes 45 min
- Evaluation of ROM at 100 frequencies takes 10 sec

Outline

- ① From wave equation to discrete operator
 - System formulation
 - Approximation of open system by PML
- ② Krylov methods for reduced order modeling
 - Construction
 - Projected solution
- ③ Examples
 - 2D box Problem TD
 - 3D cylinder Problem FD
- ④ Conclusion

Conclusion

- Krylov ROM can efficiently approximate wave propagation in open domains

Conclusion

- Krylov ROM can efficiently approximate wave propagation in open domains
- Significantly outperforms conventional methods:
 - TD: large time intervals, multiple wavelets
 - FD: multiple frequency solutions are needed

	PKS	EKS	RKS
Reduction n/m	low	medium	high

Conclusion

- Krylov ROM can efficiently approximate wave propagation in open domains
- Significantly outperforms conventional methods:
 - TD: large time intervals, multiple wavelets
 - FD: multiple frequency solutions are needed

	PKS	EKS	RKS
Reduction n/m	low	medium	high
Iteration Cost	low	medium	high

Conclusion

- Krylov ROM can efficiently approximate wave propagation in open domains
- Significantly outperforms conventional methods:
 - TD: large time intervals, multiple wavelets
 - FD: multiple frequency solutions are needed

	PKS	EKS	RKS
Reduction n/m	low	medium	high
Iteration Cost	low	medium	high
Memory Demand	$3n$	$5n$	$m \cdot n$

A Krylov Subspace Approach to Modeling of Wave Propagation in Open Domains

Delft University of Technology, CAS Group

Jörn Zimmerling

Monday 2nd February, 2015