

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero

asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain



Show & Tell Karumi

February 1st 2019



[https://github.com/asuncionjc/
Show_and_Tell_Mathematical_Proofs](https://github.com/asuncionjc/Show_and_Tell_Mathematical_Proofs)



[https://github.com/asuncionjc/
Show_and_Tell_Mathematical_Proofs](https://github.com/asuncionjc/Show_and_Tell_Mathematical_Proofs)



Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

A **theorem** is a statement that can be shown to be true.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- **Hypothesis:**
- **Thesis:**

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- **Hypothesis:**
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- **Thesis:**

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- **Hypothesis:**
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- **Thesis:**
 - $\exists c \in (a, b) : f(c) = 0$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ① $a = b$
- ② $a^2 = ab$
- ③ $a^2 - b^2 = ab - b^2$
- ④ $(a - b)(a + b) = b(a - b)$
- ⑤ $a + b = b$
- ⑥ $2b = b$
- ⑦ $2 = 1$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ❶ $a = b$
- ❷ $a^2 = ab$
- ❸ $a^2 - b^2 = ab - b^2$
- ❹ $(a - b)(a + b) = b(a - b)$
- ❺ $a + b = b$
- ❻ $2b = b$
- ❼ $2 = 1$

Where is the error?

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

Example

Theorem: $\sqrt{2}$ is irrational.

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

Example

Theorem: $\sqrt{2}$ is irrational.

Previous concepts

- **Definition:** A real number r is *rational* if there exists integers p and q , with no common factors and $q \neq 0$ such that $r = p/q$. A real number that is not rational is called *irrational*.
- **Theorem:** Let a be an integer number such that a^2 is even. Then a is also even.

Example

Theorem: $\sqrt{2}$ is irrational.

Example

Theorem: $\sqrt{2}$ is irrational.

- 1 Assume $\sqrt{2}$ is rational.

Example

Theorem: $\sqrt{2}$ is irrational.

- 1 Assume $\sqrt{2}$ is rational.
- 2 $\sqrt{2} = \frac{p}{q}$, p and q without common factors.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Contradiction!

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Contradiction!

On the one hand, we assumed p and q have no common factors. On the other hand, if p and q are even, 2 is a common factor. Therefore, $\sqrt{2}$ is irrational.

It's your turn!

Theorem: Let a be an integer number such that a^2 is even. Then a is also even.

It's your turn!

Theorem: Let a be an integer number such that a^2 is even. Then a is also even.

- ① Assume a is odd.
- ② $a = 2k + 1$ for some integer k .
- ③ $a^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$.
- ④ a^2 is odd.

Contradiction!

Since, by hypothesis, a^2 is even.

Therefore, **a is even.**

Another example?

Theorem: For every natural number $a > 2$ and a prime, then it holds that a is odd.

Another example?

Theorem: For every natural number $a > 2$ and a prime, then it holds that a is odd.

- ① Assume a is even.
- ② $a = 2k$, for some k . Since $a > 2$, then $k \neq 1$.
- ③ a is composite.

Contradiction!

Since, by hypothesis, a is prime.

Therefore, **a is odd.**

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction**
- 4 Conclusions

- ① Base case ($n = 1$).
- ② Assume the result for all $k < n$ (Induction hypothesis).
- ③ Prove for n .

- ① Base case ($n = 1$).
- ② Assume the result for all $k < n$ (Induction hypothesis).
- ③ Prove for n .

Example

Theorem (Gauss): For all $n \in \mathbb{N}$:

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

- 1 Base case ($n = 1$).
- 2 Assume the result for all $k < n$ (Induction hypothesis).
- 3 Prove for n .

Example

Theorem (Gauss): For all $n \in \mathbb{N}$:

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

1 $(n = 1) \sum_{i=1}^1 i = 1 = \frac{1(1+1)}{2}.$

- 1 Base case ($n = 1$).
- 2 Assume the result for all $k < n$ (Induction hypothesis).
- 3 Prove for n .

Example

Theorem (Gauss): For all $n \in \mathbb{N}$:

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

- 1 ($n = 1$) $\sum_{i=1}^1 i = 1 = \frac{1(1+1)}{2}$.
- 2 (Induction hypothesis) The result is true for all $k < n$.

- ❶ Base case ($n = 1$).
- ❷ Assume the result for all $k < n$ (Induction hypothesis).
- ❸ Prove for n .

Example

Theorem (Gauss): For all $n \in \mathbb{N}$:

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

- ❶ $(n = 1) \sum_{i=1}^1 i = 1 = \frac{1(1+1)}{2}$.
- ❷ (Induction hypothesis) The result is true for all $k < n$.
- ❸
$$\begin{aligned} \sum_{i=1}^n i &= \left(\sum_{i=1}^{n-1} i \right) + n = \frac{(n-1)(n-1+1)}{2} + n = \frac{(n-1)n}{2} + \frac{2n}{2} = \\ &= \frac{n^2 - n + 2n}{2} = \frac{n^2 + n}{2} = \frac{n(n+1)}{2}. \end{aligned}$$

Your turn!

Theorem (Geometric series): For all $n \in \mathbb{N}$ and $r \neq 1$:

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r}$$

Your turn!

Theorem (Geometric series): For all $n \in \mathbb{N}$ and $r \neq 1$:

$$\sum_{i=0}^n r^i = \frac{1 - r^{n+1}}{1 - r}$$

- ① $(n = 0) : \sum_{i=0}^0 r^i = r^0 = 1 = \frac{1-r}{1-r}.$
- ② (Induction hypothesis) The result is true for all $k < n$.
- ③
$$\begin{aligned} \sum_{i=0}^n r^i &= \sum_{i=0}^{n-1} r^i + r^n = \frac{1-r^n}{1-r} + r^n = \frac{1-r^n+r^n(1-r)}{1-r} = \\ &= \frac{1-r^n+r^n-r^{n+1}}{1-r} = \frac{1-r^{n+1}}{1-r}. \end{aligned}$$

Last example

Theorem: For all $n \in \mathbb{N}$:

$11^n - 6$ is divisible by 5

Last example

Theorem: For all $n \in \mathbb{N}$:

$11^n - 6$ is divisible by 5

- ① $(n = 1) : 11^1 - 6 = 5$ which is divisible by 5.
- ② (Induction hypothesis) The result is true for all $k < n$, and therefore $11^k = 5m + 6$, for some integer m .
- ③ $11^n - 6 = (11 \cdot 11^{n-1}) - 6 = 11 \cdot (5m + 6) - 6 = 11 \cdot 5m + 66 - 6 = 11 \cdot 5m + 60 = 5 \cdot (11m + 12)$, which is divisible by 5.

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

- Some basic concepts about the mathematical results and how to prove them.
- Proof by contradiction.
- Proof by induction.

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero

asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain

Thank you for your attention!



Show & Tell Karumi

February 1st 2019