

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero
asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain



Show & Tell Karumi

XX XXXX 2019

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

A **theorem** is a statement that can be shown to be true.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:

- Thesis:

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- Thesis:

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- Thesis:
 - $\exists c \in (a, b) : f(c) = 0$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ① $a = b$
- ② $a^2 = ab$
- ③ $a^2 - b^2 = ab - b^2$
- ④ $(a - b)(a + b) = b(a - b)$
- ⑤ $a + b = b$
- ⑥ $2b = b$
- ⑦ $2 = 1$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ❶ $a = b$
- ❷ $a^2 = ab$
- ❸ $a^2 - b^2 = ab - b^2$
- ❹ $(a - b)(a + b) = b(a - b)$
- ❺ $a + b = b$
- ❻ $2b = b$
- ❼ $2 = 1$

Where is the error?

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction**
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero
asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain

Thank you!



Show & Tell Karumi

XX XXXX 2019