

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero
asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain



Show & Tell Karumi

XX XXXX 2019

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

A **theorem** is a statement that can be shown to be true.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:

- Thesis:

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- Thesis:

A **theorem** is a statement that can be shown to be true.

- Hypothesis.
- Thesis.

Example

Bolzano's theorem: If f is a continuous function defined on a closed interval $[a, b]$ such that $\text{sign}(f(a)) \neq \text{sign}(f(b))$.

Then there exists a point c in the open interval (a, b) satisfying $f(c) = 0$.

- Hypothesis:
 - $f : [a, b] \rightarrow \mathbb{R}$ is continuous.
 - $\text{sign}(f(a)) \neq \text{sign}(f(b))$
- Thesis:
 - $\exists c \in (a, b) : f(c) = 0$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ① $a = b$
- ② $a^2 = ab$
- ③ $a^2 - b^2 = ab - b^2$
- ④ $(a - b)(a + b) = b(a - b)$
- ⑤ $a + b = b$
- ⑥ $2b = b$
- ⑦ $2 = 1$

A **proof** is a chain of logical deductions that establishes the truth of a theorem.

- Hypothesis.
- Axioms (*Ex: A number is equal to itself, $a = a$*).
- Previous mathematical results.

Mistakes in proofs

Conjecture: $1 = 2$.

Assume a, b are two equal positive integers:

- ❶ $a = b$
- ❷ $a^2 = ab$
- ❸ $a^2 - b^2 = ab - b^2$
- ❹ $(a - b)(a + b) = b(a - b)$
- ❺ $a + b = b$
- ❻ $2b = b$
- ❼ $2 = 1$

Where is the error?

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

Example

Theorem: $\sqrt{2}$ is irrational.

- ① Assume the thesis is false.
- ② Show that this assumption leads to a contradiction.
- ③ Hence, the thesis is true.

Example

Theorem: $\sqrt{2}$ is irrational.

Previous concepts

- **Definition:** A real number r is *rational* if there exists integers p and q , with no common factors and $q \neq 0$ such that $r = p/q$. A real number that is not rational is called *irrational*.
- **Theorem:** Let a be an integer number such that a^2 is even. Then a is also even.

Example

Theorem: $\sqrt{2}$ is irrational.

Example

Theorem: $\sqrt{2}$ is irrational.

- 1 Assume $\sqrt{2}$ is rational.

Example

Theorem: $\sqrt{2}$ is irrational.

- ① Assume $\sqrt{2}$ is rational.
- ② $\sqrt{2} = \frac{p}{q}$, p and q without common factors.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Contradiction!

Example

Theorem: $\sqrt{2}$ is irrational.

- ❶ Assume $\sqrt{2}$ is rational.
- ❷ $\sqrt{2} = \frac{p}{q}$, p and q without common factors.
- ❸ $2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$.
- ❹ p^2 is even and, by the previous theorem, p is also even.
- ❺ $p = 2s$, for some integer s .
- ❻ $(2s)^2 = 2q^2 \Rightarrow 4s^2 = 2q^2 \Rightarrow 2s^2 = q^2$.
- ❼ q^2 is even, and so it is q .

Contradiction!

On the one hand, we assumed p and q have no common factors. On the other hand, if p and q are even, 2 is a common factor. Therefore, $\sqrt{2}$ is irrational.

It's your turn!

Theorem: Let a be an integer number such that a^2 is even. Then a is also even.

It's your turn!

Theorem: Let a be an integer number such that a^2 is even. Then a is also even.

- ① Assume a is odd.
- ② $a = 2k + 1$ for some integer k .
- ③ $a^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$.
- ④ a^2 is odd.

Contradiction!

Since, by hypothesis, a^2 is even.

Therefore, a is even.

Another example?

Theorem: For every natural number $a > 2$ and a prime, then it holds that a is odd.

Another example?

Theorem: For every natural number $a > 2$ and a prime, then it holds that a is odd.

- ① Assume a is even.
- ② $a = 2k$, for some k . Since $a > 2$, then $k \neq 1$.
- ③ a is composite.

Contradiction!

Since, by hypothesis, a is prime.

Therefore, a is odd.

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction**
- 4 Conclusions

Outline

- 1 What is a proof?
- 2 Proof by contradiction
- 3 Proof by induction
- 4 Conclusions

An Introduction to Mathematical Reasoning with Applications to Induction and Contradiction Proofs

M^a Asunción Jiménez Cordero
asuncionjc@us.es

Departamento de Estadística e Investigación Operativa
and Instituto de Matemáticas de la Universidad de Sevilla,
Sevilla, Spain

Thank you!



Show & Tell Karumi

XX XXXX 2019