

Bayesian Inference of Star Formation History in the Host Galaxies of Tidal Disruption Events

Alexander S. Wheaton

Supervisor: Andy Lawrence, FRSE



March 30, 2021

Project Aims




Figure: AT2019qiz Host Galaxy

- Investigate the statistical behaviour of the star formation in the host galaxies of tidal disruption events.

Motivation: XSHOOTER Targets

| TDE | m | z |
|-------------|-----|----------|
| ASASSN-14li | 15 | 0.0206 |
| ASASSN-15oi | 16 | 0.0484 |
| AT2018fyk | 17 | 0.06 |
| AT2019ahk | 17 | 0.026211 |
| AT2019azh | 15 | 0.022 |
| AT2019dsg | 15 | 0.0512 |
| AT2019qiz | 15 | 0.01513 |
| iPTF16fni | 16 | 0.018 |
| AT2018hyz | 17 | 0.04573 |
| ASASSN-14ae | 17 | 0.0436 |

Table: The XSHOOTER targets on tidal disruption events.

Active Galactic Nuclei (AGN)

- ▶ Compact, highly luminous nuclear region.
- ▶ Broad spectral energy distribution.
- ▶ Strong (and sometimes broad) emission lines.
- ▶ Variability over short timescales.

Active Galactic Nuclei (AGN)




Figure: Accretion of matter onto surface of a black hole. Image adapted from Urry and Padovani 1995.

The Starburst-AGN Connection




Figure: Strong relationship between star formation and black hole mass.
Image adapted from Veilleux 2008.

- Cosmologically important impact on galaxy formation and evolution.

Possible mechanisms?

- ▶ Gas or radiation pressure from starbursts and/or AGN-driven wind shuts off black hole fuel supply.
- ▶ Direction of causation unclear.
- ▶ AGN is a possible source of quenching.
- ▶ Need more information...

Tidal Disruption Events



Figure: TDE impression, image credit NASA/JPL-Caltech.

- Do TDEs exhibit the same statistical behaviour?

The BAGPIPEs Module

Bayesian Analysis of Galaxies for Physical Inference and Parameter EStimation.

- ▶ Simulation of galactic spectra from SFH.
- ▶ Fit real spectra to plausible SFH.

The BAGPIEs Module




Figure: Simulated SFH and spectrum.

The BAGPIPES Module




Figure: Observed and fitted spectrum, with inferred SFH.

Stellar Population Dating

| Mass (solar masses) | Time (years) | Spectral type |
|---------------------|----------------|---------------|
| 60 | 3 million | O3 |
| 30 | 11 million | O7 |
| 10 | 32 million | B4 |
| 3 | 370 million | A5 |
| 1.5 | 3 billion | F5 |
| 1 | 10 billion | G2 (Sun) |
| 0.1 | 1000s billions | M7 |

Figure: Stellar lifetimes and spectral type.

- Poor temporal resolution in SFR beyond 1 billion years.

Stellar Population Dating

TABLE 2
Number density in the solar neighbourhood brighter than absolute magnitude +16 by
spectral type and class, per 10,000 pc³

| Class | Spectral type | | | | | | | Totals |
|---------------|---------------|----|-----|-----|-----|-----|------|--------|
| | O | B | A | F | G | K | M | |
| Giants | | | | 0.5 | 1.6 | 4 | 0.25 | 6.3 |
| Main sequence | 0.00025 | 1 | 5 | 25 | 63 | 100 | 630 | 800 |
| White dwarfs | | 63 | 100 | 50 | 50 | 25 | | 250 |

Figure: Number density of spectral types. Table adapted from Glenn 2001.

- Poor temporal resolution beyond 1 billion years.

Stellar Population Dating

- ▶ Solution? Parameterise metallicity as well.
- ▶ How do Lyman and Balmer series lines evolve with starburst age?
- ▶ What about other metal lines?




Figure: Variance in flux, over starburst evolved from $z=2.2$ to $z=0.01$.




Figure: Deltas in flux, over starburst evolved from $z=2.2$ to $z=0.01$.

SFH Inference - What is possible?

- ▶ Blind fitting of spectra with known priors.
- ▶ Fitting multiple SFH forms.
- ▶ Iterative fitting.

Exploring the SFH Parameter Space

R1: Entire Parameter Space with Two Components




Figure: Example SFH.

- ▶ Constrain z to be less than or equal to the TDE targets
- ▶ Assume separate bulk formation and burst events

Exploring the SFH Parameter Space

R2 & R3: Iterative Fitting




Figure: Example SFH.

- Insist burst component is younger than the bulk formation

Exploring the SFH Parameter Space




Figure: Prior and posterior SFH.

- “Best” fits favour single component star formation.

Selecting Physically Reasonable Priors




Figure: Example SFH.

- ▶ SPS limits burst detection.
- ▶ Precise form of old formation is unimportant.

Selecting Physically Reasonable Priors

R4: Fixed Old Component, Free Burst Component




Figure: Example SFH.

Conclusion:

- ▶ Impose a lower limit on bulk formation age and mass.
- ▶ Impose an upper limit on burst age, but let mass vary freely.

Selecting Physically Reasonable Priors

R4: Fixed Old Component, Free Burst Component

Other refined priors:

- ▶ Velocity dispersion, σ_v
- ▶ Birth cloud lifetime, τ_{BC}
- ▶ Metallicity, Z
- ▶ Extinction, A_V
- ▶ Nebular ionisation, U

Blind Fitting Results with R4 Priors




Figure: Prior and posterior SFH.

Blind Fitting Results with R4 Priors




Figure: Prior and posterior SFH.

Blind Fitting Results with R4 Priors




Figure: Prior and posterior SFH.

Blind Fitting Results with R4 Priors




Figure: Prior and posterior SFH.

Blind Fitting Results with R4 Priors





Figure: Prior and posterior SFH.

Correlated Parameters



Application to XSHOOTER Data




Figure: Posterior SFH for AT2019ahk.

Application to XSHOOTER Data




Figure: Posterior SFH for AT2019azh.

Application to XSHOOTER Data




Figure: Posterior SFH for iPTF16fnl.

Application to XSHOOTER Data




Figure: Posterior SFH for ASASSN14li.

Application to XSHOOTER Data




Figure: Posterior SFH for ASASSN15oi.

Application to XSHOOTER Data




Figure: Posterior SFH for AT2019dsg.

Conclusions



Figure: TDE impression, image credit NASA/JPL-Caltech.

- ▶ Starbusts are detectable **IF** recent
- ▶ Assumptions about host SFH history must be made
- ▶ TDE hosts exhibit strong starburst features

Future Applications

- ▶ Larger sample size.
- ▶ Tailor priors to host morphology.
- ▶ Compare with Hubble analogues, and Seyferts.