

Exercises for Tutorial 4

Gavin Salam, Edinburgh Jet Lectures
Tutors: Enrico Bothmann, Leonardo Vernazza

May/June 2017

1 Top reconstruction

Top quarks and their decays are widely studied at the LHC. They are of intrinsic interest, as the heaviest particle in the Standard Model, they appear in Higgs physics studies, as decay products of new particles in beyond-standard-model searches, and also as a background for a range of processes. Top quarks decay to a b -quark and a W and the W may decay either to a lepton and a neutrino, or to two quarks. In this tutorial, we consider the case where the W^+ decays to hadrons and the W^- decays to $\mu\bar{\nu}_\mu$. How many jets do you expect to see at “Born” level (i.e. leading order)?

If you cloned the tutorials previously, now do

```
git stash
git pull
```

and then go to the directory `tutorial-4/code`. Otherwise download a fresh zip file, unpack it and edit the `Makefile` in the above directory so that the path to Pythia points to where you built it previously.

The example here (yet again called `main01`!) illustrates many of the issues of a practical analysis using jets, including questions such as the choice of jet radius, transverse-momentum cuts, etc.

Start by reading through the code to see if you can understand what it does. Some helper code is in the other files in the directory and for a first round you shouldn't need to look into those.

By default the code generates tree-level events: i.e. in Pythia, all radiation, hadronisation and multiple-interaction steps have been turned off. Build and run the code and then generate a few plots

```
make main01
./main01 -R 0.1 -ptmin 2.0 -ymax 5.0
gnuplot all-plots.gp
```

The options specify a small jet radius, and extremely loose (unrealistic) acceptance cuts. The gnuplot command should generate a PDF file `all-plots.pdf` (but you may need to edit the terminal and replace it e.g. with “pdf” or “postscript”, depending on how gnuplot was build). Open the file to see what’s in there (it has four pages). By default 1000 events were generated. What fraction was well reconstructed, i.e. had the expected number of jets?

Now try running with a normal jet radius and acceptance cuts

```
make main01
./main01 -R 0.4 -ptmin 20.0 -ymax 2.5
gnuplot all-plots.gp
```

What fraction has the expected number of jets now?

To be continued.