



Linux : Getting feet wet

Bridge Course 2019

EE Dept IITB

24 July 2019

Outline

- 1 Introduction
 - What is Linux
 - Why Linux
- 2 Some L^AT_EX Examples
 - Tables and Figures
 - Mathematics

Linux is a family of open source Unix-like operating systems based on the Linux kernel,
Typically packaged in a Linux distribution




Why Linux




Why Linux

- 1 Access to hardware - almost no restrictions
- 2 Opensource - view code , modify , learn create


Why Linux

- 
- 1 Access to hardware - almost no restrictions
 - 2 Opensource - view code , modify , learn create
 - 3 Tailor Made - servers , fedora scientific


Why Linux

- 
- 1 Access to hardware - almost no restrictions
 - 2 Opensource - view code , modify , learn create
 - 3 Tailor Made - servers , fedora scientific
 - 4 Free

Why Linux

- 
- 1 Access to hardware - almost no restrictions
 - 2 Opensource - view code , modify , learn create
 - 3 Tailor Made - servers , fedora scientific
 - 4 Free
 - 5 Sabko Cadence chalana he

Why Linux


- 
- 1 Access to hardware - almost no restrictions
 - 2 Opensource - view code , modify , learn create
 - 3 Tailor Made - servers , fedora scientific
 - 4 Free
 - 5 Sabko Cadence chalana he
 - 6 Friend dont let friends use windows

Why Linux

- 1 Access to hardware - almost no restrictions
- 2 Opensource - view code , modify , learn create
- 3 Tailor Made - servers , fedora scientific
- 4 Free
- 5 Sabko Cadence chalana he
- 6 Friend dont let friends use windows
- 7 Microsoft gives you windows . Linux gives you whole house

Tables and Figures

- Use `tabular` for basic tables – see Table 1, for example.
- You can upload a figure (JPEG, PNG or PDF) using the files menu.
- To include it in your document, use the `includegraphics` command (see the comment below in the source code).



Item	Quantity
Widgets	42
Gadgets	13

Table 1: An example table.

Readable Mathematics

Let X_1, X_2, \dots, X_n be a sequence of independent and identically distributed random variables with $E[X_i] = \mu$ and $\text{Var}[X_i] = \sigma^2 < \infty$, and let

$$S_n = \frac{X_1 + X_2 + \dots + X_n}{n} = \frac{1}{n} \sum_{i=1}^n X_i$$

denote their mean. Then as n approaches infinity, the random variables $\sqrt{n}(S_n - \mu)$ converge in distribution to a normal $\mathcal{N}(0, \sigma^2)$.