



MITx 6.86x

Machine Learning with Python-From Linear Models to Deep Learning[Course](#)[Progress](#)[Dates](#)[Discussion](#)[Resources](#)[Course](#) / [Unit 0. Brief Prerequisite Reviews, Homework 0, and Project 0](#) / [Home](#)[← Previous](#)

8 of 15 ▾

8. 1D Optimization via Calculus

 [Bookmark this page](#)

Homework0 due Feb 8, 2023 08:59 -03 Completed

Review: 1D Optimization via Calculus

4/4 points (graded)

(For this problem, you are welcome to use any computational tools that would be helpful.)

Let $f(x) = \frac{1}{3}x^3 - x^2 - 3x + 10$ defined on the interval $[-4, 4]$.Let x_1 and x_2 be the critical points of f , and let's impose that $x_1 < x_2$. Fill in the next two values of x_1 and x_2 , respectively: (Recall that the **critical points** of f are those $x \in \mathbb{R}$ $x_1 =$ ✓ $x_2 =$ ✓Fill in the next two boxes with the values of $f''(x_1)$ and $f''(x_2)$, respectively: $f''(x_1) =$ ✓ $f''(x_2) =$ ✓

You have used 1 of 3 attempts

Review: 1D Optimization via Calculus (Continued)

3/4 points (graded)

(For this problem, you are welcome to use any computational tools that would be helpful.)

Recall that x_1 and x_2 are the critical points of the function $f(x) = \frac{1}{3}x^3 - x^2 - 3x + 10$.According to the second derivative test, x_1 is a ...☒ Local Maximum☐ Local Minimum☐ None of the above



At what value of x the (global) maximum value of $f(x)$ attained on the interval $[-2, 2]$?



Submission button: The "submit button" will activate only after you have answered all multiple choices, in this problem.

Submit

You have used 2 of 2 attempts

[< Previous](#)

[Next >](#)

Strict Concavity



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

[News](#)

Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Do Not Sell My Personal Information](#)

Connect

[Blog](#)

[Contact Us](#)