

MITx 6.86x

### **Machine Learning with Python-From Linear Models to Deep Learning**

Discussion Course **Progress** Dates Resources

☆ Course / Unit 2. Nonlinear Classification, Linear regression, Col... / Lecture 5.



# 4. Empirical Risk

☐ Bookmark this page

Exercises due Mar 8, 2023 08:59 -03 Completed

the Objective: Empirical Risk



**1**0v

#### **Video**

**▲** Download video file

#### **Transcripts**

- ▲ Download SubRip (.srt) file
- **▲** Download Text (.txt) file

## **Compute Hinge Loss**

1/1 point (graded)

The empirical risk  $oldsymbol{R_n}$  is defined as

$$R_n\left( heta
ight) = rac{1}{n} \sum_{t=1}^n \operatorname{Loss}\left(y^{(t)} - heta \cdot x^{(t)}
ight)$$

where  $(x^{(t)}, y^{(t)})$  is the tth training example (and there are n in total), and Loss is son hinge loss.


Recall from a previous lecture that the definition of hinge loss:


Lecture 5. Linear Regre	ssion   Unit 2. Nonlinear Classification, Linear regression, Collaborative Filt https://learning.edx.org/course/course-v
	Also, we have .
	Compute the value of
	1.2500
	Submit You have used 1 of 3 attempts
	Compute Squared Error Loss  1/1 point (graded)  Now, we will calculate the empirical risk with the squared error loss. Remember that the
	given by
	The 4 training examples are as in the previous problem:

# Geometrically Identifying Error

1/1 point (graded)

What type of error does the figure below depict? The blue dots are the training example the predictor





Estimation error




Submit

You have used 1 of 1 attempt

## Increasing the Number of Training Examples

0/1 point (graded)

If we increase , the number of training examples, which of the following types of error



Estimation error



Submit

You have used 1 of 1 attempt

#### (Optional) Error decomposition and the bias-variance trade-off

### Discussion

Topic: Unit 2. Non

Next >

weeks):Lecture 5. Linear Regression / 4. Empirical Risk

Show all posts

- ? Squaring only magnifies differences if they are larger than one
- ? Why do we use Hinge Loss here?

Why do we penalize for deviation that is <1 and do not penalize for deviation that is >=1? It doesn't make ser

- Error decomposition and the bias-variance trade-off
  the third term: is E [( error)^2] independent of the bias-variance trade-off. if so, then why. The whole idea of
- ? How to get the result in optional part
- Wrhat is the exact definiation of z?

In the explanation of the problems they always refer to the Loss function L(z), but nowhere is defined what z

Empirical risk

What does she mean by "projected empirical risk"?

- ? (Optional) Error decomposition and the bias-variance trade-off
- Last Problem

I believe the definition of 'large amount of training data' is quite arbitrary which, in my opinion, somewhat con

Confusing explanation about Empirical Risk Rn

Rn is defined as: Sum(Loss(y(t) - theta.X(t)) / n and then they mention Loss can be for instance the hinge los

- Confusion about the deviation
- Rationale of choosing square function

Imho, in addition to what the professor mentioned(tolerate small deviation, truly penalize large deviation), as



# edX

**About** 

**Affiliates** 

edX for Business

Open edX

Careers

<u>News</u>

# Legal

Terms of Service & Honor Code

**Privacy Policy** 

**Accessibility Policy** 

**Trademark Policy** 

**Sitemap** 

**Cookie Policy** 

Do Not Sell My Personal Information

# **Connect**

Blog

**Contact Us** 

Help Center

Security

Media Kit















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 粤ICP备17044299号-2