

Lecture 6a

Correlation

Outline

Recap: Data science vs. experimentation

Correlation between single variables

- ▶ Pearson's correlation
- ▶ Connection to predictability

Correlation between multiple variables

- ▶ Motivations
- ▶ Canonical correlation analysis (CCA)

Application 1

- ▶ Learning equations with CCA

Part 1


Data Science vs. Experimentation (Recap)

Data Science vs. Experimentation

Two main approaches to empirical science:


Experimentation

- ▶ The system of interest is available, and can be manipulated.
- ▶ Hypothesis are first stated and then guide the experimental protocol, i.e. which manipulation is applied to the system and which observables are measured.



Data science


- ▶ Data is simply there, before any hypothesis has been stated.
- ▶ Correlation between features in the data can be measured, and can be used to suggest mechanistic/causal relation between variables.



Data Science Examples

Examples:

- ▶ *Cluster Analysis*: Find cluster structures in the data can suggest meaningful categories or a taxonomy.
- ▶ *Correlation Analysis*: Find that two variables are correlated can suggest a causal or mechanistic link between these two variables, so that changing one variable would incur a change on the other variable.



Data Science vs. Experimentation


Advantages of Data Science:

- ▶ No need to manipulate the system at hand (less time-consuming, avoids ethical issues).
- ▶ Can cover a much broader hypothesis space than a single experiment.

Limitations of Data Science:


- ▶ Bound to the data that is currently available (usually not possible to collect further observations from the system of interest).
- ▶ Identified correlations do not necessarily imply a causal link between the two variables. Actual experiments need to be performed to establish this causal relationship.

Dispersion: (Lectures 3–5)



- ▶ Identifying combination of measurements along which data varies the most (PCA).
- ▶ Find clusters that capture most variation in the data (k-Means, etc.).

Correlation: (Lectures 6–7)



- ▶ Find the level of correlation between two data modalities (Pearson's correlation, CCA).
- ▶ Find how predictable one modality is from the other (regression, discriminant analysis).

Part 2

Pearson's Correlation

Pearson's Correlation


- ▶ Let $(x_i, y_i)_{i=1}^N$ be our dataset where each instance i has two associated features x_i and y_i . Let $\mu_x = E[x]$ and $\mu_y = E[y]$ be the dataset means for these two features.
- ▶ The *Pearson's correlation* associated to this dataset is defined as:

$$\rho = \frac{\overbrace{E[(x - \mu_x)(y - \mu_y)]}^{\sigma_{xy}}}{\underbrace{\sqrt{E[(x - \mu_x)^2]}}_{\sigma_x} \underbrace{\sqrt{E[(y - \mu_y)^2]}}_{\sigma_y}}$$

- ▶ It is always contained in the interval $[-1, 1]$. A value of 1 indicates that x and y are perfectly correlated, -1 indicates perfect negative correlation, and 0 indicates that x and y are decorrelated.
- ▶ Unlike the variance of the data (lectures 3–5), the Pearson's correlation is invariant to the scaling of the data.

Pearson's correlation

Examples:



- Pearson's correlation can only detect certain types of correlations (linear correlations) and may consequently underestimate the true level of correlation.

Correlation as Mutual Predictability


Assume the data is centered ($\mu_x = \mu_y = 0$). We build a homogeneous linear model that predicts y from x in a way that minimizes square errors:

$$\begin{aligned} \arg \min_{\beta} E[(y - \beta x)^2] \\ &= \arg \min_{\beta} E[\beta^2 x^2 - 2\beta xy] \\ &= \arg \min_{\beta} \beta^2 \sigma_x^2 - 2\beta \sigma_{xy} \end{aligned}$$

This is a convex optimization problem with solution found at:

$$\frac{\partial}{\partial \beta} (\beta^2 \sigma_x^2 - 2\beta \sigma_{xy}) = 0$$


which gives us the closed form $\beta = \sigma_{xy} / \sigma_x^2$.



Correlation as Mutual Predictability

Injecting the closed form $\beta = \sigma_{xy}/\sigma_x^2$ in the error function gives us the error at the optimum (or some measure of predictivity of y from x).

$$\begin{aligned}\min_{\beta} E[(y - \beta x)^2] \\&= E[(y - \frac{\sigma_{xy}}{\sigma_x^2} x)^2] \\&= E[y^2] - 2 \frac{\sigma_{xy}}{\sigma_x^2} E[xy] + \left(\frac{\sigma_{xy}}{\sigma_x^2}\right)^2 E[x^2] \\&= \sigma_y^2 - \frac{\sigma_{xy}^2}{\sigma_x^2} \\&= \sigma_y^2 \cdot \underbrace{\left(1 - \left(\frac{\sigma_{xy}}{\sigma_x \sigma_y}\right)^2\right)}_{\rho}\end{aligned}$$



Correlation as Mutual Predictability

We found that the residual prediction error of the optimal linear model is given by:

$$\begin{aligned}\min_{\beta} \mathbb{E}[(y - \beta x)^2] \\ = \sigma_y^2 \cdot (1 - \rho^2)\end{aligned}$$

Conversely, we can predict x from y with error given by

$$\begin{aligned}\min_{\beta} \mathbb{E}[(\beta y - x)^2] \\ = \sigma_x^2 \cdot (1 - \rho^2)\end{aligned}$$

Conclusion: There is a direct relation between Pearson's correlation and mutual predictability (in the least squares sense, using a linear model).

Part 3

Canonical Correlation Analysis

Suggested reading:

Borga (2001): Canonical Correlation a Tutorial

Motivations: Multivariate Data

Question:

- ▶ Are images and their associated text correlated?



"motorcycle front wheel"



"thumbnail for version as of 21
57 29 june 2010"



"file frankfurt airport
skyline 2017 05 jpg"



"file london barge race 2 jpg"



"wild boar head portrait forest
creature boar"



"st oswalds way and shops"

Image source: <https://ai.googleblog.com/2021/05/align-scaling-up-visual-and-vision.html>


Problem:

- ▶ Images and text are multivariate (composed of multiple visual features or words).
- ▶ We need to generalize correlation to modalities of more than one dimension.

Motivations: Neuro-Vascular Coupling

intercortical neural activity

FMRI/BOLD signal



haemodynamic response

Motivations: Climate Dynamics





Image source: Climate co-variability between South America and Southern Africa at interannual, intraseasonal and synoptic scales.
DOI:10.1007/s00382-016-3318-x

Canonical Correlation Analysis



Formalization:

- ▶ Let $\mathbf{x} \in \mathbb{R}^d$ be our first modality and $\mathbf{y} \in \mathbb{R}^h$ be our second modality.
- ▶ Find projections $\mathbf{w}_x \in \mathbb{R}^d$ and $\mathbf{w}_y \in \mathbb{R}^h$ of the two modalities in which the Pearson's correlation is maximized:

$$\arg \max_{\mathbf{w}} \text{Corr}(\mathbf{w}_x^\top \mathbf{x}, \mathbf{w}_y^\top \mathbf{y})$$

($\mathbf{w} = (\mathbf{w}_x, \mathbf{w}_y)$). Note: Because Pearson's correlation is invariant to the scale of the data, the vectors \mathbf{w}_x and \mathbf{w}_y don't need to be constrained to be e.g. of unit norm.

Canonical Correlation Analysis

Observation:

- ▶ The correlation can be rewritten as:

$$\begin{aligned}\text{Corr}(\mathbf{w}_x^\top \mathbf{x}, \mathbf{w}_y^\top \mathbf{y}) &= \frac{\mathbb{E}[(\mathbf{w}_x^\top (\mathbf{x} - \boldsymbol{\mu}_x)) \cdot (\mathbf{w}_y^\top (\mathbf{y} - \boldsymbol{\mu}_y))]}{\sqrt{\mathbb{E}[(\mathbf{w}_x^\top (\mathbf{x} - \boldsymbol{\mu}_x))^2]} \sqrt{\mathbb{E}[(\mathbf{w}_y^\top (\mathbf{y} - \boldsymbol{\mu}_y))^2]}} \\ &= \frac{\mathbf{w}_x^\top \mathbf{C}_{xy} \mathbf{w}_y}{\sqrt{\mathbf{w}_x^\top \mathbf{C}_{xx} \mathbf{w}_x} \sqrt{\mathbf{w}_y^\top \mathbf{C}_{yy} \mathbf{w}_y}}\end{aligned}$$

where

$$\mathbf{C}_{xy} = \mathbb{E}[(\mathbf{x} - \boldsymbol{\mu}_x)(\mathbf{y} - \boldsymbol{\mu}_y)^\top]$$

$$\mathbf{C}_{xx} = \mathbb{E}[(\mathbf{x} - \boldsymbol{\mu}_x)(\mathbf{x} - \boldsymbol{\mu}_x)^\top]$$

$$\mathbf{C}_{yy} = \mathbb{E}[(\mathbf{y} - \boldsymbol{\mu}_y)(\mathbf{y} - \boldsymbol{\mu}_y)^\top]$$

are cross-covariance and auto-covariance matrices that can be precomputed.

Canonical Correlation Analysis

Observation (2):

- ▶ Remember that the optimum of the problem:

$$\arg \max_{\mathbf{w}} \frac{\mathbf{w}_x^\top C_{xy} \mathbf{w}_y}{\sqrt{\mathbf{w}_x^\top C_{xx} \mathbf{w}_x} \sqrt{\mathbf{w}_y^\top C_{yy} \mathbf{w}_y}}$$

is specified up to a scaling of \mathbf{w}_x and \mathbf{w}_y .

Idea:

- ▶ Force a particular scale of \mathbf{w}_x and \mathbf{w}_y , and get in turn a simpler optimization problem.
- ▶ In particular, the optimization problem can be reformulated as a standard quadratic optimization problem with quadratic constraints (like for PCA):

$$\arg \max_{\mathbf{w}} \mathbf{w}_x^\top C_{xy} \mathbf{w}_y \quad \text{s.t.} \quad \begin{aligned} \mathbf{w}_x^\top C_{xx} \mathbf{w}_x &= 1 \\ \mathbf{w}_y^\top C_{yy} \mathbf{w}_y &= 1 \end{aligned}$$

Canonical Correlation Analysis

Idea (2):

- ▶ For the constrained optimization problem

$$\arg \max_{\mathbf{w}} \mathbf{w}_x^\top C_{xy} \mathbf{w}_y \quad \text{s.t.} \quad \begin{aligned} \mathbf{w}_x^\top C_{xx} \mathbf{w}_x &= 1 \\ \mathbf{w}_y^\top C_{yy} \mathbf{w}_y &= 1 \end{aligned}$$

one can obtain using the method of Lagrange multipliers a closed form solution (→ Lecture 6b for the derivation).

Final formulation:

- ▶ One gets that the solution of CCA is the first eigenvector (i.e. with highest eigenvalue) of the generalized eigenvalue problem:

$$\begin{bmatrix} 0 & C_{xy} \\ C_{yx} & 0 \end{bmatrix} \begin{bmatrix} \mathbf{w}_x \\ \mathbf{w}_y \end{bmatrix} = \lambda \begin{bmatrix} C_{xx} & 0 \\ 0 & C_{yy} \end{bmatrix} \begin{bmatrix} \mathbf{w}_x \\ \mathbf{w}_y \end{bmatrix}$$


and the latter can be solved with common libraries (e.g. scipy).

- ▶ The eigenvalue λ represent the correlation ρ of data in the two projected spaces.

Part 4

Application: Nonlinear Correlations

Application: Finding Nonlinear Correlations




Observation:

- ▶ Negligible Pearson's correlation coefficient.
- ▶ However, it is clear visually that x and y are still related in some way.

Idea:

- ▶ Nonlinearly expand x and y , and apply CCA to find correlation between x and y in expanded space.

Application: Finding Nonlinear Correlations




- ▶ CCA can explore possible nonlinear correlations.
- ▶ Correlation found by CCA should be in theory at least as good original correlation (if including x and y in the feature map).

Application: Finding Nonlinear Correlations

CCA Result:

$$\lambda_1 = 0.86 \quad \mathbf{u}_1 = \left[\underbrace{-0.13, 1.84, 0.15}_{\mathbf{w}_x}, \underbrace{0.09, -1.88, -0.07}_{\mathbf{w}_y} \right]$$

- Projecting expanded modalities on this eigenvector, gives a new scatter plot, where a linear correlation can be observed and measured:



- Correlation strength was readily given by the first eigenvalue of the CCA generalized eigenvalue problem.

Application: Finding Nonlinear Correlations

CCA Result:

$$\lambda_1 = 0.86 \quad \mathbf{u}_1 = \left[\underbrace{-0.13, 1.84, 0.15}_{\mathbf{w}_x}, \underbrace{0.09, -1.88, -0.07}_{\mathbf{w}_y} \right]$$

- An inspection of this result suggests that approximately:

$$\text{Corr}(0 + x^2 + 0, 0 - y^2 + 0) \approx 0.86$$


In other words, it suggests that the data is governed by the equation:

$$x^2 = \alpha(-y^2) + \beta$$

The true equation that describes the data would be obtained by setting $\alpha = 1$ and $\beta = 1$, i.e. $x^2 + y^2 = 1$.

Application: Finding Nonlinear Correlations


Example 2:



$$\lambda_1 = 0.88 \quad \mathbf{u}_1 = \left[\underbrace{0.16, 2.04, -0.12}_{\mathbf{w}_x}, \underbrace{-0.09, 2.00, 0.16}_{\mathbf{w}_y} \right]$$

Application: Finding Nonlinear Correlations

Example 3:



$$\lambda_1 = 0.86 \quad u_1 = \left[\underbrace{1.84, -0.11, -1.06, -1.06}_{w_x}, \underbrace{0.00, 0.15}_{w_y} \right]$$

Summary

Summary

- ▶ Finding correlations is an important component of data science as it allows to identify potential causal or mechanistic relations in some system of interest.
- ▶ Pearson's correlation is the most established measure of correlation and it has many connections to other statistical properties (e.g. explained variance, error of a linear model, etc.).
- ▶ Canonical correlation analysis generalizes Pearson's correlation to the case where the two modalities one would like to correlate are multidimensional.
- ▶ Canonical correlation analysis can also be used to discover nonlinear relations in bivariate data.