2020

PHYSICS — HONOURS

Paper: DSE-B-2

(Nuclear and Particle Physics)

Full Marks: 65

The figures in the margin indicate full marks.

Candidates are required to give their answers in their own words as far as practicable.

Group - A

1. Answer any five questions:

 2×5

- (a) Find the distance of closest approach of 1 MeV proton incident on gold nucleus (z = 79) with zero impact parameter.
- (b) What do you mean by 'charge independence' and 'charge symmetry' of nuclear force?
- (c) Find the ground state spin parity of 12Mg²⁵.
- (d) Explain why a Geiger Counter cannot measure the energy of moving charge particle.
- (e) What are the limitations of a fixed frequency cyclotron?
- (f) A hadron has a quark content ddu. Find the charge and strangeness of this hadron.
- (g) Are the following processes allowed in strong interaction?

(i)
$$\pi^- + n \rightarrow \Sigma^- + K^o$$

(ii)
$$\pi^- + p \rightarrow \wedge^{o} + K^{o}$$
.

Group - B

Answer any three questions.

5×3

- 2. The density of iron is 8×10^3 kg/m³. The neutron capture cross-section of iron is 2.5 barn. What fraction of a normally incident neutron beam is absorbed by an iron sheet of 0.01 m thick? Explain the formula used.
- 3. (a) Calculate the minimum energy required to be given to the neutron in order that the following nuclear reaction may occur

$$_{\mathrm{o}}n^{1}$$
 + $_{15}$ P 31 \rightarrow $_{14}$ Si 30 + $_{1}$ H 1

Given the masses (in a.m.u.)

$$M \begin{pmatrix} 0 & n^{1} \end{pmatrix} = 1.008665$$
 $M \begin{pmatrix} 15 & P^{31} \end{pmatrix} = 30.973766$
 $M \begin{pmatrix} 14 & Si^{31} \end{pmatrix} = 30.975349$ $M \begin{pmatrix} 1 & H^{1} \end{pmatrix} = 1.007825$

(b) What do you mean by thermal neutrons? Indicate their key role in nuclear reaction. 3+(1+1)

Please Turn Over

- 4. (a) Draw the characteristic curve of G.M. counter. Define threshold voltage.
 - (b) An organic quenched GM tube operates at 1000 V and has a wire having diameter 0.2 mm. The radius of the cathode is 2 cm. What is the maximum radial field? (2+1)+2
- **5.** Explain working principle of semiconductor detector. What is the major advantage of this detector over others?
- **6.** Explain why the following processes are not allowed.

2+2+1

- (a) $p + \pi^{o} \rightarrow p^{-} + \pi^{+} + \pi^{-}$
- (b) $n \rightarrow p + e^-$
- (c) $e^- \rightarrow v_e + \gamma$.

Group - C

Answer any four questions.

 10×4

- 7. (a) Write down the Bethe-Weiszäcker formula for binding energy of a nucleas, explain all the terms therein.
 - (b) Exlain graphically how the binding energy per nucleon varies with mass number on an average, as a result of the various terms mentioned.
 - (c) What is meant by saturation of nuclear force?
 - (d) Show that $\gamma \rightarrow e^+ + e^-$ process cannot take place in vacuum.

3+3+1+3

- 8. (a) Define threshold energy and derive an expression for the threshold energy of an endoergic reaction
 - (b) Write down an expression for the cross-section of a nuclear reaction, clearly explaining all the terms therein.
 - (c) A 0.01 mm thick ${}_{3}\text{Li}^{4}$ target is bombarded with 10^{13} protons per second. As a result 10^{6} neutrons per second are produced. What would be the cross-section for the reaction? (Density of ${}_{3}\text{Li}^{4} = 500 \text{ kg} / \text{m}^{3}$) (1+4)+2+3
- **9.** (a) Derive the Bethe-Block formula for the energy loss of a moving charged particle inside a matter due to ionisation.
 - (b) Explain the Compton wavelength shift.
 - (c) Compute the maximum energy of the Compton recoil electrons resulting from the absorption of Al of 2.19 MeV γ -rays. 4+3+3
- **10.** (a) What is the dead time of a GM counter? A GM counter has dead time of 200 μs. What are the true counting rates when the observed rates are 1000 per minute?
 - (b) Explain the basic principle of photomultiplier tube (PMT).
 - (c) What is scintillation process? Why is photomultiplier tube used in a scintilation detector? (2+2)+3+(1+2)

- 11. (a) Briefly explain the working principle of a cyclotron.
 - (b) Calculate the maximum energy of protons obtainable from a cyclotron having dees of diameter 1.2 m each and 1.5 T magnetic field. At what frequency must the cyclotron be operated? If the average energy gain per dee passage is $50 \text{ k}_0\text{V}$, how many revolutions do the proton make? 5+(2+1+2)
- **12.** (a) What is meant by Eightfold way? Explain with reference to the octate symmetry of particle physics.
 - (b) Define lepton number and baryon number. Write down the equation for muon decay. How is lepton number conserved in this process?
 - (c) The decay $\equiv^- \to \wedge^0 + \pi^-$ is observed in nature, whereas the apparently similar decay $\equiv^- \to n^0 + \pi^-$ is never observed. Why? (1+3)+(2+1+1)+2