

Supplementary Note 6: Detailed statistical document of article figures

June 13, 2023

Contents

1	Figure 2	3
1.1	Figure 2a	3
1.2	Figure 2b	3
1.3	Figure 2c	4
1.4	Figure 2d	4
1.5	Figure 2e	4
1.6	Figure 2e (With approach trials)	5
1.7	Figure 2f	5
1.8	Figure 2f (With approach trials)	6
1.9	Figure 2g	6
1.10	Figure 2g (With approach trials)	7
1.11	Figure 2h	7
1.12	Figure 2h (With approach trials)	8
2	Figure 3	8
2.1	Figure 3e	8
2.2	Figure 3f	8
2.3	Figure 3g	9
2.4	Figure 3h	9
3	Figure 5	9
3.1	Figure 5a	9
3.2	Figure 5b	9
3.3	Figure 5c	10
3.4	Figure 5d	10
3.5	Figure 5e	11
3.6	Figure 5f	11
3.7	Figure 5g	12
3.8	Figure 5i (Left)	12
3.9	Figure 5i (Right)	12
4	Figure 6	12
4.1	Figure 6b (Left)	12
4.2	Figure 6b (Right)	13
4.3	Figure 6c	13
4.4	Figure 6d (Left)	15
4.5	Figure 6d (Right)	15
4.6	Figure 6e	15
4.7	Figure 6f	16
4.8	Figure 6g	16
4.9	Figure 6h	17
4.10	Figure 6k	17

5	Supplemental Figure 1	18
5.1	Figure S.1i	18
6	Supplemental Figure 2	18
6.1	Figure S.2a	18
6.2	Figure S.2b	18
7	Supplemental Figure 4	18
7.1	Figure S.4d	18
8	Supplemental Figure 6	18
8.1	Figure S.6a	18
8.2	Figure S.6b	19
8.3	Figure S.6c	19
8.4	Figure S.6d	19
8.5	Figure S.6e	20
9	Supplemental Figure 7	20
9.1	Figure S.7a (Left)	20
9.2	Figure S.7a (Right)	21
9.3	Figure S.7b (Left)	21
9.4	Figure S.7b (Right)	22
9.5	Figure S.7d	22
9.6	Figure S.7e	23
9.7	Figure S.7f	23
9.8	Figure S.7g	24
9.9	Figure S.7h	24
9.10	Figure S.7i	25
9.11	Figure S.7j	25
9.12	Figure S.7k	26

A repeated measures analysis of variance (ANOVA) was conducted using the MATLAB `ranova` function to examine the effects of within-subject factors, such as sucrose concentration, and between-subject factors, including gender and experimental conditions (baseline vs. food deprivation). To assess the between-subject differences, a two-sample Kolmogorov-Smirnov test was employed with the MATLAB `kstest2` function. Additionally, pairwise comparisons were conducted to further explore the differences between groups using a post-hoc analysis, specifically the Tukey's honestly significant difference method, implemented with the MATLAB `multcompare` function.

1 Figure 2

1.1 Figure 2a

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 7.3397e-26.

p-value for gender: 6.3880e-02.

`kstest2` results: $h=0$, $p=8.2894e-02$, $ks2stat=0.2557$ (overall gender difference)

Post-hoc analysis:

$$0.5\% : 5.9941e - 01, \quad 2\% : 2.0283e - 02, \quad 5\% : 1.0142e - 01, \quad 9\% : 5.3385e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 0, p = 0.3032, ks2stat = 0.3788$$

$$RStest : Conc1 : h = 0, p = 0.4049, zval = 0.8329$$

$$KStest2 : Conc2 : h = 1, p = 0.0087, ks2stat = 0.6439$$

$$RStest : Conc2 : h = 1, p = 0.0187, zval = 2.3510$$

$$KStest2 : Conc3 : h = 0, p = 0.2812, ks2stat = 0.3864$$

$$RStest : Conc3 : h = 0, p = 0.1314, zval = 1.5086$$

$$KStest2 : Conc4 : h = 0, p = 0.9465, ks2stat = 0.2045$$

$$RStest : Conc4 : h = 0, p = 0.5156, zval = -0.6501$$

1.2 Figure 2b

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 2.061e-10.

p-value for gender: 0.15301.

`kstest2` results: $h=0$, $p=9.9819e-02$, $ks2stat=0.2481$ (overall gender difference)

Post-hoc analysis:

$$240lux : 1.8263e - 01, \quad 260lux : 5.1534e - 02, \quad 290lux : 8.8968e - 01, \quad 320 : 3.7194e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.5833, ks2stat = 0.3030$
 $RStest : Conc1 : h = 0, p = 0.2815$
 $KStest2 : Conc2 : h = 0, p = 0.2407, ks2stat = 0.4015$
 $RStest : Conc2 : h = 0, p = 0.0602$
 $KStest2 : Conc3 : h = 0, p = 0.6484, ks2stat = 0.2879$
 $RStest : Conc3 : h = 0, p = 1.0000$
 $KStest2 : Conc4 : h = 0, p = 0.7136, ks2stat = 0.2727$
 $RStest : Conc4 : h = 0, p = 0.4235$

1.3 Figure 2c

Statistical significance was determined using **Statistical Package for the Social Sciences (SPSS)** package (F = 12, M = 9)

p-value for concentration: <0.0001.

Sex differences across all concentrations $p = 0.8$.

Post-hoc analysis:

0.5% : $p = 0.05$, 2% : $p = 0.007$, 5% : $p = 0.8$, 9% : $p = 0.4$.

1.4 Figure 2d

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 7.8859e-04.

p-value for gender: 4.3484e-01.

kstest2 results: $h=0$, $p=3.1096e-01$, $ks2stat=0.1986$ (overall gender difference)

Post-hoc analysis:

0.5% : $8.9816e - 01$, 2% : $4.5069e - 01$, 5% : $5.3396e - 01$, 9% : $5.9227e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.7358, ks2stat = 0.2727$
 $RStest : Conc1 : h = 0, p = 0.6458, zval = -0.4597$
 $KStest2 : Conc2 : h = 0, p = 0.4896, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.6682, zval = -0.4286$
 $KStest2 : Conc3 : h = 0, p = 0.8286, ks2stat = 0.2500$
 $RStest : Conc3 : h = 0, p = 0.7169, zval = -0.3627$
 $KStest2 : Conc4 : h = 0, p = 0.7136, ks2stat = 0.2727$
 $RStest : Conc4 : h = 0, p = 0.7350, zval = -0.3385$

1.5 Figure 2e

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 8.9008e-02.

p-value for gender: 2.4720e-03.

kstest2 results: $h=1$, $p=9.4199e-06$, $ks2stat=0.5019$ (overall gender difference)

Post-hoc analysis:

$$0.5\% : 1.2707e - 03, \quad 2\% : 1.1033e - 02, \quad 5\% : 3.5299e - 03, \quad 9\% : 8.9459e - 03$$

KStest2 and wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$\begin{aligned} KStest2 : Conc1 : h = 1, p = 0.0258, ks2stat = 0.5758 \\ RStest : Conc1 : h = 1, p = 0.0051, zval = 2.8003 \\ KStest2 : Conc2 : h = 1, p = 0.0361, ks2stat = 0.5530 \\ RStest : Conc2 : h = 1, p = 0.0151, zval = 2.4311 \\ KStest2 : Conc3 : h = 1, p = 0.0230, ks2stat = 0.5833 \\ RStest : Conc3 : h = 1, p = 0.0051, zval = 2.8003 \\ KStest2 : Conc4 : h = 1, p = 0.0361, ks2stat = 0.5530 \\ RStest : Conc4 : h = 1, p = 0.0062, zval = 2.7388 \end{aligned}$$

1.6 Figure 2e (With approach trials)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 0.00016745.

p-value for gender: 0.0037713.

kstest2 results: h=1, p=7.1912e-04, ks2stat=0.4103 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 1.2870e - 02, \quad 2\% : 2.5797e - 01, \quad 5\% : 2.5247e - 02, \quad 9\% : 1.7835e - 02$$

KStest2 and wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$\begin{aligned} KStest2 : Conc1 : h = 1, p = 0.0121, ks2stat = 0.6364 \\ RStest : Conc1 : h = 1, p = 0.0104 \\ KStest2 : Conc2 : h = 0, p = 0.2270, ks2stat = 0.4167 \\ RStest : Conc2 : h = 0, p = 0.3734 \\ KStest2 : Conc3 : h = 1, p = 0.0452, ks2stat = 0.5500 \\ RStest : Conc3 : h = 1, p = 0.0192 \\ KStest2 : Conc4 : h = 1, p = 0.0289, ks2stat = 0.5682 \\ RStest : Conc4 : h = 1, p = 0.0074 \end{aligned}$$

1.7 Figure 2f

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 9.8312e-01

p-value for gender: 1.5572e-01

kstest2 results: h=1, p=2.5533e-05, ks2stat=0.4811 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 7.9690e - 02, \quad 2\% : 2.5673e - 01, \quad 5\% : 1.4691e - 01, \quad 9\% : 2.0322e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0067, ks2stat = 0.6591$
 $RStest : Conc1 : h = 1, p = 0.0028, zval = -2.9850$
 $KStest2 : Conc2 : h = 0, p = 0.2604, ks2stat = 0.3939$
 $RStest : Conc2 : h = 0, p = 0.1029, zval = -1.6310$
 $KStest2 : Conc3 : h = 1, p = 0.0323, ks2stat = 0.5606$
 $RStest : Conc3 : h = 1, p = 0.0289, zval = -2.1849$
 $KStest2 : Conc4 : h = 0, p = 0.0915, ks2stat = 0.4848$
 $RStest : Conc4 : h = 0, p = 0.0905, zval = -1.6925$

1.8 Figure 2f (With approach trials)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 0.0014948

p-value for gender: 0.4856

kstest2 results: h=0, p=6.2873e-02, ks2stat=0.2710 (overall gender difference)

Post-hoc analysis:

0.5% : $5.8083e - 02$ 2% : $8.8005e - 01$, 5% : $5.5419e - 01$, 9% : $1.6653e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.1473, ks2stat = 0.4545$
 $RStest : Conc1 : h = 0, p = 0.0878$
 $KStest2 : Conc2 : h = 0, p = 0.4896, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.5310$
 $KStest2 : Conc3 : h = 0, p = 0.2689, ks2stat = 0.4000$
 $RStest : Conc3 : h = 0, p = 0.1985$
 $KStest2 : Conc4 : h = 0, p = 0.0915, ks2stat = 0.4848$
 $RStest : Conc4 : h = 0, p = 0.0905$

1.9 Figure 2g

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 2.6239e-10.

p-value for gender: 1.7542e-03.

kstest2 results: h=1, p=3.0470e-05, ks2stat=0.4773 (overall gender difference)

Post-hoc analysis:

0.5% : $1.6629e - 04$, 2% : $2.5835e - 03$, 5% : $5.1037e - 03$, 9% : $6.0776e - 02$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0003, ks2stat = 0.8258$
 $RStest : Conc1 : h = 1, p = 0.0006, zval = -3.4158$
 $KStest2 : Conc2 : h = 1, p = 0.0059, ks2stat = 0.6667$
 $RStest : Conc2 : h = 1, p = 0.0042, zval = -2.8619$
 $KStest2 : Conc3 : h = 1, p = 0.0323, ks2stat = 0.5606$
 $RStest : Conc3 : h = 1, p = 0.0106, zval = -2.5541$
 $KStest2 : Conc4 : h = 0, p = 0.1213, ks2stat = 0.4621$
 $RStest : Conc4 : h = 0, p = 0.1481, zval = -1.4463$

1.10 Figure 2g (With approach trials)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 1.1834e-12.

p-value for gender: 0.21917.

kstest2 results: h=0, p=4.3505e-01, ks2stat=0.1793 (overall gender difference)

Post-hoc analysis:

0.5% : 1.9769e - 01, 2% : 5.3803e - 01, 5% : 4.1293e - 01, 9% : 1.8367e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.3744, ks2stat = 0.3636$
 $RStest : Conc1 : h = 0, p = 0.1486$
 $KStest2 : Conc2 : h = 0, p = 0.8286, ks2stat = 0.2500$
 $RStest : Conc2 : h = 0, p = 0.9212$
 $KStest2 : Conc3 : h = 0, p = 0.4268, ks2stat = 0.3500$
 $RStest : Conc3 : h = 0, p = 0.1985$
 $KStest2 : Conc4 : h = 0, p = 0.1328, ks2stat = 0.4545$
 $RStest : Conc4 : h = 0, p = 0.1661$

1.11 Figure 2h

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 3.0392e-07.

p-value for gender: 2.3301e-01.

kstest2 results: h=0, p=3.3508e-01, ks2stat=0.1913 (overall gender difference)

Post-hoc analysis:

0.5% : 2.6980e - 01, 2% : 7.5679e - 01, 5% : 8.5789e - 02, 9% : 3.0110e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.8067, ks2stat = 0.2500$
 $RStest : Conc1 : h = 0, p = 0.4044, zval = 0.8338$
 $KStest2 : Conc2 : h = 0, p = 0.9982, ks2stat = 0.1515$
 $RStest : Conc2 : h = 0, p = 0.8292, zval = 0.2157$
 $KStest2 : Conc3 : h = 0, p = 0.1006, ks2stat = 0.4773$
 $RStest : Conc3 : h = 1, p = 0.0483, zval = 1.9743$
 $KStest2 : Conc4 : h = 0, p = 0.4595, ks2stat = 0.3333$
 $RStest : Conc4 : h = 0, p = 0.4219, zval = 0.8031$

1.12 Figure 2h (With approach trials)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 11).

p-value for concentration: 0.00030532.

p-value for gender: 0.88032.

kstest2 results: h=0, p=7.5063e-01, ks2stat=0.1393 (overall gender difference)

Post-hoc analysis:

0.5% : $6.6119e - 01$, 2% : $4.6695e - 01$, 5% : $5.8985e - 01$, 9% : $6.5657e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.9852, ks2stat = 0.1818$
 $RStest : Conc1 : h = 0, p = 0.8422$
 $KStest2 : Conc2 : h = 0, p = 0.8848, ks2stat = 0.2333$
 $RStest : Conc2 : h = 0, p = 0.3834$
 $KStest2 : Conc3 : h = 0, p = 0.5564, ks2stat = 0.3167$
 $RStest : Conc3 : h = 0, p = 0.4681$
 $KStest2 : Conc4 : h = 0, p = 0.9610, ks2stat = 0.1970$
 $RStest : Conc4 : h = 0, p = 0.5588$

2 Figure 3

2.1 Figure 3e

Statistical significance was determined by one-way analysis of variance. (group 1 = 60, group 2 = 61, group 3 = 58, group 4 = 64, group 5 = 22, group 6 = 22, group 7 = 25, group 8 = 18)

p-value for significance of difference between the groups (utility): 0.0429.

Post-hoc analysis by Tukey's HSD method:

No group difference is statistically significant.

2.2 Figure 3f

Statistical significance was determined by one-way analysis of variance. (group 1 = 60, group 2 = 61, group 3 = 58, group 4 = 64)

p-value for significance of difference between the groups (concentration): 0.9599.

Post-hoc analysis by Tukey's HSD method:

No group difference is statistically significant.

2.3 Figure 3g

Statistical significance was determined by one-way analysis of variance. (group 1 = 22, group 2 = 22, group 3 = 25, group 4 = 18)

p-value for significance of difference between the groups (concentration): 0.5523.

Post-hoc analysis by Tukey's HSD method:

No group difference is statistically significant.

2.4 Figure 3h

Statistical significance was determined by one-way analysis of variance. (group 1 = 243, group 2 = 87)
p-value for significance of difference between the groups (concentration): 0.0012.

3 Figure 5

3.1 Figure 5a

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: 1.0842e-55

p-value for BL vs FD: 8.0411e-05.

kstest2 results: h=1, p=1.0816e-02, ks2stat=0.2386 (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : 6.8154e - 01, 2% : 5.2118e - 01, 5% : 5.0500e - 04, 9% : 4.8848e - 03

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : h = 0, p = 0.1746, ks2stat = 0.3182

RStest : Conc1 : h = 0, p = 0.3038, zval = 1.0283

KStest2 : Conc2 : h = 0, p = 0.3320, ks2stat = 0.2727

RStest : Conc2 : h = 0, p = 0.3820, zval = 0.8743

KStest2 : Conc3 : h = 1, p = 0.0138, ks2stat = 0.4545

RStest : Conc3 : h = 1, p = 0.0011, zval = -3.2659

KStest2 : Conc4 : h = 1, p = 0.0000, ks2stat = 0.6818

RStest : Conc4 : h = 1, p = 0.0003, zval = -3.6583

3.2 Figure 5b

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 12, FD N = 9).

p-value for concentration: 1.3979e-09

p-value for BL vs FD: 4.2358e-01.

kstest2 results: $h=0$, $p=1.0000e+00$, $ks2stat=0.0556$ (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : $4.2051e-01$, 2% : $1.2090e-02$, 5% : $8.9340e-01$, 9% : $8.4742e-01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : $h = 0, p = 0.7495, ks2stat = 0.2778$
RStest : Conc1 : $h = 0, p = 0.5274$
KStest2 : Conc2 : $h = 1, p = 0.0065, ks2stat = 0.6944$
RStest : Conc2 : $h = 1, p = 0.0034$
KStest2 : Conc3 : $h = 0, p = 0.1417, ks2stat = 0.4722$
RStest : Conc3 : $h = 0, p = 0.5421$
KStest2 : Conc4 : $h = 0, p = 0.1915, ks2stat = 0.4444$
RStest : Conc4 : $h = 0, p = 0.2015$

3.3 Figure 5c

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: $6.3645e-10$.

p-value for BL vs FD: $1.9462e-01$.

kstest2 results: $h=0$, $p=3.5436e-01$, $ks2stat=0.1435$ (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : $1.0418e-02$, 2% : $3.2611e-01$, 5% : $7.3375e-01$, 9% : $1.0336e-01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : $h = 0, p = 0.0647, ks2stat = 0.4286$
RStest : Conc1 : $h = 1, p = 0.0127, zval = -2.4921$
KStest2 : Conc2 : $h = 1, p = 0.0395, ks2stat = 0.4286$
RStest : Conc2 : $h = 0, p = 0.0883, zval = -1.7044$
KStest2 : Conc3 : $h = 0, p = 0.7388, ks2stat = 0.1991$
RStest : Conc3 : $h = 0, p = 0.6885, zval = 0.4009$
KStest2 : Conc4 : $h = 0, p = 0.3320, ks2stat = 0.2727$
RStest : Conc4 : $h = 0, p = 0.1625, zval = 1.3966$

3.4 Figure 5d

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: $2.8777e-07$.

p-value for BL vs FD: $1.6989e-03$

kstest2 results: $h=1$, $p=7.5537e-08$, $ks2stat=0.4318$ (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : $3.7073e-04$, 2% : $7.5759e-04$, 5% : $3.3233e-02$, 9% : $1.3234e-02$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0015, ks2stat = 0.5455$
 $RStest : Conc1 : h = 1, p = 0.0008, zval = -3.3683$
 $KStest2 : Conc2 : h = 1, p = 0.0356, ks2stat = 0.4091$
 $RStest : Conc2 : h = 1, p = 0.0028, zval = -2.9928$
 $KStest2 : Conc3 : h = 1, p = 0.0138, ks2stat = 0.4545$
 $RStest : Conc3 : h = 1, p = 0.0151, zval = -2.4294$
 $KStest2 : Conc4 : h = 1, p = 0.0356, ks2stat = 0.4091$
 $RStest : Conc4 : h = 1, p = 0.0151, zval = -2.4294$

3.5 Figure 5e

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: 2.7791e-01.

p-value for BL vs FD: 4.3141e-02

kstest2 results: h=1, p=7.5537e-08, ks2stat=0.4318 (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : 3.9999e - 02, 2% : 3.3272e - 02, 5% : 5.5036e - 02, 9% : 5.9155e - 02

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0001, ks2stat = 0.6364$
 $RStest : Conc1 : h = 1, p = 0.0003, zval = 3.6265$
 $KStest2 : Conc2 : h = 1, p = 0.0138, ks2stat = 0.4545$
 $RStest : Conc2 : h = 1, p = 0.0028, zval = 2.9928$
 $KStest2 : Conc3 : h = 0, p = 0.0828, ks2stat = 0.3636$
 $RStest : Conc3 : h = 1, p = 0.0109, zval = 2.5468$
 $KStest2 : Conc4 : h = 1, p = 0.0138, ks2stat = 0.4545$
 $RStest : Conc4 : h = 1, p = 0.0032, zval = 2.9458$

3.6 Figure 5f

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: 6.6926e-20

p-value for BL vs FD: 2.4891e-02

kstest2 results: h=1, p=1.0816e-02, ks2stat=0.2386 (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : 7.3584e - 04, 2% : 1.0250e - 01, 5% : 2.0052e - 01, 9% : 1.2278e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0015, ks2stat = 0.5455$
 $RStest : Conc1 : h = 1, p = 0.0012, zval = 3.2275$
 $KStest2 : Conc2 : h = 0, p = 0.0828, ks2stat = 0.3636$
 $RStest : Conc2 : h = 0, p = 0.0689, zval = 1.8191$
 $KStest2 : Conc3 : h = 0, p = 0.3320, ks2stat = 0.2727$
 $RStest : Conc3 : h = 0, p = 0.4455, zval = 0.7629$
 $KStest2 : Conc4 : h = 0, p = 0.1746, ks2stat = 0.3182$
 $RStest : Conc4 : h = 0, p = 0.1424, zval = 1.4670$

3.7 Figure 5g

Statistical significance was determined by Repeated measures analysis of variance. (BL N = 22, FD N = 22).

p-value for concentration: 8.1104e-22

p-value for BL vs FD: 1.2044e-01

kstest2 results: h=1, p=1.7572e-02, ks2stat=0.2273 (overall difference in BL vs FD)

Post-hoc analysis:

0.5% : 5.6817e - 03, 2% : 1.9624e - 02, 5% : 5.4119e - 01, 9% : 7.4789e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0049, ks2stat = 0.5000$
 $RStest : Conc1 : h = 1, p = 0.0068, zval = -2.7047$
 $KStest2 : Conc2 : h = 1, p = 0.0356, ks2stat = 0.4091$
 $RStest : Conc2 : h = 1, p = 0.0186, zval = -2.3536$
 $KStest2 : Conc3 : h = 0, p = 0.3320, ks2stat = 0.2727$
 $RStest : Conc3 : h = 0, p = 0.2485, zval = 1.1541$
 $KStest2 : Conc4 : h = 0, p = 0.3320, ks2stat = 0.2727$
 $RStest : Conc4 : h = 0, p = 0.9156, zval = -0.1059$

3.8 Figure 5i (Left)

Statistical significance was determined using SPSS software package (F = 12, M = 9)

p-value for gender: <0.001)

p-value for acceptance rate: 0.723

3.9 Figure 5i (Right)

Statistical significance was determined using SPSS software package (F = 12, M = 10)

p-value for gender: 0.873)

p-value for acceptance rate: 0.018

4 Figure 6

4.1 Figure 6b (Left)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 2.5457e-16.

p-value for gender: 4.7989e-01.

kstest2 results: h=0, p=8.0438e-01, ks2stat=0.1333 (overall gender difference)

Post-hoc analysis:

0.5% : $8.0114e - 01$, 2% : $8.8708e - 01$, 5% : $5.7254e - 01$, 9% : $5.0164e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : h = 0, p = 0.8848, ks2stat = 0.2333

RStest : Conc1 : h = 0, p = 0.8621, zval = 0.1736

KStest2 : Conc2 : h = 0, p = 1.0000, ks2stat = 0.1167

RStest : Conc2 : h = 0, p = 1.0000, zval = -0.0000

KStest2 : Conc3 : h = 0, p = 0.9304, ks2stat = 0.2167

RStest : Conc3 : h = 0, p = 0.9467, zval = 0.0668

KStest2 : Conc4 : h = 0, p = 0.8848, ks2stat = 0.2333

RStest : Conc4 : h = 0, p = 1.0000, zval = -0.0000

4.2 Figure 6b (Right)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 10, Male N = 10).

p-value for concentration: 6.1134e-18.

p-value for gender: 2.4633e-02.

kstest2 results: h=0, p=3.6131e-01, ks2stat=0.2000 (overall gender difference)

Post-hoc analysis:

0.5% : $8.8484e - 01$, 2% : $1.0123e - 01$, 5% : $7.5526e - 02$, 9% : $3.3494e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : h = 0, p = 0.9748, ks2stat = 0.2000

RStest : Conc1 : h = 0, p = 0.5004, zval = 0.6739

KStest2 : Conc2 : h = 0, p = 0.3129, ks2stat = 0.4000

RStest : Conc2 : h = 0, p = 0.1315, zval = 1.5083

KStest2 : Conc3 : h = 0, p = 0.6751, ks2stat = 0.3000

RStest : Conc3 : h = 0, p = 0.1233, zval = 1.5411

KStest2 : Conc4 : h = 0, p = 0.9748, ks2stat = 0.2000

RStest : Conc4 : h = 0, p = 0.4201, zval = 0.8062

4.3 Figure 6c

After Alcohol Analysis: Statistical significance was determined by Repeated measures analysis of variance. (Female N = 10, Male N = 10).

p-value for concentration: 3.374e-25.

p-value for gender: 0.13625.

kstest2 results: h=0, p=1.0000e+00, ks2stat=0.0000 (overall gender difference)

Post-hoc analysis:

0.5% : $8.1246e - 01$, 2% : $8.0739e - 01$, 5% : $1.3206e - 01$, 9% : $3.5295e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.1108, ks2stat = 0.5000$
 $RStest : Conc1 : h = 0, p = 0.5648$
 $KStest2 : Conc2 : h = 0, p = 0.1108, ks2stat = 0.5000$
 $RStest : Conc2 : h = 0, p = 0.5648$
 $KStest2 : Conc3 : h = 0, p = 0.1108, ks2stat = 0.5000$
 $RStest : Conc3 : h = 0, p = 0.7882$
 $KStest2 : Conc4 : h = 0, p = 0.1108, ks2stat = 0.5000$
 $RStest : Conc4 : h = 0, p = 0.0713$

Before and After Alcohol Analysis of Female: Statistical significance was determined by Repeated measures analysis of variance. (Before Alcohol N = 12, After Alcohol N = 10).
p-value for concentration: 6.4554e-17.
p-value for before and after alcohol: 0.11585.
kstest2 results: h=0, p=9.8992e-01, ks2stat=0.0917 (overall difference in before and after alcohol)

Post-hoc analysis:

0.5% : $6.8311e - 04$, 2% : $3.1640e - 03$, 5% : $2.8801e - 01$, 9% : $1.8980e - 05$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0017, ks2stat = 0.7500$
 $RStest : Conc1 : h = 1, p = 0.0016$
 $KStest2 : Conc2 : h = 1, p = 0.0076, ks2stat = 0.6667$
 $RStest : Conc2 : h = 1, p = 0.0059$
 $KStest2 : Conc3 : h = 1, p = 0.0282, ks2stat = 0.5833$
 $RStest : Conc3 : h = 0, p = 0.0789$
 $KStest2 : Conc4 : h = 1, p = 0.0001, ks2stat = 0.9167$
 $RStest : Conc4 : h = 1, p = 0.0002$

Before and After Alcohol Analysis of Male: Statistical significance was determined by Repeated measures analysis of variance. (Before Alcohol N = 12, After Alcohol N = 10).
p-value for concentration: 3.6982e-15.
p-value for before and after alcohol: 0.76845.
kstest2 results: h=0, p=1.0000e+00, ks2stat=0.0472 (overall difference in before and after alcohol)

Post-hoc analysis:

0.5% : $1.1779e - 01$, 2% : $9.7611e - 01$, 5% : $6.6111e - 01$, 9% : $8.3969e - 02$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.1273, ks2stat = 0.5000$
 $RStest : Conc1 : h = 0, p = 0.0884$
 $KStest2 : Conc2 : h = 0, p = 0.5732, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.9681$
 $KStest2 : Conc3 : h = 0, p = 0.2031, ks2stat = 0.4556$
 $RStest : Conc3 : h = 0, p = 0.7785$
 $KStest2 : Conc4 : h = 0, p = 0.0667, ks2stat = 0.5556$
 $RStest : Conc4 : h = 0, p = 0.0762$

4.4 Figure 6d (Left)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 5.4017e-02.

p-value for gender: 4.0729e-01.

kstest2 results: h=0, p=4.6263e-01, ks2stat=0.2016 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 7.1363e - 01, \quad 2\% : 7.6787e - 01, \quad 5\% : 7.5464e - 02, \quad 9\% : 6.6465e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 0, p = 0.2141, ks2stat = 0.5000$$

$$RStest : Conc1 : h = 0, p = 0.3462$$

$$KStest2 : Conc2 : h = 0, p = 0.3180, ks2stat = 0.5000$$

$$RStest : Conc2 : h = 0, p = 0.3701$$

$$KStest2 : Conc3 : h = 1, p = 0.0032, ks2stat = 0.7167$$

$$RStest : Conc3 : h = 1, p = 0.0192$$

$$KStest2 : Conc4 : h = 0, p = 0.2503, ks2stat = 0.4545$$

$$RStest : Conc4 : h = 0, p = 0.5360$$

4.5 Figure 6d (Right)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 10, Male N = 10).

p-value for concentration: 3.8602e-01.

p-value for gender: 7.0494e-01.

kstest2 results: h=0, p=1.7336e-01, ks2stat=0.2745 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 5.2868e - 01, \quad 2\% : 4.4209e - 01, \quad 5\% : 5.2319e - 01, \quad 9\% : 4.8906e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 1, p = 0.0204, ks2stat = 0.8000$$

$$RStest : Conc1 : h = 1, p = 0.0303$$

$$KStest2 : Conc2 : h = 0, p = 0.5070, ks2stat = 0.4250$$

$$RStest : Conc2 : h = 0, p = 0.2844$$

$$KStest2 : Conc3 : h = 0, p = 0.4892, ks2stat = 0.3556$$

$$RStest : Conc3 : h = 0, p = 0.3562$$

$$KStest2 : Conc4 : h = 0, p = 0.1076, ks2stat = 0.5417$$

$$RStest : Conc4 : h = 1, p = 0.0274$$

4.6 Figure 6e

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 4.1083e-02.

p-value for gender: 8.2510e-01.

kstest2 results: h=0, p=7.7095e-01, ks2stat=0.2000 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 8.0968e - 01, \quad 2\% : 2.1173e - 01, \quad 5\% : 4.2256e - 01, \quad 9\% : 2.2622e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 0, p = 0.6974, ks2stat = 0.4000$$

$$RStest : Conc1 : h = 0, p = 0.6429$$

$$KStest2 : Conc2 : h = 0, p = 0.2090, ks2stat = 0.6000$$

$$RStest : Conc2 : h = 0, p = 0.2063$$

$$KStest2 : Conc3 : h = 0, p = 0.6974, ks2stat = 0.4000$$

$$RStest : Conc3 : h = 0, p = 0.6349$$

$$KStest2 : Conc4 : h = 0, p = 0.6974, ks2stat = 0.4000$$

$$RStest : Conc4 : h = 0, p = 0.3016$$

4.7 Figure 6f

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 8.1732e-01.

p-value for gender: 1.2629e-01.

kstest2 results: h=1, p=8.1617e-03, ks2stat=0.5000 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 2.1755e - 01, \quad 2\% : 9.1185e - 02, \quad 5\% : 1.8516e - 01, \quad 9\% : 9.4228e - 02$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 0, p = 0.6974, ks2stat = 0.4000$$

$$RStest : Conc1 : h = 0, p = 0.2222$$

$$KStest2 : Conc2 : h = 1, p = 0.0361, ks2stat = 0.8000$$

$$RStest : Conc2 : h = 1, p = 0.0317$$

$$KStest2 : Conc3 : h = 0, p = 0.2090, ks2stat = 0.6000$$

$$RStest : Conc3 : h = 0, p = 0.1508$$

$$KStest2 : Conc4 : h = 0, p = 0.2090, ks2stat = 0.6000$$

$$RStest : Conc4 : h = 0, p = 0.0952$$

4.8 Figure 6g

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 2.6307e-07.

p-value for gender: 6.6366e-01.

kstest2 results: h=0, p=6.2161e-01, ks2stat=0.2083 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 7.5735e - 01, \quad 2\% : 3.7013e - 01, \quad 5\% : 8.0930e - 01, \quad 9\% : 6.4244e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc1 : h = 0, p = 1.0000$
 $KStest2 : Conc2 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc2 : h = 0, p = 0.3095$
 $KStest2 : Conc3 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc3 : h = 0, p = 0.8182$
 $KStest2 : Conc4 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc4 : h = 0, p = 0.5887$

4.9 Figure 6h

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 4.2378e-01.

p-value for gender: 3.0986e-01.

kstest2 results: h=1, p=9.3124e-04, ks2stat=0.5417 (overall gender difference)

Post-hoc analysis:

0.5% : 3.2581e - 01, 2% : 3.9929e - 01, 5% : 2.7256e - 01, 9% : 2.6367e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.0766, ks2stat = 0.6667$
 $RStest : Conc1 : h = 0, p = 0.1320$
 $KStest2 : Conc2 : h = 1, p = 0.0122, ks2stat = 0.8333$
 $RStest : Conc2 : h = 0, p = 0.0649$
 $KStest2 : Conc3 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc3 : h = 0, p = 0.3095$
 $KStest2 : Conc4 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc4 : h = 0, p = 0.2403$

4.10 Figure 6k

Statistical significance was determined by one-way analysis of variance. (Female BL = 10, Male BL = 10, Female Oxy = 5, Male Oxy = 5, Female Incub = 6, Male Incub = 6, Female PA = 10, Male PA = 10)

p-value for significance of difference between the groups: 1.0469e-09.

Post-hoc analysis by Tukey's HSD method:

$FemaleBLandMaleBL : 0.9981$
 $FemaleBLandFemaleOxy : 6.5947e - 06$
 $FemaleBLandFemaleIncub : 0.0132$
 $FemaleBLandFemalePA : 0.9993$
 $MaleBLandMaleOxy : 4.0017e - 06$
 $MaleBLandMaleIncub : 3.3984e - 04$
 $MaleBLandMalePA : 0.3631$

5 Supplemental Figure 1

5.1 Figure S.1i

statistical significance was determined by paired t-test using **SPSS** software package ($F = 12$, $M = 11$).
p-value for gender difference: 0.01.

6 Supplemental Figure 2

6.1 Figure S.2a

Statistical significance was determined by one-way analysis of variance. ($N = 5$).
p-value for light level: 0.0011.

6.2 Figure S.2b

Statistical significance was determined by one-way analysis of variance. ($N = 23$).
p-value for light level: 0.0028.

7 Supplemental Figure 4

7.1 Figure S.4d

statistical significance was determined by chi-squared test using **SPSS** software package ($F = 12$, $M = 11$).
p-value for Sigmoidal and U-shape for initial 1-3 months: 0.016.
p-value for Sigmoidal and U-shape after a year: 0.0009.

8 Supplemental Figure 6

8.1 Figure S.6a

Statistical significance was determined by Repeated measures analysis of variance. (Female $N = 12$, Male $N = 10$).
p-value for concentration: 4.3911e-30.
p-value for gender: 1.5870e-01.
kstest2 results: $h=0$, $p=9.3097e-01$, $ks2stat=0.1125$ (overall gender difference)

Post-hoc analysis:

$$0.5\% : 7.8880e - 01, \quad 2\% : 2.2787e - 01, \quad 5\% : 2.6929e - 01, \quad 9\% : 7.6084e - 01$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 0, p = 0.9636, ks2stat = 0.2000$$

$$RStest : Conc1 : h = 0, p = 0.5631$$

$$KStest2 : Conc2 : h = 0, p = 0.8286, ks2stat = 0.2500$$

$$RStest : Conc2 : h = 0, p = 0.2840$$

$$KStest2 : Conc3 : h = 0, p = 0.6961, ks2stat = 0.2833$$

$$RStest : Conc3 : h = 0, p = 0.2892$$

$$KStest2 : Conc4 : h = 0, p = 0.8848, ks2stat = 0.2333$$

$$RStest : Conc4 : h = 0, p = 0.8391$$

8.2 Figure S.6b

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 2.6791e-06.

p-value for gender: 1.6014e-04.

kstest2 results: h=1, p=1.3139e-08, ks2stat=0.6375 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 4.0070e - 04, \quad 2\% : 1.4910e - 05, \quad 5\% : 9.2678e - 03, \quad 9\% : 2.5464e - 03$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 1, p = 0.0076, ks2stat = 0.6667$$

$$RStest : Conc1 : h = 1, p = 0.0033$$

$$KStest2 : Conc2 : h = 1, p = 0.0001, ks2stat = 0.9167$$

$$RStest : Conc2 : h = 1, p = 0.0003$$

$$KStest2 : Conc3 : h = 0, p = 0.0567, ks2stat = 0.5333$$

$$RStest : Conc3 : h = 1, p = 0.0229$$

$$KStest2 : Conc4 : h = 1, p = 0.0076, ks2stat = 0.6667$$

$$RStest : Conc4 : h = 1, p = 0.0051$$

8.3 Figure S.6c

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 1.1616e-02.

p-value for gender: 7.3997e-03.

kstest2 results: h=1, p=1.8518e-05, ks2stat=0.5000 (overall gender difference)

Post-hoc analysis:

$$0.5\% : 1.8703e - 02, \quad 2\% : 4.5455e - 04, \quad 5\% : 2.3150e - 01, \quad 9\% : 1.1937e - 02$$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$$KStest2 : Conc1 : h = 1, p = 0.0076, ks2stat = 0.6667$$

$$RStest : Conc1 : h = 1, p = 0.0111$$

$$KStest2 : Conc2 : h = 1, p = 0.0003, ks2stat = 0.8333$$

$$RStest : Conc2 : h = 1, p = 0.0014$$

$$KStest2 : Conc3 : h = 0, p = 0.2270, ks2stat = 0.4167$$

$$RStest : Conc3 : h = 0, p = 0.4098$$

$$KStest2 : Conc4 : h = 1, p = 0.0452, ks2stat = 0.5500$$

$$RStest : Conc4 : h = 1, p = 0.0092$$

8.4 Figure S.6d

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 7.6444e-13.

p-value for gender: 2.4455e-02.

kstest2 results: h=1, p=6.8336e-03, ks2stat=0.3500 (overall gender difference)

Post-hoc analysis:

0.5% : $1.1326e - 02$, 2% : $1.2677e - 04$, 5% : $9.3261e - 01$, 9% : $1.6538e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : $h = 1, p = 0.0358, ks2stat = 0.5667$

RStest : Conc1 : $h = 1, p = 0.0111$

KStest2 : Conc2 : $h = 1, p = 0.0001, ks2stat = 0.9167$

RStest : Conc2 : $h = 1, p = 0.0003$

KStest2 : Conc3 : $h = 0, p = 0.9989, ks2stat = 0.1500$

RStest : Conc3 : $h = 0, p = 0.9212$

KStest2 : Conc4 : $h = 0, p = 0.5564, ks2stat = 0.3167$

RStest : Conc4 : $h = 0, p = 0.2485$

8.5 Figure S.6e

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: $2.3011e-15$.

p-value for gender: $3.8916e-01$.

kstest2 results: $h=0, p=2.6677e-01, ks2stat=0.2083$ (overall gender difference)

Post-hoc analysis:

0.5% : $2.8309e - 01$, 2% : $1.3761e - 02$, 5% : $8.4288e - 01$, 9% : $5.6653e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

KStest2 : Conc1 : $h = 0, p = 0.4896, ks2stat = 0.3333$

RStest : Conc1 : $h = 0, p = 0.3891$

KStest2 : Conc2 : $h = 0, p = 0.0567, ks2stat = 0.5333$

RStest : Conc2 : $h = 1, p = 0.0149$

KStest2 : Conc3 : $h = 0, p = 0.9636, ks2stat = 0.2000$

RStest : Conc3 : $h = 0, p = 0.8940$

KStest2 : Conc4 : $h = 0, p = 0.3689, ks2stat = 0.3667$

RStest : Conc4 : $h = 0, p = 0.5716$

9 Supplemental Figure 7

9.1 Figure S.7a (Left)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: $1.4050e-01$.

p-value for gender: $3.6754e-01$.

kstest2 results: $h=0, p=1.2139e-01, ks2stat=0.2458$ (overall gender difference)

Post-hoc analysis:

0.5% : $7.7450e - 01$, 2% : $9.1269e - 01$, 5% : $3.6064e - 03$, 9% : $5.6181e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.9304, ks2stat = 0.2167$
 $RStest : Conc1 : h = 0, p = 0.7667$
 $KStest2 : Conc2 : h = 0, p = 0.8286, ks2stat = 0.2500$
 $RStest : Conc2 : h = 0, p = 0.7667$
 $KStest2 : Conc3 : h = 1, p = 0.0101, ks2stat = 0.6500$
 $RStest : Conc3 : h = 1, p = 0.0111$
 $KStest2 : Conc4 : h = 0, p = 0.1072, ks2stat = 0.4833$
 $RStest : Conc4 : h = 0, p = 0.3734$

9.2 Figure S.7a (Right)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 10, Male N = 10).

p-value for concentration: 6.8275e-01.

p-value for gender: 1.9427e-01.

kstest2 results: h=0, p=1.3925e-01, ks2stat=0.2500 (overall gender difference)

Post-hoc analysis:

0.5% : $4.4173e - 01$, 2% : $1.5443e - 01$, 5% : $4.3851e - 01$, 9% : $8.8666e - 02$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.3129, ks2stat = 0.4000$
 $RStest : Conc1 : h = 0, p = 0.3075$
 $KStest2 : Conc2 : h = 0, p = 0.6751, ks2stat = 0.3000$
 $RStest : Conc2 : h = 0, p = 0.1620$
 $KStest2 : Conc3 : h = 0, p = 0.6751, ks2stat = 0.3000$
 $RStest : Conc3 : h = 0, p = 0.9698$
 $KStest2 : Conc4 : h = 0, p = 0.1108, ks2stat = 0.5000$
 $RStest : Conc4 : h = 0, p = 0.0890$

9.3 Figure S.7b (Left)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 12, Male N = 10).

p-value for concentration: 7.3371e-02.

p-value for gender: 4.0968e-04.

kstest2 results: h=1, p=2.4759e-06, ks2stat=0.5417 (overall gender difference)

Post-hoc analysis:

0.5% : $1.2714e - 02$, 2% : $5.9126e - 02$, 5% : $8.5187e - 04$, 9% : $1.1225e - 03$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0220, ks2stat = 0.6000$
 $RStest : Conc1 : h = 1, p = 0.0092$
 $KStest2 : Conc2 : h = 0, p = 0.0567, ks2stat = 0.5333$
 $RStest : Conc2 : h = 0, p = 0.0806$
 $KStest2 : Conc3 : h = 1, p = 0.0076, ks2stat = 0.6667$
 $RStest : Conc3 : h = 1, p = 0.0022$
 $KStest2 : Conc4 : h = 1, p = 0.0076, ks2stat = 0.6667$
 $RStest : Conc4 : h = 1, p = 0.0041$

9.4 Figure S.7b (Right)

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 10, Male N = 10).

p-value for concentration: 5.5161e-01.

p-value for gender: 4.4971e-02.

kstest2 results: h=1, p=1.0793e-02, ks2stat=0.3500 (overall gender difference)

Post-hoc analysis:

0.5% : $9.3255e - 02$, 2% : $8.9902e - 02$, 5% : $5.4953e - 01$, 9% : $6.7305e - 03$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0310, ks2stat = 0.6000$
 $RStest : Conc1 : h = 0, p = 0.0890$
 $KStest2 : Conc2 : h = 0, p = 0.3129, ks2stat = 0.4000$
 $RStest : Conc2 : h = 0, p = 0.1212$
 $KStest2 : Conc3 : h = 0, p = 0.3129, ks2stat = 0.4000$
 $RStest : Conc3 : h = 0, p = 0.6232$
 $KStest2 : Conc4 : h = 1, p = 0.0069, ks2stat = 0.7000$
 $RStest : Conc4 : h = 1, p = 0.0113$

9.5 Figure S.7d

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 8.7784e-02.

p-value for gender: 3.1525e-02.

kstest2 results: h=1, p=2.3213e-02, ks2stat=0.4500 (overall gender difference)

Post-hoc analysis:

0.5% : $3.1246e - 03$, 2% : $5.8901e - 01$, 5% : $3.6169e - 02$, 9% : $9.2633e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 1, p = 0.0038, ks2stat = 1.0000$
 $RStest : Conc1 : h = 1, p = 0.0079$
 $KStest2 : Conc2 : h = 0, p = 0.9996, ks2stat = 0.2000$
 $RStest : Conc2 : h = 0, p = 0.6905$
 $KStest2 : Conc3 : h = 1, p = 0.0361, ks2stat = 0.8000$
 $RStest : Conc3 : h = 0, p = 0.0556$
 $KStest2 : Conc4 : h = 0, p = 0.6974, ks2stat = 0.4000$
 $RStest : Conc4 : h = 0, p = 1.0000$

9.6 Figure S.7e

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 6.8473e-01.

p-value for gender: 1.5292e-01.

kstest2 results: h=1, p=2.3213e-02, ks2stat=0.4500 (overall gender difference)

Post-hoc analysis:

0.5% : 3.2600e - 01, 2% : 1.2515e - 01, 5% : 3.5598e - 01, 9% : 1.0995e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc1 : h = 0, p = 0.3095$
 $KStest2 : Conc2 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc2 : h = 0, p = 0.2222$
 $KStest2 : Conc3 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc3 : h = 0, p = 0.3968$
 $KStest2 : Conc4 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc4 : h = 0, p = 0.2222$

9.7 Figure S.7f

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 6.2169e-01.

p-value for gender: 8.7263e-02.

kstest2 results: h=1, p=8.1617e-03, ks2stat=0.5000 (overall gender difference)

Post-hoc analysis:

0.5% : 1.9249e - 01, 2% : 1.3288e - 01, 5% : 5.9820e - 02, 9% : 5.9578e - 02

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.6974, ks2stat = 0.4000$
 $RStest : Conc1 : h = 0, p = 0.2222$
 $KStest2 : Conc2 : h = 0, p = 0.6974, ks2stat = 0.4000$
 $RStest : Conc2 : h = 0, p = 0.2222$
 $KStest2 : Conc3 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc3 : h = 0, p = 0.0952$
 $KStest2 : Conc4 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc4 : h = 0, p = 0.0952$

9.8 Figure S.7g

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 5, Male N = 5).

p-value for concentration: 4.5703e-01.

p-value for gender: 6.3326e-02.

kstest2 results: h=1, p=7.2529e-04, ks2stat=0.6000 (overall gender difference)

Post-hoc analysis:

0.5% : $1.7344e - 01$, 2% : $9.6526e - 02$, 5% : $5.1225e - 02$, 9% : $4.6853e - 02$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc1 : h = 0, p = 0.2222$
 $KStest2 : Conc2 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc2 : h = 0, p = 0.0952$
 $KStest2 : Conc3 : h = 1, p = 0.0361, ks2stat = 0.8000$
 $RStest : Conc3 : h = 0, p = 0.0556$
 $KStest2 : Conc4 : h = 0, p = 0.2090, ks2stat = 0.6000$
 $RStest : Conc4 : h = 0, p = 0.0952$

9.9 Figure S.7h

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 3.0232e-04.

p-value for gender: 4.9003e-01.

kstest2 results: h=0, p=8.6076e-01, ks2stat=0.1667 (overall gender difference)

Post-hoc analysis:

0.5% : $6.9253e - 01$, 2% : $6.3994e - 01$, 5% : $6.0961e - 01$, 9% : $7.4470e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc1 : h = 0, p = 0.5887$
 $KStest2 : Conc2 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.6991$
 $KStest2 : Conc3 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc3 : h = 0, p = 0.4848$
 $KStest2 : Conc4 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc4 : h = 0, p = 0.8182$

9.10 Figure S.7i

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 1.1029e-03.

p-value for gender: 3.4076e-01.

kstest2 results: h=0, p=5.0588e-02, ks2stat=0.3750 (overall gender difference)

Post-hoc analysis:

0.5% : 5.1260e - 01, 2% : 3.1392e - 01, 5% : 2.5092e - 01, 9% : 3.6439e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 1.0000, ks2stat = 0.1667$
 $RStest : Conc1 : h = 0, p = 0.8182$
 $KStest2 : Conc2 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc2 : h = 0, p = 0.2403$
 $KStest2 : Conc3 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc3 : h = 0, p = 0.1797$
 $KStest2 : Conc4 : h = 1, p = 0.0122, ks2stat = 0.8333$
 $RStest : Conc4 : h = 1, p = 0.0411$

9.11 Figure S.7j

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 1.8477e-06.

p-value for gender: 3.2473e-01.

kstest2 results: h=0, p=2.1598e-01, ks2stat=0.2917 (overall gender difference)

Post-hoc analysis:

0.5% : 4.5320e - 01, 2% : 4.5178e - 01, 5% : 3.0428e - 01, 9% : 2.1404e - 01

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc1 : h = 0, p = 0.4848$
 $KStest2 : Conc2 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.4848$
 $KStest2 : Conc3 : h = 0, p = 0.0766, ks2stat = 0.6667$
 $RStest : Conc3 : h = 0, p = 0.3939$
 $KStest2 : Conc4 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc4 : h = 0, p = 0.2403$

9.12 Figure S.7k

Statistical significance was determined by Repeated measures analysis of variance. (Female N = 6, Male N = 6).

p-value for concentration: 5.9005e-04.

p-value for gender: 1.4048e-01.

kstest2 results: h=0, p=2.1598e-01, ks2stat=0.2917 (overall gender difference)

Post-hoc analysis:

0.5% : $3.2825e - 01$, 2% : $4.1626e - 01$, 5% : $5.1375e - 02$, 9% : $3.6430e - 01$

KStest2 and Wilcoxon rank sum test Results (complementary to post-hoc analysis)

$KStest2 : Conc1 : h = 0, p = 0.3180, ks2stat = 0.5000$
 $RStest : Conc1 : h = 0, p = 0.3939$
 $KStest2 : Conc2 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc2 : h = 0, p = 0.5887$
 $KStest2 : Conc3 : h = 0, p = 0.0766, ks2stat = 0.6667$
 $RStest : Conc3 : h = 0, p = 0.0649$
 $KStest2 : Conc4 : h = 0, p = 0.8096, ks2stat = 0.3333$
 $RStest : Conc4 : h = 0, p = 0.4848$