

Distribution of Entry Ratios in Sequences of Powers of Transition Matrices

Ali Taqi

2/4/2021

Notation

Random Batches

Let \mathbb{F} be a field, and fix some $M \in \mathbb{N}$. Let $\mathcal{B}_\lambda \subset \mathbb{F}^M$ be a uniformly random batch of points in \mathbb{F}^M in the M -hypercube of length λ . That is,

$$\mathcal{B}_\lambda = \{\vec{x} \mid x_i \sim \text{Unif}(-\lambda, \lambda) \text{ for } i = 1, \dots, M\}$$

Note: If $\mathbb{F} = \mathbb{C}$, then take $\vec{x} \in \mathcal{B}_\lambda$ to mean $\vec{x} = a + bi$ where $a, b \sim \text{Unif}(-\lambda, \lambda)$.

Finite Evolution Sequences

Suppose we sample a random point from \mathcal{B}_λ to emulate a random point $\vec{v} \in \mathbb{F}^M$. Additionally, let $Q \in \mathbb{F}^{M \times M}$ be a transition matrix over \mathbb{F} . Fixing a maximum power (‘time’) $T \in \mathbb{N}$, define the evolution sequence of \vec{v} as follows:

$$\mathcal{S}(v, Q, T) = (\alpha_n)_{n=1}^T \text{ where } \alpha_k = vQ^k$$

If we do not impose a finiteness constraint on the sequence, we consider powers for $n \in \mathbb{N}$ or $t = \infty$

Consecutive Ratio Sequences

Accordingly, define the consecutive ratio sequence (CST) of \vec{v} as follows:

$$\mathcal{R}(v, Q, T) = (r_n)_{n=2}^T \text{ where } (r_n)_j = \frac{(\alpha_n)_j}{(\alpha_{n-1})_j} \text{ for } j = 1, \dots, M$$

In other words, the consecutive ratio sequence of v can be obtained by performing **component-wise division** on consecutive elements of the evolution sequence of v .

Near Convergence

Because these sequences may never truly converge to eigenvectors of the matrix, we formalize a notion of “near convergence”. As a preliminary, we first define ε -equivalence. Let \mathbb{F} be a field, and fix $\varepsilon \in \mathbb{R}^+$. Suppose we have vectors $v, v' \in \mathbb{F}^M$. Then, $v \sim_\varepsilon v'$ if $\|v - v'\| < \varepsilon$ where $\|\cdot\|$ is the norm on \mathbb{F} .

Let $\varepsilon \in \mathbb{R}^+$, and suppose we have an evolution sequence $(a[\vec{v}])_n$. Then, a_n ε -converges at N if:

$$\exists N \in \mathbb{N} : \forall n \geq N \mid a_N \sim_\varepsilon a_n$$

Questions

1. How are the entries of the CRS distributed? Are they normal, and if so, what is its mean?
2. Are the entries of the CRS i.i.d as $t \rightarrow \infty$?
3. For an Erdos-Renyi matrix, is the mixing time t dependent on the parameter p ?
4. What impact does the running time parameter T have on σ (the variance of the distribution of the CRS entries)?

Initial Findings

It seems to be the case that the **log-transformed** entries of the CRS are normally distributed about $\log \lambda_1$ where $\lambda_1 = \max(\sigma(Q))$, the largest eigenvalue of Q . That is,

$$r_i \sim \mathcal{N}(\ln \lambda_1, \sigma) \text{ for } i = 1, \dots, M$$