

HOMEWORK 1

A. BOLSHAKOV

1. PROBLEM 1

Many constraint surfaces can be locally linearized. Let us examine the behavior of the dynamical system

$$\dot{\mathbf{x}} = P_A(\mathbf{x}) - P_B(\mathbf{x})$$

for this large class of systems by just looking at a simple, 2 dimensional case. Let constraint A be $x_2 = mx_1$ and constraint B be $x_2 = -mx_1$ for some slope m . (see Fig. ??).

The system is simple enough that we can explicitly solve for P_A and P_B :

$$P_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \frac{1}{m^2 + 1} \begin{pmatrix} x_1 + mx_2 \\ mx_1 + m^2 x_2 \end{pmatrix}$$
$$P_B \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \frac{1}{m^2 + 1} \begin{pmatrix} x_1 - mx_2 \\ m^2 x_2 - mx_1 \end{pmatrix}$$

We can ignore the $m^2 + 1$ coefficient as we only care about long term behavior. So, we get

$$\dot{\mathbf{x}} = P_A(\mathbf{x}) - P_B(\mathbf{x}) = \begin{pmatrix} 2mx_2 \\ 2mx_1 \end{pmatrix} = \begin{pmatrix} 0 & 2m \\ 2m & 0 \end{pmatrix} \mathbf{x}$$

This is a classic linear hyperbolic system. The two eigenvalues are $\pm 2m$, so all trajectories except a set of measure 0 will flow away.

2. PROBLEM 2

3. PROBLEM 3