

# Pacific saury *Cololabis saira* report

Andrew Chin

6/22/2021

## **LH, distribution, stock structure**

Additional summaries of Pacific saury ecology and fisheries can be found in Yatsu (2019) and in NPFC Report SSC PS06, Annex H (2020).

Pacific saury (*Cololabis saira* Brevoort, 1856) occur throughout the subarctic and subtropical North Pacific Ocean from inshore waters of Japan and Kuril Islands to eastward to Gulf of Alaska and southward to Mexico. Pacific saury is an especially important commercial fish in the Western North Pacific Ocean (Parin 1968; Hubbs and Wisner 1980).

Pacific saury have a two-year life cycle, migrating between summer feeding grounds in the Oyashio Current, the Kuroshio Current, the Kuroshio-Oyashio Transition Zone (KOTZ), and the Japan Sea (Yatsu 2019) and the breeding grounds in the Kuroshio Current off southern Japan (Fukushima 1979; Kosaka 2000). Spawning occurs over a broad spatial and temporal period, during the winter in Kuroshio and subtropical areas and the spring/summer in the KOTZ (Watanabe and Lo 1989). Pacific saury in offshore regions (east of 160E) also migrate westward toward the coast of Japan after October every year (Suyama et al. 2012). There is little genetic evidence for distinct stock structure (Chow et al. 2009), so Pacific saury is managed as one stock.

Pacific saury mature about 8 months after hatching, at around 22cm (Suyama et al. 2016), though there is growth variation depending where larvae have hatched. Age-1 fish grow to 27cm by June and July when Japanese fishery-independent surveys are conducted, and to 30cm when fishery occurs between August and December (Suyama et al. 2006).

## **Fishery**

Pacific saury are managed by the North Pacific Fisheries Commission (NPFC) by its six of its eight member countries: China, Japan, Korea, Russia, Chinese Taipei, and Vanuatu. The two other member countries, Canada and the United States, do not have a sustained commercial history with Pacific saury and are rarely caught as bycatch.

Japan was one of the first countries to develop its commercial industrialized fishery for Pacific saury in the 1940's using stick-held dip nets. Stick-held dip nets are the dominant gear for capturing Pacific saury by all countries. Russian and Korean fisheries were subsequently began in the 1950's, and Chinese Taipei commercial fishery in 1975, some of which concurrently fished squid but then switched over to Pacific saury only in 1990s. China and Vanuatu commenced their fisheries in 2003 and 2004, respectively. China has rapidly developed their fisheries capacity in 2012.


## **Catch and Stock Status**

Historical Japanese data extends to the 1950s, CPUE data by country from the 1980s, and summary statistics have been reported to the NPFC from 1995 to present. CPUE is standardized for all countries using a

generalized linear model, and CPUE from Japan (early and late datasets differentiated), China, Chinese Taipei, Korea, and Russia are used in stock assessment models. Japanese and Russian catches mainly occur within their respective Exclusive Economic Zones (EEZs), which China, Chinese Taipei, and Vanuatu fisheries mostly occur within the NPFC high seas Convention Area (CA). Catches in the CA have been increasing, and are largely driven by China and Chinese Taipei (Figure...). An increasing proportion of catch is occurring in the NPFC CA waters as well (Figure ...).


Pacific saury were not managed intensively until a 2015 NPFC resolution to assess the stock was established, and the first stock assessments in 2019 and 2020 were conducted using a Bayesian state-space surplus production model. The ratio between estimated biomass and biomass required for Maximum Sustainable Yield (B:B<sub>msy</sub>) was 0.46 (80%CI 0.30-0.67), indicating the stock was overfished (North Pacific Fisheries Commission 2021a). Examining the CPUE used in the stock assessment models, CPUE has been declining across all countries (Figure ...).

In 2019, a Total Allowable Catch (TAC) limit of 330,000 metric tons in all areas was established for 2020 (North Pacific Fisheries Commission 2019). Total TAC was raised slightly for 2021 and 2022 to 333,750 tons in all areas, and 198,000 metric tons in the high-seas CA (North Pacific Fisheries Commission 2021b). These new TACs are a 52.9% and 56.2% reduction from the maximum catch from the total areas and NPFC convention area, respectively. However, the 2021-2022 TAC is only a 89.2% reduction from the average total catch.



Total Pacific saury catch reported to the NPFC. Black dashed line indicates 333,750 metric ton TAC instituted for 2021 and 2022.


### Pacific saury catch in NPFC high seas convention area



Data from <https://www.npfc.int/statistics>.


Reported Pacific saury catches in the NPFC high seas convention area. The black dashed line indicates 198,000 metric ton TAC instituted for the high seas in 2021 and 2022.

## Comparison between national waters and NPFC convention area



Data from <https://www.npfc.int/statistics>.


Comparison of Pacific saury catch within National Waters (NW, within 200 nautical miles of land) and in the high seas NPFC Convention Area (CA). Black dashed line indicates 333,750 metric ton TAC instituted for 2021 and 2022.



Time series of standardized CPUE indices used by the NPFC to estimate Pacific saury stock abundance and status. X's and O's in the Japan figure represent catch data from “early” and “late” catch record periods.

## Overlap with IYS 2022 expedition

Zone 1 of the IYS Expedition overlaps strongly with the summer feeding grounds and autumn fishing grounds of Pacific saury. In addition, coho salmon are known to be predators of saury as well. Changes in the Kuroshio and the KOTZ could influence production or interactions between salmon and Pacific saury.



Schematic diagram of life history of Pacific saury and ocean circulations in the northwestern Pacific. Solid lines indicate dominant current systems and dashed lines the migration routes. from Ito et al. 2004.




Figure ... Planned sampling zones for the IYS 2022 Pan-Pacific Winter Expedition.

## References

- Chow S., Suzuki N., Brodeur R.D., Ueno Y. 2009. Little population structuring and recent evolution of the Pacific saury (*Cololabis saira*) as indicated by mitochondrial and nuclear DNA sequence data. *J Exp Mar Biol Ecol* 369:17–21.
- Fukushima S. 1979. Synoptic analysis of migration and fishing conditions of saury in northwest Pacific Ocean. *Bull. Tohoku Reg. Fish. Res. Lab.* 41, 1-70.
- Hubbs C.L., Wisner R.L. 1980. Revision of the sauries (Pisces, Scomberesocidae) with descriptions of two new genera and one new species. *Fish Bull US* 77: 521–566.
- Kosaka S. 2000. Life history of the Pacific saury *Cololabis saira* in the northwest Pacific and considerations on resource fluctuations based on it. *Bulletin of Tohoku National Fisheries Research Institute* 63: 1–96.
- North Pacific Fisheries Commission. 2019. Conservation and Management Measure for Pacific Saury. CMM 2019-08. 2 pp. (Available at [www.npfc.int](http://www.npfc.int))
- North Pacific Fisheries Commission. 2021a. North Pacific Ocean Pacific Saury 2019 Stock Assessment Update Report. NPFC-2021-SCsm01-WP03. (Available at [www.npfc.int](http://www.npfc.int)).
- North Pacific Fisheries Commission. 2021b. Conservation and Management Measure for Pacific Saury. CMM 2021-08. 3 pp. (Available at [www.npfc.int](http://www.npfc.int))
- Parin N.V. 1968. Scomberesocidae (Pisces, Synentognathi) of the eastern Atlantic Ocean. *Atlantide Rep.* 10: 275- 290.
- Small Scientific Committee on Pacific Saury. 2020. 6th Meeting Report. NPFC-2020-SSC PS06-Final Report. 68 pp. (Available at [www.npfc.int](http://www.npfc.int))
- Suyama S., Kurita Y., Ueno Y. 2006. Age structure of Pacific saury *Cololabis saira* based on observations of the hyaline zones in the otolith and length frequency distributions. *Fish Sci* 72: 742–749.
- Suyama S., Nakagami M., Naya M., Ueno Y. 2012. Migration route of Pacific saury *Cololabis saira* inferred from the otolith hyaline zone. *Fisheries Science* 78(6): 1179-1186.
- Suyama S., Shimizu A., Isu S., Ozawa H., Morioka T., Nakaya M., Nakagawa T., · Murakami N., Ichikawa T., Ueno Y. 2016. Determination of the spawning history of Pacific saury *Cololabis saira* from rearing experiments: identification of post-spawning fish from histological observations of ovarian arterioles. *Fisheries Science* 82(3): 445-457.
- Watanabe Y., Lo N.C.H. 1989. Larval production and mortality of Pacific saury, *Cololabis saira*, in the northwestern Pacific Ocean. *Fish Bull US* 87: 601–613.
- Yatsu A. 2019. Review of population dynamics and management of small pelagic fishes around the Japanese Archipelago. *Fish Science*. 85: 611-639.