# Introduction to Databases Problem Set #2

### Si Kai Lee s13950@columbia.edu

February 29, 2016

## Problem 2

#### $\mathbf{a}$

```
Let liquor be L and sales be S.

Let \rho(C(1->\text{lid}1,7->\text{lid}2),L\times S) be X.

Names = \pi_{\text{name}}(\sigma_{\substack{\text{lid}1 = \text{lid}2\\ \text{month} = \text{December}\\ \text{county} = \text{Polk}\\ \text{quantity} >= 1}}(X)) \cap \pi_{\text{name}}(\sigma_{\substack{\text{lid}1 = \text{lid}2\\ \text{month} = \text{December}\\ \text{county} = \text{Linn}\\ \text{quantity} >= 1}}(X))
```

#### b

Let liquor be L and sales be S.

Let  $L \bowtie S$  be X.

$$\begin{aligned} \text{Manufacturers} &= \pi_{\text{manufacturer}} \sigma_{1 \neq 9} ((\sigma_{\text{month} = \text{January}}(X) \bowtie_{\text{manufacturer}} \sigma_{\text{month} = \text{January}}(X))) \\ & \text{county} &= \text{Polk} \\ & \text{quantity} >= 1 \end{aligned}$$

### Problem 3

 $\mathbf{a}$ 

B D

x c

y a

x a

Si Kai Lee sl3950

### $\mathbf{b}$

A В D 1 1 Х  $\mathbf{c}$ 3 1 у a 3 1  $\mathbf{X}$ a 1 3  $\mathbf{X}$  $\mathbf{c}$ 3 3 a 3 3 a

### $\mathbf{c}$

### $\mathbf{d}$

Since T1 as T1 - T2 = T1, hence T1- (T1 - T2) results in an empty table.

#### $\mathbf{e}$

Since T1 does not have a column D, there would not be a common column to natural join, hence an empty table

Si Kai Lee sl3950 2