




POLITECNICO
MILANO 1863

**SCUOLA DI INGEGNERIA INDUSTRIALE
E DELL'INFORMAZIONE**



The Earth's Bow Shock

LAUREA MAGISTRALE IN AERONAUTICAL ENGINEERING - INGEGNERIA AERONAUTICA

Author: ALESSANDRO TEDESCO - 10801442

Advisor: PROF. GIULIO GORI

Academic year: 2025-2026

1. Introduction

The Earth's bow shock is the most famous example of a plasma shock. It is a standing magnetohydrodynamic shock that consists of an axisymmetric, parabolically shaped surface that develops around the Sun-Earth line as a result of the interaction of the Earth's magnetosphere with the hypersonic plasma flow coming from the Sun, known as the solar wind. Indeed, the magnetosphere acts as a blunt obstacle at rest, which decelerates the solar wind flow. It forms on the dayside and extends for approximately 13 to 14 Earth radii outside the planet [13], dividing the incoming solar wind into two regions: the undisturbed plasma flow upstream and the disturbed magnetosheath flow downstream.

The bow shock acts as a discontinuity: even though it is characterized by a finite thickness, this is negligible compared to the characteristic length scale of the problem. As the velocity suddenly decreases across the shock, the density and magnetic field intensity increase by up to a factor of four, as shown by experimental measurements [13, 15, 16]. Clearly, both the intensity and the shape of the bow shock are strongly influenced by solar activity, which drives variations in the solar wind.

In conventional shocks (collisional shocks), the dissipation mechanism is provided by the viscosity due to collisions between molecules, which redistributes energy among all the particles. How-

ever, a viscosity mechanism cannot occur in a rarefied plasma, where collisions between particles are essentially negligible. Instead, wave-particle interactions are responsible for the exchange of momentum, resulting in a collisionless shock. Nonetheless, the concepts regarding the fundamental nature of such shocks are the same as in a gas-dynamic shock, and so are the basic conservation laws.

Hence, the main purpose of this report is to analyze the fluid-dynamic properties and dissipation mechanisms of these collisionless shocks.

2. Solar Wind

The solar wind is a continuous flow of hypersonic ionized solar plasma, carrying a remnant of the solar magnetic field that pervades the entire heliosphere¹. The plasma is driven outward by the huge pressure difference between the solar corona and interplanetary space, which overcomes the restraining influence of solar gravity. Its existence was first proposed in the 1950s based on the evidence that small variations in Earth's geomagnetic behavior were related to solar activity, as well as on theoretical models for the equilibrium state of the solar corona. It was then experimentally confirmed in the early 1960s by several missions such as Luna-1, -2, and -3, Mariner-2 and American Explorer-10 [14].

¹The sphere of influence of the Sun in interplanetary space.

2.1. Properties

The most extensive and detailed observations of the solar wind have been made from spacecraft near Earth's orbit [12]. The main properties of the plasma and magnetic field at this distance (1 Astronomical Unit, or AU) are summarized in Table 1. This plasma stream consists largely of protons and electrons in nearly equal numbers, with a minor admixture (a few percent) of ionized helium He^{2+} and heavier elements in traces [5].

It is important to note that these properties are highly dynamic and depend strongly on the solar cycle and solar activity; moreover, they evolve significantly as the flow expands from the Sun's atmosphere to our planet.

Property	Value
Proton Density	6.6 cm^{-3}
Electron Density	7.1 cm^{-3}
He^{2+} Density	0.25 cm^{-3}
Average Flow Speed	450 km/s
Proton Temperature	$1.2 \cdot 10^5 \text{ K}$
Electron Temperature	$1.4 \cdot 10^5 \text{ K}$
Magnetic Induction	7 nT

Table 1: Observed properties of solar wind near the orbit of the Earth [11].

The solar wind is categorized into two distinct states — the fast and the slow wind — which differ primarily in their origin, physical properties, and stability. The fast solar wind originates in the coronal holes, which are cooler, darker regions of the corona dominated by open magnetic field lines. Fast wind streams are generally stable over extended time periods (spanning several solar rotations), and differences between successive streams are usually small. At 1 AU, the fast wind is characterized by flow speeds between 400 km/s and 800 km/s . Conversely, the slow solar wind exhibits lower speeds, ranging from 250 km/s to 400 km/s [11].

An intrinsic feature of the solar wind is the Interplanetary Magnetic Field (IMF) it carries. Because the solar wind is a highly electrically conductive plasma, magnetic diffusion is negligible compared to advection. Consequently, there is no relative motion between the magnetic field

lines and the plasma flow; the magnetic field is effectively "frozen-in" to the plasma [10]. As the plasma expands radially outward, it drags the magnetic field lines with it. Combined with the Sun's rotation, this radial expansion winds the field lines into an Archimedean spiral [12]. At 1 AU, this results in the IMF being oriented at an angle of approximately 45° relative to the Sun-Earth line.

3. Plasma Waves

A wave is a disturbance propagating through a continuous medium, resulting in a periodic motion of the fluid properties. In a magnetized plasma, wave propagation depends on the orientation of the wave vector $\mathbf{k}$ with respect to the undisturbed background magnetic field $\mathbf{B}_0$ [11]. The magnetohydrodynamic (MHD) governing equations form a set of coupled non-linear partial differential equations. By considering small perturbations, these equations can be linearized to analyze plane wave solutions of the form:

$$\mathbf{B}(\mathbf{r}, t) = \mathbf{B}_0 \exp i(\mathbf{k} \cdot \mathbf{r} - \omega t),$$

described by the angular frequency ω and the wave vector $\mathbf{k}$. The phase velocity of a wave front is defined as $v_{ph} = \omega/k$, with k being the wavenumber.

Because a sinusoidal wave extends indefinitely in space and cannot transport information, physical disturbances are described by wave packets moving at the group velocity $v_g = \partial\omega/\partial k$, which is determined from the dispersion relation between the frequency and the wavenumber, $\omega = \omega(k)$. The group velocity is always bounded by the speed of light, and it represents the propagation speed of wave energy and information through the medium.

3.1. Alfvén Waves

Alfvén waves are transverse waves propagating parallel to the magnetic field ($\mathbf{k} \parallel \mathbf{B}$). Their motion is similar to that of a plucked string: magnetic tension acts as a restoring force, and a transverse wave propagates along the field line. The fluctuating quantities are the electromagnetic field and the current density.

Furthermore, Alfvén waves are non-dispersive waves: phase and group velocities are the same. Indeed, the dispersion relation $\omega_A = \pm k_{\parallel} v_A$ is

linear. The Alfvén speed is defined as [7]

$$v_A = \frac{B_0}{\sqrt{\mu_0 \rho_0}} \quad (1)$$

This is a crucial parameter for plasmas, representing the maximum disturbance speed along the magnetic field, playing an analogous role to the sound speed in a gas. In hypersonic space flows, the ratio of the flux speed u and the Alfvén speed defines the *Alfvénic Mach number* ($M_A = u/v_A$). For $M_A > 1$, a shock wave forms to decelerate the plasma. Typical Alfvén speeds range from tens of km/s in the interplanetary medium to hundreds of km/s in the solar atmosphere.

3.2. Magnetosonic Waves

Magnetosonic waves are compressive waves that involve both thermal and magnetic pressure as restoring forces. Unlike Alfvén waves, they can propagate across the magnetic field. For a propagation perpendicular to the field, the two restoring forces act together, and the wave travels at the magnetosonic speed, which is determined by the sound and Alfvén speeds (1):

$$v_{ms}^2 = \frac{\omega^2}{k^2} = v_s^2 + v_A^2, \quad (2)$$

which is independent of the wavenumber k , meaning that the wave is non-dispersive. Moreover, as the magnetic field vanishes, the Alfvén speed approaches zero, and the magnetosonic phase speed converges to the acoustic sound speed.

3.3. Ion Acoustic Waves

While high-frequency electron plasma oscillations assume stationary ions due to their high inertia [11], ion motion becomes significant at lower frequencies. In ordinary fluids, sound waves represent the counterparts of ion waves. However, in collisionless plasmas, unlike neutral fluids driven by binary collisions, the restoring force for ion waves is provided by electron thermal pressure. As light, energetic electrons expand ahead of the heavier ions, they set up a microscopic charge separation that generates an electric field. This drags the ions along, whose mass provides the flow inertia.

Ion waves are non-dispersive, with equal group

and phase velocities. Indeed, the dispersion relation² [7]:

$$\omega_{ia}^2 = k^2 \frac{k_B(T_e + \gamma T_i)}{m_i}$$

shows that ion wave frequency is proportional to the wave vector. The ion-acoustic sound speed is then obtained from this relationship as:

$$c_{ia} = \sqrt{\frac{k_B(T_e + \gamma T_i)}{m_i}} \quad (3)$$

4. Physics of Collisionless Shocks

A shock wave is a macroscopic discontinuity separating two different regimes in an otherwise continuous medium. Space plasmas host both traveling shocks, driven by coronal mass ejections propagating through interplanetary space, and standing bow shocks, formed when supersonic solar wind decelerates against a planetary magnetosphere.

In both classical gas-dynamics and MHD, a shock is broadly defined by three main characteristics:

- The relative flow velocity must exceed the local signal speed. In space plasmas, information propagates via both acoustic and magnetic waves, governed by the magnetosonic speed (2).
- At the shock front, the properties of the medium change abruptly over a very thin layer. While hydrodynamic shocks exhibit jumps in pressure, temperature, and density, MHD shocks are additionally characterized by a sharp increase in magnetic field strength.
- Upstream kinetic energy is irreversibly converted into thermal energy downstream, increasing the entropy of the system.

In traditional gas-dynamic shocks, this irreversible dissipation is driven by binary molecular collisions, manifesting as viscosity. These collisions establish a local Maxwellian velocity distribution, equalizing temperatures across different species and allowing a single thermodynamic temperature to be defined.

Conversely, space plasmas are extremely rarefied

²The subscript e refers to electrons, while the i to ions.

gases: close to Earth's orbit, the mean free path of a solar wind proton is on the order of 1 AU [6]. Compared to the length scale of the magnetosphere, this yields $K_n \gg 1$, that places the flow in the free-molecular region. Under such conditions, collisions are virtually negligible; hence, we talk about *collisionless shocks*. The lack of collisions has some relevant implications: electrons and protons can maintain different temperatures, and their velocity distributions can significantly differ from a Maxwellian distribution, making the concept of thermodynamic temperature inadequate. Furthermore, the presence of the magnetic field might even lead to highly anisotropic particle distributions, and processes of dissipation involve complex interactions between particles and fields. Nonetheless, shocks frequently occur in space plasmas. While collisional coupling between the particles is negligible, the electromagnetic field acts as a coupling medium. Observations show that the collective behavior of the plasma is mainly due to wave-particle interactions. Thus, collisionless shocks represent a profound example of a macroscopic phenomenon regulated entirely by microscopic, non-collisional kinetic processes [11].

4.1. Conservation Laws

Plasma properties such as flow velocity $\mathbf{u}$, magnetic field $\mathbf{B}$, mass density ρ , and pressure p differ significantly between the upstream and downstream regions. The relationship between these two sets of parameters is established by conservation laws known as the Rankine-Hugoniot equations.

4.1.1 Reference Frame

The simplest description of a gas-dynamic shock uses the shock's stationary reference frame (see Figure 1). In this frame, gas with a speed larger than the signal speed flows into the shock from the upstream medium, which has not yet received any information about the approaching obstacle. At the shock front, irreversible processes lead to the compression of the gas and a sudden drop in velocity. Across this discontinuity, mass conservation must be satisfied; thus, the mass flux through each element of the shock surface is constant.

A traveling shock and a standing shock are




Figure 1: Reference frames for MHD shocks: normal incidence frame (*left*) and de Hoffmann-Teller frame (*right*). Adapted from [11].

equivalent, differing only by a Galilean change of the reference frame.

The crucial difference between an MHD shock and an ordinary shock is the presence of the magnetic field. Hence, the conservation laws must be expanded to also accommodate this field, and the geometry becomes more complex because the flow is not necessarily parallel to it. To simplify the analysis, a special reference frame is often used, known as the *de Hoffmann-Teller frame*, shown in Figure 1. In the normal incidence frame, the upstream plasma flow is perpendicular to the shock but oblique to the magnetic field, while the downstream flow is oblique to both the magnetic field and the shock normal. In the de Hoffmann-Teller frame [8], the plasma flow is parallel to the magnetic field on both sides of the shock, and the $\mathbf{u} \times \mathbf{B}$ induction field in the shock front vanishes: the reference frame moves parallel to the shock front with the de Hoffmann-Teller velocity $\mathbf{v}_{\text{HT}} \times \mathbf{B} = -\mathbf{E}$.

4.1.2 Rankine-Hugoniot Relations

For an MHD shock, the Rankine-Hugoniot relations can be inferred in the same way as in gas-dynamic shocks³. In this section, the abbreviation $[X] = X_u - X_d$ denotes the difference of a quantity X between the upstream and downstream media.

Let $\mathbf{n}$ be the unit vector along the shock normal; the Rankine-Hugoniot equations are [11]:

- Mass balance, which is identical to the ordinary shock,

$$[\rho \mathbf{u} \cdot \mathbf{n}] = 0$$

- Momentum balance, where the additional terms describe the magnetic pressure per-

³A detailed derivation of the classical relations can be found in [1].

pendicular to the shock front,

$$\left[\rho \mathbf{u}(\mathbf{u} \cdot \mathbf{n}) + \left(p + \frac{B^2}{2\mu_0} \right) \mathbf{n} - \frac{(\mathbf{B} \cdot \mathbf{n})\mathbf{B}}{\mu_0} \right] = \mathbf{0}$$

- Energy balance, where the additional terms account for the electromagnetic energy flux $\mathbf{E} \times \mathbf{B}/\mu_0$ with the electric field expressed by $\mathbf{E} = -\mathbf{v} \times \mathbf{B}$,

$$\left[\mathbf{u} \cdot \mathbf{n} \left(\frac{\rho u^2}{2} + \frac{\gamma}{\gamma - 1} p + \frac{B^2}{\mu_0} \right) - \frac{(\mathbf{B} \cdot \mathbf{n})(\mathbf{B} \cdot \mathbf{u})}{\mu_0} \right] = 0$$

- Maxwell's equations,

$$[\mathbf{B} \cdot \mathbf{n}] = 0 ,$$

which follows from $\nabla \cdot \mathbf{B} = 0$, stating that the normal component of the magnetic field is continuous ($B_n = \text{const}$), and

$$[\mathbf{n} \times (\mathbf{u} \times \mathbf{B})] = \mathbf{0} ,$$

stating that the tangential component of the electric field must be continuous.

The Rankine-Hugoniot equations are, therefore, a set of five conservation relations, sufficient to determine the downstream plasma properties (ρ , $\mathbf{u}$, p , and $\mathbf{B}$) from the upstream conditions across an MHD shock.

4.1.3 Shock Geometry

A critical parameter in the description of an MHD shock is the local geometry; specifically, the angle θ_{Bn} between the magnetic field direction and the shock normal. Shocks are geometrically classified as follows:

- Perpendicular shock, propagating perpendicularly to the magnetic field: $\theta_{Bn} = 90^\circ$;
- Parallel shock, which propagates parallel to the magnetic field: $\theta_{Bn} = 0^\circ$;
- Oblique shock, that propagates at any θ_{Bn} between 0° and 90° . Oblique shocks can be subdivided into: quasi-parallel shocks with $0^\circ < \theta_{Bn} < 45^\circ$ and quasi-perpendicular shocks with $45^\circ < \theta_{Bn} < 90^\circ$.

For example, the configuration shown in Figure 1 represents a quasi-parallel shock.

Notice that in a parallel shock, the tangential

magnetic field is zero ($B_t = 0$), and the magnetic field passes through unchanged. Conversely, in perpendicular shocks, the normal component of the magnetic field vanishes ($B_n = 0$), and both the plasma pressure and the magnetic field strength increase across the shock.

4.1.4 Fast and Slow Shocks

In a plasma, different modes of MHD waves can steepen into shocks: fast, slow, and intermediate waves. Among these, only the fast and slow waves are compressive, while the intermediate shock, often called Alfvén shock, exists only in an anisotropic medium [11].

Physical compressive shocks are thereby formed exclusively by fast and slow magnetosonic waves. In both modes, the plasma density and pressure increase across the shock. The phase speeds of these modes are [16]:

$$2v_{\text{fast,slow}}^2 = v_{ms}^2 \pm \sqrt{v_{ms}^4 - 4v_A^2 v_s^2 \cos^2 \theta_{Bn}}$$

with the (+) sign referring to the fast, and the (−) sign to the slow mode.

The change in the magnetic field is different in fast and slow shocks: in a fast shock, the magnetic field increases and is bent away from the shock normal because the normal component of the field is constant. The normal component of the upstream (downstream) flow speed is larger (smaller) than the propagation speed of a fast MHD wave, and both upstream and downstream flow speeds exceed the Alfvén speed. In a slow shock, the upstream speed exceeds the sound speed, but not the Alfvén speed. In addition, the magnetic field strength decreases across the shock, and the field is hence bent towards the shock normal (see Figure 2 and Figure 3). Traveling interplanetary shocks, in general, and planetary bow shocks, in particular, are always fast MHD shocks.

In the shock frame, the Alfvén Mach number is defined relative to the shock velocity v_{sh} and the upstream fluid velocity u_u :

$$M_A = \frac{v_{sh} - u_u}{v_A}$$

Because of geometric dependencies, even fast MHD shocks can occasionally yield $M_A < 1$.




Figure 2: Changes in magnetic field and plasma parameters across a fast (left) and a slow (right) shock. Reproduced from [4].




Figure 3: Change in magnetic field direction across a fast and slow MHD shock. Adapted from [11].

5. The Bow Shock

The Earth's bow shock is a standing wave that forms as the hypersonic solar wind is decelerated by the planet magnetosphere, which acts as a blunt obstacle at rest. As illustrated in Figure 4, the flow decreases from super-magnetosonic to sub-magnetosonic across the parabolic shock, separating the undisturbed upstream region from the turbulent magnetosheath downstream.




Figure 4: The Earth's bow shock.

5.1. Mach Number

The thermal gas pressure in a fully ionized gas with equal ion and electron number densities ($n_i = n_e = n$) is given by:

$$p_{gas} = nk_B(T_i + T_e)$$

Using the characteristic solar wind parameters reported in Table 1, the computed pressure is approximately 25 pPa . Acoustic waves in an ionized gas with thermal pressure p_{gas} and mass density $\rho \approx nm_i$ propagate at the ion-acoustic sound speed (3). Assuming an ideal monoatomic plasma ($\gamma = 5/3$) and the parameters from Table 1, the ion-acoustic speed is on the order of 60 km/s . The ratio of the average solar wind velocity to this sound speed provides a free-stream Mach number $M_\infty \approx 8$ [16], confirming that the flow is hypersonic.

Furthermore, the magnetic field embedded within the solar wind exerts a magnetic pressure $p_{mag} = B^2/2\mu_0$, which amounts to about 20 pPa at 1 AU. This value is directly comparable to the thermal gas pressure, demonstrating that magnetic effects play a role as crucial as the thermal pressure forces in controlling the solar wind dynamics. Consequently, Alfvén waves (see subsection 3.1) propagate at speeds comparable to the sound speed. The magnetosonic Mach number, defined as the ratio of the solar wind speed to the magnetosonic speed, can also be estimated: since the bow shock nose is quasi-perpendicular, the relevant signal speed is the perpendicular fast speed of (2); combining (1) and (3) then yields $v_{ms} \approx 85 \text{ km/s}$, giving $M_{ms} \approx 5 - 6$ and confirming that magnetic and thermodynamic effects are comparable.

5.2. Shock Properties and Geometry

Because the solar wind at 1 AU is a high-Mach number stream, the bow shock manifests as a fast-magnetosonic shock. The collisionless nature of the shock can be confirmed by estimating the Knudsen number, $K_n = \lambda/L$. Near 1 AU, the mean free path of a solar wind proton is on the order of 1 AU itself ($\lambda \approx 1.5 \times 10^8 \text{ km}$), whereas the shock thickness is on the order of the ion inertial length ($L \approx 100 \text{ km}$) [2, 6]. This yields $K_n \approx 10^6$, placing the flow firmly in the free-molecular regime, where binary collisions are negligible. Despite the absence of collisions, the shock still compresses the plasma

strongly: both the plasma density and magnetic field strength increase abruptly across the discontinuity, jumping by a factor of roughly 4 [13, 15, 16] - the classical theoretical upper limit for strong shocks in a $\gamma = 5/3$ gas.

In situ measurements reveal a variety of magnetic field fluctuations and complex structural transitions caused by the interaction of the solar wind with the geomagnetic field. Spacecraft measurements place the nominal subsolar stagnation point of this collisionless MHD shock wave at approximately 13.4 Earth radii from the center of our planet [13].

The shock exists only over a spatially limited region in front of the Earth. This is because the effective Mach number governing shock formation depends strictly on the solar wind velocity component normal to the local shock surface ($v_n = v \cos \theta$). The super-magnetosonic condition ($M_{ms} > 1$) is satisfied only as long as the angle θ between the solar wind velocity vector and the shock normal obeys the geometric constraint $\theta < \arccos(1/M_{ms})$ [3, 11]. For $M_{ms} \approx 5$, the maximum angle up to which the shock can be maintained is $\theta_{max} \approx 78^\circ$. Consequently, the bow shock forms a spatially restricted curved shield in front of the magnetosphere, undergoing a continuous transition from a high-Mach number, quasi-perpendicular shock at the nose, to a weak, low-Mach number shock along its distant flanks.

5.3. Currents System

Since binary collisions cannot provide the dissipation, the compression is instead driven by the electric currents and self-generated fields within the shock layer. Evidence of this structure appears in the magnetic field profile of Figure 5, which shows an initial slight increase, known as the *foot*, followed by a *ramp* across the discontinuity, and at the end, an *overshoot* [2, 3, 17].

The jump in the tangential component of the magnetic field across the shock indicates that the bow shock itself is a current layer. It possesses an internal surface current density, j_{sh} , which accounts for the change in the magnetic field. This can be estimated as:

$$j_{sh} = \frac{\|B_t\|}{\mu_0 d_{sh}}$$

where d_{sh} represents the shock width, as depicted in Figure 6. This current increases the




Figure 5: Shock profiles of electron density, proton velocity and magnetic induction determined from spacecraft measurements.

magnetic field strength behind the shock.

Instead, the slight increase in the magnetic field strength in the foot region is primarily due to the appearance of reflected ions in front of the shock. Solar wind ions and electrons encountering the compressed perpendicular magnetic field of the shock will have different gyro-radii [15]; notably, ions can penetrate deeper into the field than the electrons. This difference in penetration depth generates a charge separation electric field, $\mathbf{E}_{sh}$, in the shock normal direction, pointing toward the Sun. This field reflects a certain number of ions back into the solar wind while attracting and capturing electrons. The fraction of reflected ions returns to the solar wind in front of the shock and performs another gyration in its magnetic field; hence, they carry a current in the foot region ($\mathbf{j}_f$ in Figure 6), which closes the shock current in front of the discontinuity.

Finally, the magnetic overshoot in the shock profile is related to another current layer that develops inside the shock transition region. This current is a pure electron drift current [11], indicated as $\mathbf{j}_e$ in Figure 6, resulting from the presence of the charge separation field $\mathbf{E}_{sh}$. This field is restricted to a thin layer, narrower than the ion gyro-radius in the compressed shock magnetic field, but wider than the electron gyro-radius. Consequently, within this layer, the electrons undergo an electric $\mathbf{E} \times \mathbf{B}$ drift motion [10] in the crossed electric and magnetic fields, while the ions remain largely unaffected. This drift generates an electron current ($\mathbf{j}_e$) that flows in the same direction as the main shock current $\mathbf{j}_{sh}$. This locally amplifies the current, thereby causing the magnetic overshoot appearing in Figure 5.




Figure 6: Current system in the bow shock thickness.

5.4. Heating Mechanism

At a shock front, irreversible dissipation occurs. This can be described as the transformation of ordered energy into random thermal energy among the particles. In an ordinary fluid, this dissipation is provided by viscosity, which arises from collisions between molecules that redistribute energy among all particles. However, a viscosity mechanism cannot occur in a rarefied, cold plasma because particle collisions are negligible. Instead, wave-particle interactions are responsible for the exchange of momentum [9]. As explained in subsection 5.3, a fraction of the ions approaching the shock front is reflected back into the upstream region. The interaction between these reflected ions and the incoming flow creates turbulence, resulting in the generation of waves [18]; in particular, ion acoustic and Alfvén waves. Among the randomly moving particles in the plasma, some travel at a velocity that matches the phase velocity of the plasma wave. These particles are in resonance with the wave, allowing an intense exchange of energy to take place. This mechanism, known as *Landau damping* [9], transfers energy from the wave to the resonant particles. Due to the Lorentz force acting on an ion, its trajectory is curved, causing it to interact repeatedly with the same wave, as shown in Figure 7. Each encounter adds more energy to the particle. Furthermore, electrons can interact with the ion acoustic and Alfvén waves, which transfer their momentum and change their motion. This process retards the electron flow in the plasma, creating a resistance to the electric current. This represents an-




Figure 7: Wave-particle interaction.

other dissipation mechanism that operates without involving direct particle collisions.

Ultimately, through these collisionless processes, particles exchange energy not through direct impacts, but via the waves generated at the shock by electromagnetic field fluctuations. This shock heating relies on mechanisms unique to plasmas; yet it effectively broadens the velocity distribution function, thermalizing the plasma.

6. Conclusions

In conclusion, the Earth's bow shock represents a clear example of a collisionless MHD shock, differing from conventional gas-dynamic shocks due to the extreme rarefaction of space plasmas. The interaction between the hypersonic solar wind and the Earth's magnetosphere produces a standing, fast-magnetosonic shock wave. Despite the negligible role of binary molecular collisions, the flow decelerates, and plasma properties such as density and magnetic field intensity increase abruptly across the discontinuity, governed by the modified Rankine-Hugoniot relations. Crucially, the irreversible dissipation of kinetic energy into thermal energy is achieved not through viscosity, but via complex, microscopic wave-particle interactions and local electric current systems within the shock layer. Ultimately, understanding these non-collisional processes is crucial for comprehending the macroscopic behavior of the bow shock.

Nonetheless, many aspects remain open — notably the structure of the quasi-parallel shock and its turbulent foreshock, and the question of whether the shock front is truly stationary or continuously reforms — making the bow shock a natural laboratory for collisionless shock physics that continues to be probed by dedicated spacecraft missions.

References

- [1] John D. Anderson, Jr. *Modern Compressible Flow: With Historical Perspective*. McGraw-Hill, 3rd edition, 2003.
- [2] Stuart Bale, MA Balikhin, T. Horbury, V. Krasnoselskikh, H. Kucharek, E. Möbius, Simon Walker, Andre Balogh, David Burgess, Bertrand Lembège, Elizabeth Lucek, M. Scholer, and Michelle Thomsen. Quasi-perpendicular shock structure and processes. *Space Science Reviews*, 118:161–203, 06 2005.
- [3] Wolfgang Baumjohann and Rudolf Treumann. *Basic Space Plasma Physics*, volume 15-22. 09 1996.
- [4] Wolfgang Baumjohann and Rudolf Treumann. *Basic Space Plasma Physics - Revised Edition*. 03 2012.
- [5] P. Bochsler. Solar wind composition. *Encyclopedia of Astronomy and Astrophysics*, -1:2303, 10 2000.
- [6] Antoine Bret and Asaf Pe'er. Bridging the gap between collisional and collisionless shock waves. *Journal of Plasma Physics*, 87(2):905870204, 2021.
- [7] Francis Chen. *Introduction to Plasma Physics and Controlled Fusion*. 01 2016.
- [8] F. De Hoffmann and E. Teller. Magneto-hydrodynamic shocks. *Phys. Rev.*, 80:692–703, Nov 1950.
- [9] R. J. Goldston and P. H. Rutherford. Introduction to plasma physics. *Journal of Plasma Physics*, 59(2):387–389, 1998.
- [10] Umran S Inan and Marek Gołkowski. *Principles of Plasma Physics for Engineers and Scientists*. Cambridge University Press, 2010.
- [11] Russell Kivelson. *Introduction to Space Physics*. Cambridge University Press, 1995.
- [12] P. Manoharan. *The Solar Wind*, pages 299–326. 10 2008.
- [13] Norman F. Ness, Clell S. Searce, and Joseph B. Seek. Initial results of the imp 1 magnetic field experiment. *Journal of Geophysical Research (1896-1977)*, 69(17):3531–3569, 1964.
- [14] V. Obridko and Oleg Vaisberg. On the history of the solar wind discovery. *Solar System Research*, 51:165–169, 03 2017.
- [15] G. Parks, Eungmok Lee, Suiyan Fu, Naiguo Lin, Y. Liu, and Zhongwei Yang. Shocks in collisionless plasmas. *Reviews of Modern Plasma Physics*, 1, 06 2017.
- [16] John R. Spreiter, Audrey L. Summers, and Alberta Y. Alksne. Hydromagnetic flow around the magnetosphere. *Planetary and Space Science*, 14(3):223–253, 1966.
- [17] Rudolf Treumann and Wolfgang Baumjohann. *Advanced Space Plasma Physics*. 01 1997.
- [18] D. Winske and N. Omidi. A nonspecialist's guide to kinetic simulations of space plasmas. *Journal of Geophysical Research: Space Physics*, 101(A8):17287–17303, 1996.