

The Sir Ivan Fire: observed pyroCb formation and features, ground-plume coupling and coldfront impact - Article Frame

Alex Terrasson

May 30, 2017

1 Abstract

2 Introduction

Points to be covered:

- Presentation of the fire (date, quick overview of the meteorological context, area burnt, pyroCb presence)
- Context and objectives
 - Extreme fire with pyroCb generation with a lot of observations: good case study
 - Focus of the study: We apply known methods (use of doppler radar to get axial/absolute doppler velocity and turbulence) to a different kind of plume: a pyroCb
 - focusing on plume study, the goal is to generalize to some extent what kind of turbulence/vorticity can be expected in a pyroCb.
- State of the art:
 - Radar Doppler are widely used in storm studies to detect large vortices/turbulence (couple of references to add)
 - PyroCb have been studied in Australia and America through several case studies (Chislom fire, 2013 Rim fire), mostly via satellite.
 - Vortices often develop within the plume at different scales (Development of vortices on prescribed fires, review of vortices in wildland fire), but these vortices have not been watched many times (Smoke Column observation from two forests fires using doppler lidar and doppler radar)
 - The turbulence field within a pyroCb has not been studied

3 Methods used

- Namoi radar characteristics (WSR 74 S band doppler), highlight on the spec width
- NCAR Turbulence Detection Algorithm (to get the turbulence field from the reflectivity and the spec width)
- Single Doppler algorithm

4 Results

4.1 Meteorological and ground context

Meteorological context

Main points:

- Perfect ground/low-level condition for an extreme fire behaviour: surface winds, RH, Temp
- Good pyroCb condition: Mid-level moisture/upper-level instability
- a coldfront was present, link to even bigger event

Figure here to summarize the meteorological conditions: The center of the figure could be a couple of top views of the area at key time with the surface winds and the coldfront passage, the sides of the figure could be available soundings to see the stability of the atmosphere

Ground fire results

Main points:

- Show the dramatic increase in spreading/intensity on Sunday afternoon

1 here: The center will be the top view of the burnt area evolution, and a graph with the FRP (fire rate power) and the burnt area over time would complete it.

4.2 Plume Structure and developments

Main points:

- The plume has three steps of development
 1. Before pyroCb
 2. After pyroCb/ before ColdFront
 3. After Coldfront

- The plume has a structure resulting of both buoyancy driven effects and wind driven effects.
- Structure description:
 - Vertical column (buoyancy driven)
 - Horizontal spread at low/mid elevations in the south east (wind driven)
 - Horizontal spread at low elevation in the north direction after the coldfront (wind driven)

Figure 2 here: himawari and 2-3 reflectivity signatures and 3 key timesteps to show the pyroCb structure and its evolution

- PyroCb impact on the plume (top height and reflectivity distribution) **Figure here:** top height over time and reflectivity distribution at key time steps **Note: I still have to plot the reflectivity distribution in the whole volume**
- Litghnings (?)

4.3 Turbulence/ Vortices within the plume

Main points:

- Before the pyroCb: expected features:
 - The divergence line on the doppler plots follow the center line: no large vortices
 - The turbulences are maximum at the edges of the plume
- After the pyroCb we see unexpected features:
 - Large vortices signatures along the centerline of the plume
 - Turbulence field disturbed, maxima o turbulence along the centerline of the plume
 - The center of the plume presents both large scale (cf doppler signature) and smaller scale (cf turbulence field) vortices.

Figure here: Showing over 1 time step of the "normal" velocity field/turbulence and a series of the unexpected turbulence and vortices (3-4 between 17:20 and 18:10 to show the motion of the vortices)

5 Discussions

5.1 Discussion about the pyroCb generation

The atmospheric/ground conditions are they consistent with the litterature ? **Note: The point would be to show that our case is not that different from others, to then support a generalization**

5.2 Analysis of the vortices/ turbulence within the plume

What is the part of the pyroCb ? what is the part of the coldfront ? can some other parameters explain the unexpected features ? **Note: this part is not clear to me yet, it is definitely where there still is the more to do**

6 Conclusion

Note: Will depend on the discussion

7 References

8 figures

Note: These figures are not finished yet, figure 1 needs some more data from the NFS, figure 2 needs some presentation work from me, figure 3 might need additional elevation, and maybe the single doppler field. Since this last figure is crucial for the article, I was considering splitting it in two one for the doppler velocity, the other part for the turbulence.

Note: Two more figures are to be done as well: The one presenting the atmospherical condition (I m waitng for data front the NFS, and the one presenting the top height and the reflectivity distribution)

Figure 1: Fire spread during the event (main), area burnt and FRP over time (bottom right).
 Note: the point of this figure is to show the massive increase in area and FRP on sunday. The graph does not display the right information yet (I m waiting for the ground data from the NFS)




Figure 2: Evolution of the plume on sunday. The each column corresponds to a time step: the first is 15:00pm, before the pyroCb, the second is 17:00, after the apparition of the pyroCb and during the coldfront passage, the last columns corresponds to 18:10pm, after the coldfront passage. The first line shows the visible Himawari view, and the two other rows the reflectivity signature for 0.5 and 1.8 degree of elevation. **Note: the point of this figure is to show the plume evolution: first a soft expansion to the south-east, then the apparition of the pyroCb, and finally the coldfront creating a new expansion to the north east, but only at low level**




Figure 3: Axial Doppler velocity and turbulence field at four time steps: 15:40, before the apparition of the pyroCb ; 17:00, during the pyroCb event, and at the begining of the large vortices signatures and the move of the maximum of turbulence toward the centre ; 17:50 and 18:10 with the motion toward south east of the large vortex. **note: the point of this figure is to show The apparition and the development of the vortexes and the turbulence displacement.** An interesting featur is that the vortex doesn't follow the coldfront wind direction, but the mid level wind direction. So I m considering adding another radar elevation to show that the same signature on vortex and turbulence are present as well

