

Assessing the Impact of the Anacostia Watershed Restoration Plan on the Water Quality of the Northeast Branch

https://github.com/atf35/Fischer_WDA2022_FinalProject

Atalie Fischer

1 Rationale and Research Questions

The Anacostia River watershed is a heavily urbanized watersheds located in the Baltimore-DC area. It contains 14 major subwatersheds and a tidal portion covering approximately 176 square miles (USACE, 2022). The Northeast Branch and the Northwest Branch are the two main tributaries of the Anacostia River, which flows into the Potomac River and into the Chesapeake Bay. This watershed is one of the main priorities for restoration in the Chesapeake Bay Program. The Army Corps of Engineers (Corps) developed the Anacostia Restoration Plan in 2010 to improve and restore the Anacostia Watershed. It identified more than 3,000 projects (USACE, 2022). Some of the strategies of these projects include stormwater controls, stream restoration, wetland creation and restoration, fish blockage removal, reforestation, trash and toxic contaminant control, and parkland acquisition (Metropolitan Washington Council of Governments, 2010). These projects aim to improve water quality and reduce flooding.

The goal of this report is to assess the progress of the Anacostia Restoration Plan on water quality of the Northeast Branch. Due to limitations on data availability, this report will focus only on the Plan’s effectiveness in using green stormwater controls in removing suspended sediment, removing contaminants, and reducing the temperature from urban stormwater runoff (See the “Dataset Information” section for details on these specific parameters).

The main research question is: *Has there been an improvement in water quality of the Northeast Branch since the implementation of the Anacostia Restoration Plan in 2010?* This study will be guided by the following question comparing time periods before and after 2010:

How has water quality changed in terms of turbidity, temperature, specific conductance, and dissolved oxygen?

2 Dataset Information

Data were retrieved from the the United States Geological Survey (USGS) National Water Information System (NWIS). The dataRetrieval package in R was used to pull data directly from NWIS without the need to download any data files. Data were pulled from the gage on the Northeast Branch of the Anacostia River near Riverdale, Maryland (USGS gage #01649500). The water quality dataset was wrangled into two separate time periods: before the Anacostia Watershed Restoration Plan implementation in 2010 and after. The water quality data were sampled monthly and their sampling dates were rounded to the first of the month for even time steps. Linear interpolations were conducted to fill in any missing values.

Table 1: Summary of raw data used.

| Dataset | Information |
|---------------------|--|
| NEBranch_flow | Discharge data collected at USGS gage #01649500 from 1/1/2003 through 4/22/2022 |
| NEBranchWQ_raw_full | Water quality data collected at USGS gage #01649500 from 1/1/2003 through 4/22/2022. Parameters collected were turbidity, temperature, specific conductance, and dissolved oxygen. |

3 Exploratory Analysis

Turbidity was plotted over time to visualize any trends in suspended sediment (Figure 1). There are no obvious trends in the data from this visualization. However, there are six missing values. A summary of the data show that the average turbidity in the Northeast Branch is 107.920 FNU.




Figure 1: Turbidity of the North East Branch over time.

Temperature was plotted to visualize any trends over time (Figure 2). There are no obvious trends in temperature over time and a summary of the data show that there are no missing values in the dataset. However, around 2006 and 2007, a high temperature is not observed like the rest of the time period. The mean temperature over the full time period is 14.45 degrees C.




Figure 2: Temperature of the North East Branch over time.

Specific conductance was plotted over time to visualize any major trends in ionic concentration over time (Figure 3). The plot shows eight major peaks in specific conductance from 2003 through 2022, with the highest peak in 2021. A summary of the data show that the mean specific conductance is 390.8 $\mu\text{S}/\text{cm}$ and the maximum is 4640.0 $\mu\text{S}/\text{cm}$, which is an order of magnitude larger. There are no missing values for specific conductance in this dataset.




Figure 3: Specific conductivity of the North East Branch over time.

Dissolved oxygen was plotted to visualize any major trends over time (Figure 4). It appears that dissolved oxygen concentrations may be less variable in recent years, but it is difficult to tell in the plot. A summary of the data show that the average dissolved oxygen concentration is 10.400 mg/L and that there is one missing value for dissolved oxygen in this dataset.




Figure 4: Dissolved oxygen concentration of the North East Branch over time.

4 Analysis

4.1 Question: How has water quality changed?

4.1.1 Turbidity

Time series analyses were conducted on turbidity for the time period before the implementation of the Anacostia Watershed Restoration Plan (2003 to 2009) (Figure 5) and after its implementation (2010-2022) (Figure 6). A seasonal Mann-Kendall's test was conducted on both time periods to test for stationarity. Both time periods display stationarity and are neither significantly increasing or decreasing over time with p-values greater than 0.05 (Table 2). While these data are not statistically significant, it is worthwhile to consider the sign of the tau values. Turbidity prior to the Plan's implementation was generally increasing, while it is generally decreasing post-implementation. This suggests that over a longer time period, stormwater control projects could potentially reduce turbidity significantly.




Figure 5: Time series decomposition of turbidity from 2003 to 2009.

NULL

NULL




Figure 6: Time series decomposition of turbidity from 2010 to 2022.

4.1.2 Temperature

Time series analyses were conducted on temperature for the time period before the implementation of the Anacostia Watershed Restoration Plan (2003 to 2009) (Figure 7) and after its implementation (2010-2022) (Figure 8). A seasonal Mann-Kendall's test was conducted on both time periods to test for stationarity. Both time periods display stationarity and are neither significantly increasing or decreasing over time with p-values greater than 0.05 (Table 2). The Anacostia Watershed Restoration Plan did not significantly affect temperature in the Northeast Branch.




Figure 7: Plot of turbidity over time with linear models from 2003 to 2009 and 2010 to 2022.




Figure 8: Time series decomposition of temperature from 2003 to 2009.




Figure 9: Time series decomposition of temperature from 2010 to 2022.

4.1.3 Specific Conductivity

Time series analyses were conducted on specific conductivity for the time period before the implementation of the Anacostia Watershed Restoration Plan (2003 to 2009) (Figure 9) and after its implementation (2010-2022) (Figure 10). A seasonal Mann-Kendall's test was conducted on both time periods to test for stationarity. Both time periods display stationarity and are neither significantly increasing or decreasing over time with p-values greater than 0.05 (Table 2). While these data are not statistically significant, it is worthwhile to consider the sign of the tau values. Specific conductivity prior to the Plan's implementation was generally increasing, while it is generally decreasing post-implementation. This suggests that over a longer time period, stormwater control projects could potentially reduce specific conductivity and the concentration of pollutants significantly.




Figure 10: Plot of temperature over time with linear models from 2003 to 2009 and 2010 to 2022.




Figure 11: Time series decomposition of specific conductivity from 2003 to 2009.




Figure 12: Time series decomposition of specific conductivity from 2010 to 2022.

4.1.4 Dissolved Oxygen

Time series analyses were conducted on dissolved oxygen for the time period before the implementation of the Anacostia Watershed Restoration Plan (2003 to 2009) (Figure 11) and after its implementation (2010-2022) (Figure 12). A seasonal Mann-Kendall's test was conducted on both time periods to test for stationarity. Both time periods display stationarity and are neither significantly increasing or decreasing over time with p-values greater than 0.05 (Table 2). The Anacostia Watershed Restoration Plan did not significantly affect dissolved oxygen in the Northeast Branch.




Figure 13: Plot of specific conductivity over time with linear models from 2003 to 2009 and 2010 to 2022.




Figure 14: Time series decomposition of dissolved oxygen from 2003 to 2009.




Figure 15: Time series decomposition of dissolved oxygen from 2010 to 2022.

Table 2: Summary of results from stationarity tests.

| Parameter | Time Period | tau | p-value |
|-----------------------|-------------|---------|----------|
| Turbidity | Before | 0.13 | 0.17993 |
| . | After | -0.0992 | 0.12894 |
| Temperature | Before | 0.144 | 0.13719 |
| . | After | 0.0245 | 0.70806 |
| Specific Conductivity | Before | 0.0926 | 0.33815 |
| . | After | -0.0258 | 0.69284 |
| Dissolved Oxygen | Before | -0.181 | 0.061497 |
| . | After | -0.0626 | 0.33838 |




Figure 16: Plot of dissolved oxygen over time with linear models from 2003 to 2009 and 2010 to 2022.

5 Summary and Conclusions

Time series analyses and linear models conducted on turbidity, temperature, specific conductivity, and dissolved oxygen on the time periods both before and after the implementation of the Anacostia Watershed Restoration Plan in 2010 were not significant. Therefore, no concrete conclusions can be drawn from this analysis. However, we can speculate that turbidity and specific conductivity may decrease in the future given longer datasets.

6 References

Metropolitan Washington Council of Governments. (2010, April 19). Officials Release Landmark Anacostia Watershed Plan. Retrieved from Newsroom: <https://www.mwcog.org/about-us/newsroom/2010/04/19/officials-release-landmark-anacostia-watershed-plan-anacostia-restoration-water-quality/>

US Army Corps of Engineers (USACE). (2022, April 26). Anacostia Watershed Restoration. Retrieved from Baltimore District Website: <https://www.nab.usace.army.mil/Missions/Environmental/Anacostia-Watershed-Restoration/>