

Chapter 6: ODEs

Due date:

A correct answer without proper explanation will not receive full credit.

6.1 IVPs

- 3 Use separation of variables to find solutions of the IVP given by $y(0) = 1$ and the following differential equations:

(a) $y' = t$

(b) $y' = 2(t + 1)y$

(c) $1/y^2$

For which of the IVPs in the Exercise 3 does the advanced theorem of Existence and Uniqueness of Solutions guarantee a unique solution?

(a) Find the Lipschitz constants if they exist.

(b) What is the largest interval $[0, b]$ for which the solutions exist?

- 7 (a) Show that $y = \tan(t + c)$ is a solution of the differential equation $y' = 1 + y^2$ for each c .
(b) For each real number y_0 , find c in the interval $(-\pi/2, \pi/2)$ such that the initial value problem $y' = 1 + y^2$, $y(0) = y_0$ has a solution $y = \tan(t + c)$.

Computer Problems. Use the code snippet provided on Canvas

1. Apply Euler's Method with step size $h = 0.1$ on $[0, 1]$ to the initial value problems in Exercise 3. Print a table of the t values, Euler approximations, and error (difference from exact solution) at each step.
2. Plot the Euler's Method approximate solutions for the IVPs in Exercise 3 on $[0, 1]$ for step sizes $h = 0.1, 0.05$, and 0.025 , along with the exact solution.
- 4 For the IVPs in Exercise 3, make a log-log plot of the error of Euler's Method at $t = 1$ as a function of $h = 0.1 \times 2^{-k}$ for $0 \leq k \leq 5$.
- 7 Plot the Euler's Method approximate solution on $[0, 1]$ for the differential equation $y' = 1 + y^2$ and initial condition $y_0 = 1$, along with the exact solution (see Exercise 7). Use step sizes $h = 0.1$ and 0.05 .