

Exam Practice 7

Section: 014

Name: _____

Print off this assignment and write in your answers. **Turn this in Friday, April 17th in class (1:10pm)!** Each answer is one point unless noted differently.


- 1) A random sample of 157 bills from a local Bozeman restaurant were collected and researchers recorded the bill amount and the tip amount for each to create the given scatterplot. The regression line was calculated to be

$$\widehat{Tip} = -0.292 + 0.182(Bill)$$

- a) What is the correlation of the plot?

A. -0.89
B. 0.92
C. -0.03
D. 0.36

- b) Predict the tip amount on a bill costing \$23.70.



- c) One of the 157 bills was for \$23.70. The tip left was \$10. Calculate the residual.

- d) Which of the circled points would have the largest negative residual?

A. B. C. D.

- e) Interpret the slope of the line in the context of the problem.

- f) The y-intercept in this regression is not useful. What term best describes this problem?

A. Extrapolation
B. Regression
C. P-value
D. Residual

- 2) In June 2013, the mayor of New York City proposed a limit of at most 16 ounces on sugary beverages sold in restaurants. He argued that limiting sugar intake would lower rates of obesity and diabetes. A Gallup poll used a random sample of 1015 US adults on June 15-16. They asked if people support a ban on sugary beverages with over 16 ounces in restaurants. Only 485 respondents said they favored the ban.
- a) We want to know if less than half of NYC residents favor the ban. What are our null and alternative hypotheses in proper notation? (2 pts)
 H_0 :

 H_a :
 - b) Is the sample size sufficiently large to use the Normal distribution? Show your work to check this assumption.
 - c) Compute the standard error of \hat{p} under the null hypothesis. You must show your work to receive credit.
 - d) Compute the test statistic. Again, show your work.
 - e) The p-value was found to be 0.080. Give your conclusion in the context of the problem.
- 3) An article in *Journal of Periodontics* studied 52 young people with tongue piercings and found that 18 of them had receding gums, which could result in later tooth loss. The participants were basically a convenience sample, but the intended population is all “young people” with tongue piercings.
- a) Is this a sufficient large sample to use the normal distribution? Explain your reasoning.
 - b) Compute the point estimate (\hat{p}).
 - c) We want to build a 99% confidence interval. What multiplier from your table will you use?
 - d) Build a 99% CI for the true proportion of “young people” with tongue piercings who have receding gums. Show all work!
 - e) To what group of people does this interval apply?