خدمتكم عبادة نتقرب بها الى الله

2021

ملخص

د. محمد فؤاد

إحصاء واحتمالات

No.

* Elements of probability :-

- Definitions:

- Sample space: The set of all expected outcomes resulting from the prob. experiment denoted by Ω

e.g: In the experiment of tossing a coin one time,
find Ω

$\Omega = \{H, T\}$ → H: Head, T: tail

eg. In the experiment of tossing a coin two times,
find Ω

$\Omega = \{HH, HT, TH, TT\}$

- Event: Any subset of Ω. denoted by any letter
(A, B, C, ----)

eg let $\Omega = \{1, 2, 3\}$, Find a subset of Ω

$\{1\}, \{2\}, \{3\}, \{1,2\}, \{1,3\}\{2,3\}, \{1,2,3\}, \phi$ 8 events

- Probability of event:

The probability of an event A is denoted by

$$P(A) = \frac{\#\text{ of elements in } A}{\#\text{ of elements in } \Omega}$$

Event → impossible $P(A) = 0$
→ sure $P(A) = 100\% = 1$ certain event
→ ordinary $0 < P(A) < 1$

Scanned with CamScanner

Scanned with CamScanner

No.

e.g: In the experiment of tossing a coin 2 times.
find:

a) the prob. of getting only one head.

b) the prob. of getting at least one head.

Sol:

$\Omega = \{HH, HT, TH, TT\}$

a) $A = \{HT, TH\}$ $P(A) = \frac{2}{4} = 0.5$

b) $B = \{HT, TH, HH\}$ $p(B) = 3/4 = 0.75$

e.g: In the exp of tossing a coin 3 times, Find:

a) the prob. of getting only 2 heads

b) the prob. of getting at least 2 heads.

c) the prob. of getting at most 1 head

d) the prob of getting no heads

Sol:

$\Omega = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$

a) $A = \{HHT, HTH, THH\}$ $P(A) = 3/8$

b) $B = \{HHT, HTH, THH, HHH\}$ $P(B) = 4/8$

c) $C = \{HTT, THT, TTH, TTT\}$ $P(C) = 4/8$

d) $D = \{TTT\}$ $p(D) = 1/8$

Scanned with CamScanner

Scanned with CamScanner

e.g: In the exp. of rolling one dice, find the prob.
a) of getting an odd no.
b) of getting a prime no.

$\Omega = \{1, 2, 3, 4, 5, 6\}$

Sol: a- $A = \{1, 3, 5\}$ $P(A) = 3/6$

b- $\{1, 2, 3, 5\}$ $P(B) = 3/6$

- rolling one dice twice
=
rolling two dices once.

e.g: In the exp. of rolling two dices, find the prob. of:
a) the sum of faces is 9
b) one of the faces is twice the other
c) the 2nd entry is triple the first
d) the 1st entry is prime
e) the sum of the faces is a prime less than 4

Sol:

$\Omega = \{(1,1), (1,2), (1,3), \ldots (1,6)$
$(2,1), (2,2), (2,3) \ldots (2,6)$
$\vdots$
$(6,1), (6,2) (6,3) \ldots (6,6)\}$

$|\Omega| = 36$

a) $A = \{(3,6), (4,5), (5,4), (6,3)\}$

$P(A) = 4/36 = 1/9$

* $|\Omega|$ in the exp. of tossing a coin n times is 2^n.

* $|\Omega|$ in the exp. of rolling a dice n times is 6^n.

b) $B = \{(1,2), (2,1), (2,4), (4,2), (3,6), (6,3)\}$

$P(B) = 6/36 = 1/6$

Scanned with CamScanner

Scanned with CamScanner

c) $C = \{(1,3), (2,6)\}$

$P(C) = 2/36 = 1/18$

d) $D = \{(1,1), (1,2) \cdots (4,6), (2,1), (2,2) \cdots (2,6), \cdots (5,6)\}$

$P(D) = 18/36 = 2/3$

e) $e = \{(1,1), (1,2), (2,1)\}$

$P(D) = 3/36 = 1/12$

* Prob. rules :-

1) $P(\phi) = 0$, $P(\Omega) = 1$

2) $0 \leq P(A) \leq 1$

3) Multiplicative law of prob:

- if A, B are two independant events, then

$P(A \cap B) = P(A) \cdot P(B)$

** $A \cap B$ = A intersects B ≡ A and B ≡ both of A, B

Def : The two events are indep. if there is no relation between them and they come from the same background.

Def : The two events are called disjoint if there is no relation between them but they come from diff. backgrou

- if A, B are disjoint, then $P(A \cap B) = 0$

** Events → indep.

→ disjoint

→ dependent → $P(A \cap B) \neq 0 \neq P(A) \cdot P(B)$

Scanned with CamScanner

Scanned with CamScanner

No.

4) Additive law of prob. :

→ $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

** $A \cup B \equiv A$ union $B \equiv A$ or $B \equiv$ at least one of them $\equiv$ A or B or both

** if A, B are indep.

$$P(A \cup B) = P(A) + P(B) - P(A) \cdot P(B)$$

** if A, B are disjoint.

$$P(A \cup B) = P(A) + P(B)$$

e.g: Two fire detecters A, B are working independently

s.t. $P(A) = 0.2$, $P(B) = 0.4$

a) find the prob. that both of them will work

b) find the prob. that at least one of them will work.

Sol:

a- $P(A \cap B) = P(A) \cdot P(B) = (0.2)(0.4) = 0.08$

b- $P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.2 + 0.4 - 0.08 = 0.52$



$P(A \cup B) = P(A) + P(B) - P(A \cap B)$



$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

Scanned with CamScanner

Scanned with CamScanner

No.

e.g: A fire fighting system consists of 3 indep. component with prob. of working of:

$P(A) = 0.2$, $P(B) = 0.4$, $P(C) = 0.6$

Find the prob. that the system will work

a- if the components are connected in series → indep

b- if the components are connected in parallel → dep

c- if only A is connected in series

d- if only B is connected in parallel

Sol: a)



$P(\text{system works}) = P(A \cap B \cap C)$

$= P(A) \cdot P(B) \cdot P(C) = (0.2)(0.4)(0.6) = 0.0$

b)



$P(\text{system works}) = P(A \cup B \cup C)$

$= P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

$= 0.2 + 0.4 + 0.6 - 0.08 - 0.24 - 0.12 + 0.048 = 1.0 - 792.8$

** $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

} → Demorgan law

Scanned with CamScanner

Scanned with CamScanner

No.

Wednesday
27

c) $P(A) = 0.2$, $P(B) = 0.4$, $P(C) = 0.6$



$P(\text{system work}) = P(A \cap (B \cup C))$

$= P(\underline{A \cap B}) \cup P(\underline{A \cap C}))$

$= P(A \cap B) + P(A \cap C) - P(\underline{(A \cap B) \cap (A \cap C)})$ → $A \cap B \cap C$

$- P(A) \cdot P(B) + P(A) \cdot P(C) - P(A) \cdot P(B) \cdot P(C)$

$= (0.2)(0.4) + (0.2)(0.6) - (0.2)(0.4)(0.6)$

d)



$P(\text{system works}) = P(B \cup (A \cap C))$

$= P(B) + P(A \cap C) - P(B \cap (A \cap C))$

$= P(B) + P(A) \cdot P(C) - P(A) \cdot P(B) \cdot P(C)$

$= 0.4 + (0.2)(0.6) - (0.2)(0.4)(0.6)$

Scanned with CamScanner

5) Complement of Prob. 3

$\bar{A}$ = the complement of A.

= $\Omega - A$ → All items in Ω except of A



1) $A \cup \bar{A} = \Omega$

2) $A \cap \bar{A} = \emptyset$ → disjoint

3) $P(A \cap \bar{A}) = 0$

4) $P(A \cup \bar{A}) = P(A) + P(\bar{A})$

$P(\Omega) = P(A) + P(\bar{A})$

$1 = P(A) + P(\bar{A})$

$P(\bar{A}) = 1 - P(A)$

Theorem: If A, B are indep., then

1) $A, \bar{B}$ are indep.

2) $\bar{A}, B$ " "

3) $\bar{A}, \bar{B}$ " "

Proof:

3) we want to show that $\bar{A}, \bar{B}$ are indep.

i.e. we want to show that

$P(\bar{A} \cap \bar{B}) = P(\bar{A}) \cdot P(\bar{B})$

L.H.S

$P(\bar{A} \cap \bar{B}) = P(\overline{A \cup B})$

$= 1 - P(A \cup B)$

$= 1 - [P(A) + P(B) - P(A \cap B)]$

$= 1 - P(A) - P(B) + P(A) \cdot P(B)$

$= 1 - P(B)(1 + P(A))$

الدكتور ما بده الاثبات

"عشان التعلم عن بعد"

smile

Scanned with CamScanner

Scanned with CamScanner

$= P(\bar{A}) - P(B)[1 - P(A)]$ (2)

$P(\bar{A}) - P(B)\,P(\bar{A})$

$= P(\bar{A})[1 - P(B)]$

$= P(\bar{A}) \cdot P(\bar{B})$ #

---

* Demorgans laws :

1) $\bar{A} \cap \bar{B} = \overline{A \cup B}$
2) $\bar{A} \cup \bar{B} = \overline{A \cap B}$
3) $\overline{A \cap B \cap C} = \bar{A} \cup \bar{B} \cup \bar{C}$
4) $\overline{A \cup B \cup C} = \bar{A} \cap \bar{B} \cap \bar{C}$

Ex] A, B are indep. Components such that

$P(A \text{ works}) = 0.3$, $P(B \text{ works}) = 0.6$

Find the prob. that :

a) B will fail.

b) at least one of them will fail.

c) A will fail & B will fail

d) both of them will fail

Sol:

a) $P(B \text{ fail}) = P(\bar{B}) = 0.4$

b) $P(\bar{A} \cup \bar{B}) = P(\overline{A \cap B})$

$= 1 - P(A \cap B)$

$= 1 - P(A) \cdot P(B)$

$= 1 - (0.3)(0.6) = .82$

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

$\overline{A \cup B}$

c* ~~$A \not\subseteq B$~~ will fail

~~$P(\overline{A \cup B})$~~ $= P(\bar{A} \cap \bar{B})$

$= P(\bar{A}) \cdot P(\bar{B})$ $= (0.7)(0.4)$

~~X d)~~ ~~$P(\overline{A \cap B})$~~ $= 1 - P(A \cap B)$

$= 1 - P(A) \cdot P(B)$

$= 1 - (0.3)(0.6)$

6 Exception law Prob. :- قانون الطرح

$P(A - B) = P(A \cap \bar{B})$

$= P(A) - P(A \cap B)$ بنوخذ P(A) عامل مشترك

$\sim P(\bar{A} \cap B) = P(B) - P(A \cap B)$



مثل سي if
عنطقت التقاطع

* IF A,B are indep. , then

$P(A \cap \bar{B}) = P(A) - P(A \cap B)$

$= P(A) - P(A)\,P(B)$

$= P(A) \cdot (1 - P(B))$

$= P(A) \cdot P(\bar{B})$

المنطقة اللي
تحتوي على نونيز
من الخطر



لو كان عنا مجموعة تقاطع مجموعة ثانية Complement بتكون هي بدها المجموعة اللي ما

عليها Complement ناقص اللي تحت Complement

Scanned with CamScanner

Scanned with CamScanner

6 Exception law prob. قانون الطرح

$$P(A-B) = P(A \cap \bar{B})$$
$$= P(A) - P(A \cap B)$$
$$\sim P(\bar{A} \cap B) = P(B) - P(A \cap B)$$



المنطقة المظللة هي $P(A \cap \bar{B})$

* IF A,B are indep., then

$$P(A \cap \bar{B}) = P(A) - P(A \cap B)$$
$$= P(A) - P(A)\,P(B)$$
$$= P(A).(1 - P(B))$$
$$= P(A).P(\bar{B})$$

المنطقة التي تحتوي على توحيد هي حاصل ضرب



Scanned with CamScanner

No.

Ex1 If $P(A) = 0.2$, $P(\bar{A} \cap B) = 0.3$,
find $P(A \cup B)$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
$$= P(A) + P(\bar{A} \cap B)$$
$$= 0.2 + 0.3 = 0.5$$

Ex2 if $P(A) = 0.4$, $P(A \cap \bar{B}) = 0.1$
Find $P(\bar{A} \cup \bar{B})$

$$P(\bar{A} \cup \bar{B}) = P(\overline{A \cap B})$$
$$= 1 - P(A \cap B)$$

But $P(A \cap \bar{B}) = P(A) - P(A \cap B)$
$0.1 = 0.4 - P(A \cap B)$
$P(A \cap B) = 0.3$

So, $P(\bar{A} \cup \bar{B}) = 1 - P(A \cap B)$
$= 1 - 0.3 = 0.7$

Ex3 If $P(A \cup B) = 0.9$, $P(A) = 0.7$, $P(B) = 0.4$ Find $P(A \cup \bar{B})$

$0.9 = P(A \cup B) = P(A) + P(B) - P(A \cap B)$
$0.9 = 0.7 + 0.4 - P(A \cap B)$
$\rightarrow$ $P(A \cap B) = 0.2$

Now; $P(\overline{A \cup B}) = P(\bar{A} \cap \bar{B}) = P(\bar{A} \cap B)$
$= P(B) - P(A \cap B)$
$= 0.4 - 0.2 = 0.2$

مشطوب

smile

Scanned with CamScanner

Scanned with CamScanner

قانون الشرط : احتمالية اشي شرط اشي ثاني يساوي احتمالية التقاطع على احتمالية الشرط

[7] Conditional probability: الاحتمال المشروط

~ P(A), P(B), P(C) ... Marginal prob. الاحتمال اللي يعتبر احتمال حادث واحد (هامشي).

~ P(A∩B) ≡ P(AB) = P(ABC), P(AC), ... → Joint Prob. احتمالية حدوث حادثين.

~ P(A|B), P(B|C), P(A|C), ... → Conditional Prob.
given

[*1] P(A|B) = Pro. of A given B = Prob. of event A s.t event B is done

تعني احتمال الحادث الذي يقع خلف | بشرط حدوث الثاني.

↝ If A, B are indep. then لا يؤثر الحادث بعضهم

P(A|B) = P(A)

هاي الشحطة تعبر عن الـ indp

[*1] P(A|B) = P(A∩B) / P(B) ... في المقام يكون احتمال وشرط

↳ P(A∩B) = P(A|B) · P(B)

P(B|A) = P(B∩A) / P(A) → P(A∩B) = P(B|A) · P(A)

↝ If A, B are indep. then:

P(A|B) = P(A∩B) / P(B) = P(A) · P(B) / P(B) = P(A)

الشرط بنشطب لنه ما اله معنى بسبب انه ما اله علاقة على الـ A

Scanned with CamScanner

نستطيع معرفة لو كانت الاحتمالية لا تعتمد على الشرط عن طريق جواب ضرب الاحتمالية الأولى بالشرط مثلا لو ضربنا بالمثال الاتي
P(A) x P(B) الجواب 0.2" وهيك بعرف غير معتمدين عن بعض عنه لجواب بساوي ناتجهم

Ex] In some population, it's found that
$P(A) = 0.4$, $P(B) = 0.5$, $P(A \cap B) = 0.2$
where A denotes to the hypertension ارتفاع ضغط الدم
& B denotes to the over weight
If a person is chosen at random, find the prob. that he has a hypertension Sol! he has an over weight.

Sol:
$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.2}{0.5} = 0.4$$

حادثين مستقلين

$P(A|B) = P(A)$, So, A is indep of B كحادثين منفصلين

If $P(A) = 0.2$ → لو كانت ~ → يكونو بينهم علاقة → dep.

* Rules:

1. $P(\phi|A) = 0$
2. $P(\Omega|A) = 1$
3. $P(\bar{A}|B) = 1 - P(A|B)$

احتمالية خاي بدون شرط

Proof:

$$P(\bar{A}|B) = \frac{P(\bar{A} \cap B)}{P(B)}$$

$$= \frac{P(B) - P(A \cap B)}{P(B)}$$ ← وزعنا البسط على المقام

$$= 1 - \frac{P(A \cap B)}{P(B)}$$ الاحتمال الشرطي

$$= 1 - P(A|B)$$ #

4. $P(A|\bar{B}) \neq 1 - P(A|B)$

$$P(A|\bar{B}) = \frac{P(A \cap \bar{B})}{P(\bar{B})}$$

$$= \frac{P(A) - P(A \cap B)}{1 - P(B)}$$

مشطوب

smile
Scanned with CamScanner
Scanned with CamScanner

No.

* permutations and combinations:- التباديل والتوافيق والتجاميع

* No. of ways of chosing r elements out of n elements s.t. the order is important is denoted by

$$nP_r = \frac{n!}{(n-r)!}$$

تعد العناصر اذا كانة الترتيب مهم نستخدم

* No. of ways of chosing r elements out of n s.t the order is not important is denoted by

$$nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Ex| How many 3 digits car plates could you form from $\{0,1,2,\dots,9\}$

→ No. of plates $= (10)^3$

Ex| How many car plates could you form from $\{0,1,\dots,9\}$ without duplicated any number

بدون تكرار

no. of plates $= 10 \times 9 \times 8 = \boxed{10P3} = \frac{10!}{(10-3)!} = \frac{10!}{7!} = \frac{10\times9\times8\times7!}{7!}$

= Chosing 3 out of 10 without replacement

smile

Scanned with CamScanner

Scanned with CamScanner

ترتيب غير مهم

Ex) In the exp. of drawing two balls from a box with 3 red and 2 black balls,

a) Find no. of ways of choosing 1 red ball
b) find no. of ways of Choosing 2 red balls.
c) find no. of ways of Choosing 2 balls.
d) find the prob. of Choosing 1 red ball
e) find the prob. of Choosing 2 red balls.

Sol :-

$\Omega = \{R_1R_2, R_1R_3, R_2R_3, R_1B_1, R_1B_2, R_2B_1, R_2B_2, R_3B_1, R_3B_2, B_1B_2\}$



a) No. of way of choosing 1 red ball is $= \binom{3}{1}\binom{2}{1} = 6$

3 choose 1

b) No. of ways of choosing 2 red $= \binom{3}{2}\binom{2}{0} = 3$

c) No. of ways of choosing 2 balls $= \binom{5}{2} = 10 = \#$ dem

d) $p(1\text{ red}) = \frac{\#(1\text{ red})}{\#(\Omega)} = \frac{\binom{3}{1}\binom{2}{1}}{\binom{5}{2}} = 0.3$

Scanned with CamScanner

Scanned with CamScanner

No.

e) $P(2\text{ red}) = \frac{\#(2\text{ red})}{\#\Omega} = \frac{\binom{3}{2}\binom{2}{0}}{\binom{5}{2}} = 0.3$

[٧] 10 students — بدي اختار 3 طلاب عشوائيا



a) Find the prob. of choosing 2 sci student.

$$P(2S) = \frac{\binom{5}{2}\binom{5}{1}}{\binom{10}{3}}$$

b) Find the prob. of choosing at least 2 Eng. students

٣ أو ٢ من الاثنين

$$P(\text{at least } 2E) = P(2E) + P(3E)$$

$$= \frac{\binom{3}{2}\binom{7}{1}}{\binom{10}{3}} + \frac{\binom{3}{3}\binom{7}{0}}{\binom{10}{3}}$$

c) Find the prob. of choosing 1 sci. and at least 1 Eng.

$$P(1S \cap \text{at least } 1E) = P(1S \cap \text{at } 1E) + P(1S \cap 2E)$$

$$= \frac{\binom{5}{1}\binom{3}{1}\binom{2}{1}}{\binom{10}{3}} + \frac{\binom{5}{1}\binom{3}{2}\binom{2}{0}}{\binom{10}{3}}$$

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

d) Find the prob. of chosing at least 1 Medical Student

$$P(\text{at least } 1M) = P(1M) + P(2M)$$

$$= \frac{\binom{2}{1}\binom{8}{2}}{\binom{10}{3}} + \frac{\binom{2}{2}\binom{8}{1}}{\binom{10}{3}}$$

* Random Variables:-

Def: The Random Variables (r.v) is a function defined from Ω to $\mathbb{R}$.

Range

i.e $X: \Omega \longrightarrow \mathbb{R}$ → real no. ج

الأعداد

~ The Range of the r.v X is denoted by R_X which is the set of all possible values of the r.v X.

e.g) X : No. of student who pass the exam o

$X = \{0, 1, \dots, 10\} = R_X$

Scanned with CamScanner

Scanned with CamScanner

No.

* The r.v's have two main types?

(I) Discret r.v : الفيزياء

↳ It takes values as distinct no.'s.

e.g. Y: No. of heads in the exp. of tossing a coin 5 times

$Y = \{0, 1, 2, \ldots, 5\}$

(II) Cont. r.v : الفيزياء

It takes values as intervals.

e.g. Z: The weight of boxing championship.

$Z = \{(49, 52), (52, 56), \ldots, (91, 206)\}$.

Z_1 Z_2

Def:

The prob. density function (p.d.f): is denoted by f (maybe f(x) in case of cont. r.v's) which is prob. function satisfies the following :-

2

Scanned with CamScanner

Scanned with CamScanner

Cont. / discrete كالسّ في pdf لـ حواهب *

↳ discrete:

1) $P(x) \geq 0$, $\forall x \in R_X$

2) $\sum_{x \in R_X} P(x) = 1$ مجموعة من النقاط تجمع

3) $P(X \in A) = \sum_{x \in A} P(x)$, where $A \subseteq R_X$

~~4) $P(X \in A) \cap (X \in B) = P(X \in A \cap B) = \sum_{x \in A \cap B} P(x)$~~

5) $P(X \in A \mid X \in B) = \frac{P(X \in A \cap B)}{P(X \in B)}$

↳ Cont. :-

1) $f(x) \geq 0$, $\forall x \in R_X$

لـ interval فترة

2) $\int_{R_X} f(x)\,dx = 1$ تتكون من فترة او عدد لانهائي من الفترات

3) $P(X \in A) = \int_{(A)} f(x)\,dx$

فترة او اتحاد فترات

4) $P(a \leq x \leq b) = \int_a^b f(x)\,dx$

5) $P(X = a) = \int_a^a f(x)\,dx = 0$

Scanned with CamScanner

Scanned with CamScanner

↳ discrete :-

6) The expected value of the r.v X is denoted by:

$$EX = \sum_{x \in R_x} x\,p(x)$$

7) $EX^2 = \sum_{x \in R_x} x^2 p(x)$

$EX^3 = \sum_{x \in R_x} x^3 p(x)$ / $Eg(x) = \sum_{x \in R_x} g(x)\,p(x)$

8) $E(aX+b) = aEx + b$ / $E(ax^2+bx+c) = aEx^2 + bEX + C$

9) $\text{Var } X = \sigma_x^2 = EX^2 - (EX)^2$

10) $\text{Var}(ax+b) = a^2 \text{Var} X$

↳ Cont. :

مطلوب

6) $P(x \in A \cap x \in B) = \int_{A \cap B} f(x)\,dx$

7) $P(x \in A \mid x \in B) = \dfrac{P(x \in A \cap B)}{P(x \in B)} = \dfrac{\int_{A \cap B} f(x)\,dx}{\int_B f(x)\,dx}$

8) $EX = \int_{R_x} x\,f(x)\,dx$

9) $EX^2 = \int_{R_x} x^2 f(x)\,dx$ / $EX^3 = \int_{R_x} x^3 f(x)\,dx$ / $Eg(x) = \int_{R_x} g(x) f(x)\,dx$

10) $E(aX+b) = aEx + b$ / $E(ax^2+bx+c) = aEx^2 + bEx + C$

11) The variance of the r.v. X is

$\text{Var } X = \sigma_x^2 = EX^2 - (EX)^2$

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

الانحراف المعياري
Standard deviation

Discrete:

11) $std(x) = \sqrt{\sigma_x^2}$ → varian

12) $std(ax+b) = |a|\, std(x)$ → الـ B يروح

مثال:
لو قدمنا امتحان وكان في خطا واجا الدكتور رفعنا كلنا
10 علامات شو بأثر بتباين العلامات ؟
ولا شي بنهضلزي هو

Standard deviation
لو ربعناه بعطيني الـ **variance**

Cont:

12) $Var(aX+b) = a^2 VarX$

13) $std(x)$ = standard deviation of $x = \sqrt{\sigma_x^2}$

14) $std(ax+b) = |a|\, std(x)$.

Scanned with CamScanner

Scanned with CamScanner

#Fida'a_Abbadi

No.

11/10/2017

* EX ≡ expectation of X
≡ mean of x = average of x ≡ $\bar{x}$ (المتوسط الحسابي)

~ If $y = ax + b$, then

1) $Ey = E(ax+b) = aEX + b$

$\bar{y} = a\bar{x} + b$

2) $Var\ y = Var(ax+b)$

$= a^2\ Var\ X$.

(التشتت / التباين)

لا يتأثر بالمقدار الثابت المضاف للمتغير العشوائي X

e.g:-



Scanned with CamScanner

Scanned with CamScanner

* $\binom{n}{r} = \binom{n}{n-r}$ $\binom{10}{1} = \binom{10}{9}$ $\binom{10}{2} = \binom{10}{8}$ ...

No.

Ex) Two balls are drawn from a box with 3 red and 2 black balls. let x be the no. of red balls. Find every things:



$X = \{0, 1, 2\}$

احتمال اصحاب ←

$P(X=0) = P(0) = \frac{\binom{3}{0}\binom{2}{2}}{\binom{5}{2}} = 0.1$ کدشش ... سودای دو قرمز

$P(X=1) = P(1) = \frac{\binom{3}{1}\binom{2}{1}}{\binom{5}{2}} = 0.6$

$P(X=2) = P(2) = \frac{\binom{3}{2}\binom{2}{0}}{\binom{5}{2}} = 0.3$

* $\sum_{x \in R_x} P(x) = P(0) + P(1) + P(2) = 1$

X	0	1	2	
P(x)	0.1	0.6	0.3	1

← Prob. distribution table

جدول توزیع احتمالی ← مجموع احتمال

discrete

X	P(x)	X P(x)	$X^2 P(x)$	$X^3 P(x)$
0	0.1	0	0	0
1	0.6	0.6	0.6	0.6
2	0.3	0.6	1.2	2.4
مجموع	1	1.2	1.8	3

Scanned with CamScanner

Scanned with CamScanner

No.

$\rightsquigarrow Ex = \sum_{x \in Rx} x p(x) = 1.2$

$\rightsquigarrow Ex^2 = \sum_{x \in Rx} x^2 p(x) = 1.8$

$Ex^3 = \sum_{x \in Rx} x^3 p(x) = 3$

$\rightsquigarrow$ $\text{Var } X = Ex^2 - (Ex)^2 = 1.8 - (1.2)^2 = 0.36$

$\rightsquigarrow$ $\text{std}(x) = 0.6 = \sqrt{0.36}$

- If $y = -2x + 3$, then

$Ey = E(-2x+3) = -2Ex + 3$

$= (-2)(1.2) + 3 = 0.6$

$\rightsquigarrow$ $\text{Var } y = \text{Var}(-2x+3) = 4 \text{ var } x = (4)(0.36) = 1.44$

$\rightsquigarrow$ $\text{std}(y) = \text{std}(-2x+3) = |-2| \text{ std}(x)$

$= (2)(0.6) = 1.2$

Scanned with CamScanner

Scanned with CamScanner

You can do it

No.

ما الـ dis. كانت اذا كانت Can او

Ex) let X be a r.v. with $EX = 2$, $\text{Var}X = 3$
find $E(-2X^2 + 3X - 5)$

Sol:

$$3 = \text{Var}X = EX^2 - (EX)^2$$
$$3 = EX^2 - (2)^2 \longrightarrow EX^2 = 7$$
$$\text{So, } E(-2X^2 + 3X - 5)$$
$$= -2EX^2 + 3EX - 5$$
$$= (-2)(7) + (3)(2) - 5 = -13$$

Ex): let X be a discrete r.v with range $R_X = \{1, 2, 3\}$ s.t $EX = 2$, $\text{Var}X = 1.3$, find $p(X \in A)$; $A = \{1, 3\}$.

Sol: $EX^2 = \text{Var}X + (EX)^2$

$EX^2 = 1 + 4 = 5$

$2 = EX = \sum_{x \in R_X} x p(x)$

$5 = EX^2 = \sum_{x \in R_X} x^2 p(x)$

$5 = (1)p(1) + 4p(2) + 9p(3)$ — (2)

$\sum_{x \in R_X} p(x) = 1$

$p(1) + p(2) + p(3) = 1$ — (3)

محلول غلط الدكتور حله بالصفحات الجاي

الفكرة: انه احنا بنعوف توقع X^2 فنستخدم الحذف والتعويض عشان نطلع احتمالية {1,2,3}

(1)

Scanned with CamScanner

Scanned with CamScanner

You strong enough

No.

$P(1) = 0.5$, $P(2) = 0$, $P(3) = 0.5$

$P(x \in \{1,3\}) = P(1) + P(3) = 0.5 + 0.5 = 1$

Ex) let X be a discrete r.v with p.d.f

$$P(x) = \begin{cases} c, & x = 1, 2, 3, 4 \\ 0.1, & x = 5, 6 \\ 0, & o.w \end{cases}$$

Find c.

Sol:

$$\sum_{x \in R_X} p(x) = 1$$

$$P(1) + P(2) + P(3) + P(4) + P(5) + P(6) = 1$$

$$C + C + C + C + 0.1 + 0.1 = 1$$

$$4C = 0.8 \qquad \boxed{C = 0.2}$$

Ex): let x be a Cont. r.v with p.d.f

$$F(x) = \begin{cases} c, & 0 \le x < 1/2 \\ 2c, & 1/2 < x \le 1 \\ 0, & o.w \end{cases}$$

Find c.

Sol:

$$\int_{R_X} f(x)\,dx = 1 \longrightarrow \int_0^{1/2} c\,dx + \int_{1/2}^{1} 2c\,dx = 1$$

$$\tfrac{1}{2}c + 2\left(\tfrac{1}{2}c\right) = 1$$

$$\tfrac{1}{2}c + c = 1$$

$$\tfrac{3}{2}c = 1 \qquad \boxed{c = \tfrac{2}{3}}$$

smile for life

Scanned with CamScanner

Scanned with CamScanner

Eg. X discrete r.v $x \in [1,2,3]$

$E(X) = 2$, $Var(X) = 1$

find $P(x \in A)$ where $A = [1,3]$

Sol.

$$Var(X) = E(X^2) - (E(X))^2$$

$$\rightarrow 1 = E(X^2) - 2^2$$

$$\rightarrow E(X^2) = 5 .$$

Since $2 = E(X) = \sum_{x \in [1,2,3]} x\, p(x) = (1)\,p(1) + (2)\,p(2) + 3\,p(3) = 2$

$$\rightarrow \boxed{p(1) + 2p(2) + 3p(3) = 2} \quad (1)$$

Since $E(x^2) = 5$, we have

$$E(x^2) = \sum x^2 p(x) = \boxed{p(1) + 4p(2) + 9p(3) = 5} \quad (2)$$

Since $\sum p(x) = 1 \rightarrow \boxed{p(1) + p(2) + p(3) = 1} \quad (3)$

Scanned with CamScanner

Scanned with CamScanner

by solving ①, ② & ③, we have

$p(1) = \frac{1}{2}$, $p(2) = 0$, $p(3) = \frac{1}{2}$

$\rightarrow p(x \in \{1,3\}) = p(1) + p(3)$

$= \frac{1}{2} + \frac{1}{2} = 1$

Scanned with CamScanner

Scanned with CamScanner

No.

Ex) let X be a Cont. r.v with p.d.f :-

$$f(x) = \frac{3}{2}C^2 e^{-cx}, \quad 0 < x < \infty$$

a) Find C

$$\int_{Rx} f(x)\,dx = 1 \longrightarrow \int_0^\infty \frac{3}{2}C^2 e^{-cx}\,dx = 1$$

$$\longrightarrow c^2 \frac{-1}{c} e^{-cx}\Big]_0^\infty = 1$$

C عدد موجب

$$-c\, e^{-cx}\Big]_0^\infty = 1 \qquad -c[0-1] = 1 \longrightarrow \boxed{C = +1}$$

b) find $P(X \in (0,1))$

$$P(X \in (0,1)) = P(0 < x < 1)$$

$$= \int_0^1 f(x)\,dx = \int_0^1 e^{-x}\,dx = -e^{-x}\Big]_0^1$$

$$= -(e^{-1} - e^{0}) = 1 - 1/e$$

$f(x) = e^{-x}$

احتمالية اكس تنتمي لفترة هي التكامل على الفترة

- $P(X \in (-1,2) \cap X \in (1,15))$

$$= P(X \in (1,2)) = \int_1^2 e^{-x}\,dx = \text{as part b.}$$

Scanned with CamScanner

Scanned with CamScanner

No.

16/10/2017

c) Find $P(X=2)$

$$P(X=2) = \int_2^2 f(x)\,dx = 0.$$

كس نس اوي عدد احتماليتهابتصيد 0 لنلاون ص24 اخر سطر

d) Find EX

$$Ex = \int_{R_x} x\, f(x)\,dx = \int_0^\infty x\, e^{-x}\,dx$$

$$= -x e^{-x} - e^{-x} \Big]_0^\infty = 0 - (0-1) = 1$$

e) Find Ex^2

$$Ex^2 = \int_0^\infty x^2 e^{-x}\,dx \qquad (*) \qquad (*)$$

$$= -x^2 e^{-x} - 2x e^{-x} - 2e^{-x} \Big]_0^\infty$$

$$= 0 - (0 - 0 - 2) = 2$$

f) Find $E(2x^2 - 3x + 5)$

$$E(2x^2 - 3x + 5)$$

$$= 2Ex^2 - 3Ex + 5$$

$$= (2)(2) - (3)(1) + 5 = 6$$

Scanned with CamScanner

Scanned with CamScanner

g) Find Var (x)

$\text{Var } X = EX^2 - (EX)^2 = 2 - (1)^2 = 1$

h) Find $E(2X-3)$, Var $(2X-3)$

$E(2X-3) = 2EX - 3$

$= (2)(1) - 3 = -1$

$\text{Var}(2X-3) = (2)^2 \text{Var} X = (4)(1) = 4$.

Ex) let x be a Cont. r.v with p.d.f

$$f(x) = \begin{cases} C, & 0 \le x < 1 \\ 2C, & 1 \le x < 2 \\ 0, & \text{o.w} \end{cases}$$

H.W

a) Find c b) Find $P(X=1)$

c) Find $P(\frac{1}{2} < x < 1.5)$

d) Find EX, EX^2, $\text{Var } X$

e) Find $E(2-3X)$, $\text{Var}(2-3X)$.

Scanned with CamScanner

Scanned with CamScanner

Def: let X be a r.v. with range R_x, then the Cumulative distribution function (c.d.f) ← التوزيع التراكمي or Simply the distribution function (d.f) is given by

$$F(x) = \begin{cases} \sum_{u \le x} p(u) & ; \ X \text{ is discrete} \\ \int_{-\infty}^{x} f(u)\,du & ; \ X \text{ is Cont.} \end{cases}$$

هون حطينا حرف U لنه ما بصير يكون حد التكامل جوا القوسين f(x)

Ex) let X be a discrete r.v with the following:-

X	0	1	2
P(x)	1/4	1/2	1/4



a) Find the c.d.f
b) Find $P(1 \le x < 2)$
c) Find $p(0.5 < x < 1.5)$

Sol:

a) $F(x) = 0 \ ; \ x < 0$

$F(x) = 1/4 \ ; \ 0 \le x < 1$

$F(x) = \frac{1}{4} + 1/2 \ ; \ 1 \le x < 2$

$F(x) = \frac{1}{4} + 1/2 + \frac{1}{4} \ ; \ 2 \le x$

$$\rightarrow F = \begin{cases} 0 & ; \ x < 0 \\ \frac{1}{4} & ; \ 0 \le x < 1 \\ \frac{3}{4} & ; \ 1 \le x < 2 \\ 1 & ; \ x \ge 2 \end{cases}$$

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

b) $P(1 \leq x \leq 2) =$

$= P(1) + P(2) = \frac{1}{2} + \frac{1}{4} = \frac{3}{4}$

علامة اليساوي مهمة بالدسكريت
بس مش مهمة بالـ cont

c) $P(0.5 < x < 1.5) =$

$= P(1) = \frac{1}{2}$

* Properties:-

1) F(x) is a step function when x is discrete

2) F(x) is Cont. when x is Cont.

3) $F(-\infty) = 0$, $F(\infty) = 1$

4) $P(a \leq x \leq b) = F(b) - F(a)$ if x Cont. r.v

5) $f(x) = \frac{d}{dx}(F(x))$; if x is Cont. cdf is pdf. cont.

6) $F(x_1) \leq F(x_2)$; if $x_1 \leq x_2$

i.e. F is increasing.

Scanned with CamScanner

Scanned with CamScanner

No.

Ex) let x be a discrete r.v with c.d.f

$$F(x) = \begin{cases} 0 & x<4 \\ 1/4 & 4 \le x<5 \\ 1/2 & 5 \le x<6 \\ 1 & x \ge 6 \end{cases}$$



لو فاهمين طريقة الحل
مافي داعي للتفصيل

a) Find a p.d.F
b) Find $P(4 \le x \le 6)$

خطوات الرسم p.d.F ←

a) $P(X=4) = F(4) = 1/4$

$P(x=5) = F(5) - F(4) = 1/2 - 1/4 = 1/4$

$P(X=6) = F(6) - F(5) = 1 - 1/2 = 1/2$

$F(4.5) = 1/4$ / $P(X=4.5) = 0$ ← لانه بنزعم عند نقطه

عبارة عن ارتفاع الدرجة عند الدرجة اللي قبلها

ال p.d.F عند الدرجة ← ال c.d.F عند نفس الدرجة

b) $P(4 \le x \le 6) = P(4) + P(5) + P(6)$

$= \frac{1}{4} + \frac{1}{4} + \frac{1}{2}$

$= 1$

smile for life
Scanned with CamScanner
Scanned with CamScanner

Ex) let x be a Cont. r.v. with p.d.f

$$f(x) = e^{-\lambda x}, \quad x \in (0, \infty)$$

Find. a) The c.d.f

b) $P(1 \leq x \leq 2)$

c) $P(1 < x < 2)$

Sol:

a) $$F(x) = \int_{-\infty}^{x} f(u)\,du = \int_{0}^{x} e^{-\lambda u}\,du$$

$$= \frac{-1}{\lambda} e^{-\lambda u} \Big]_0^x = \boxed{\frac{-1}{\lambda}\left[e^{-\lambda x} - 1\right]}$$

b) $$P(1 \leq x \leq 2) = P(1 < x < 2)$$

$$= \int_1^2 f(x)\,dx$$

or

$$= F(2) - F(1)$$

$$= -\frac{1}{\lambda}\left[e^{-2\lambda} - 1\right] - \frac{1}{\lambda}\left[e^{-\lambda} - 1\right]$$

Scanned with CamScanner

Scanned with CamScanner

ملاحظة: لما يكون الاف كبيتل F بكون CDF و r.v cont ——— No. ——— CDF (F)

طرقة الحل عشان اوجد الpdf بشتق الcdf

Ex) let $F(x) = \ln(1-\alpha x)$,
find $f(x)$.

a) $f(x) = \frac{d}{dx}(F(x))$

$= -\frac{-\alpha}{1-\alpha x}$

$= \frac{\alpha}{1-\alpha x}$

b) Find $P(0<x<0.5) =$

$P(0<x<0.5) = \int_0^{0.5} f(x)\,dx$

$= \int_0^{0.5} \frac{\alpha}{1-\alpha x}\,dx$

or $= F(0.5) - F(0)$

$= 1 - \ln(1-0.5\alpha) - (1 - \ln(1))$

$= -\ln(1-0.5\alpha)$.

c) Find $P(X=0.25) = 0$

smile for life

Scanned with CamScanner

Scanned with CamScanner

Wednesday

18-10-2017

No.

عنا 2 ر**r.v** ممكن يكونو ال2 **Discrete** او 2 **cont** وممكن يكون حالة تالتة بس مش مطلوب منا غير الحالتين

* Bivariate random variables :-

Def:- let x, y be two discrete r.v's with ranges R_x, R_y, then the two dimensional range of x, y is given by

هون راح نتعامل مع ازواج مرتبة

$$R_{xy} = \{(x,y) \; ; \; x \in R_x, \; y \in R_y\}$$ discrete

الرسمات مش كثير مهمة



Cont. $R_{xy} = \{(x,y)$
$1<x<2$
$2<y<3$



Range الـ x / x نقاط

أصغر وحدة نتعامل معها هي النقطة

* أصغر وحدة مساحة نتعامل معها هي وحدة واحدة

$(dx) = \lim \Delta x$

يتكامل على أصغر وحدة . وهو ما يقترب عند الصفر .

Scanned with CamScanner

Scanned with CamScanner

بستخدم "joint" عشان
اجمع بين p.d.f تبع
X & Y

- with joint p.d.f :-

$$f(x,y) = P_{\text{R}}\{X=x \cap Y=y\}$$

which satisfies the following :-

بنعوض كل النقاط بالاقتران X & Y
وبنجمع الأجوبة هاي بحالة Discrete
اما بالـ cont بصير دبل انتقريشن بدنا
نكامل بالنسبة لـ X وبالنسبة لـ Y

1) $f(x,y) \geq 0, \forall (x,y) \in R_{xy}$.

لـ في حالة
تقطع
Region

2) $\sum_{x\in R_x}\sum_{y\in R_y} f(x,y) = 1$, $\iint_{R_{xy}} f(x,y)\,dx\,dy = 1$

$(x,y)\in R_{xy}$

cont.

3) The marginal p.d.f. of X is given by

$$f_x(x) = \sum_{y\in R_y} f(x,y).$$

cont.

$$f_x(x) = \int_{R_y} f(x,y)\,dy$$

$f(x,y) = x+y$
$\int (x+y)\,dy$
$= xy + \frac{y^2}{2}$
$= x + \frac{1}{2} - 0$

هون بس نكامل بالنسبة لـ X
بنعتبر الـ Y ثابت والعكس
صحيح

4) The marginal p.d.f of Y is

cont

$$f_y(y) = \sum_{x\in R_x} f(x,y)$$

$$f_y(y) = \int_{R_x} f(x,y)\,dx.$$

اذا بدي الـ marginal
p.d.f بالنسبة للـ Y
بكامل على X والعكس

smile for life

Scanned with CamScanner

Scanned with CamScanner

لما يطلب اشي بدلالة الـ X
ممكن نستعمل marginal p.d.f
للـ X واذا طلب للـ Y نفس الاشي

No.

5) $EX = \sum_{x \in R_x} x f_x(x)$ → Cont. $EX = \int_{R_x} x f_x(x)\, dx$

- $EX^2 = \sum_{x \in R_x} x^2 f_x(x)$ $\int_{R_x} x^2 f_x(x)\, dx$

* $Eg(x) = \int_{R_x} g(x) f_x(x)\, (dx)$

6) $EY = \sum_{R_y} y f_y(y)$ cont. → $\int_{R_y} y f_y(y)\, dy$

- $EY^2 = \sum_{y \in R_y} y^2 f_y(y)$ → $= \int_{R_y} y^2 f_y(y)\, dy$

7) $EXY = \sum_{x \in R_x} \sum_{y \in R_y} xy f(x,y)$

لو بدي توقع X و Y
ما بقدر استعمل
X sub X
A sub Y
لنه التوقع مكس

cont:

$$EXY = \iint_{R_{xy}} xy f(x,y)\, dx\, dy$$

- $Eg(x,y) = \iint g(x,y) f(x,y)\, (dx\, dy)$

independent Y و X اذا كان
وطلعنا
marginal p.d.f x
marginal p.d.f y
وضربناهم ببعض
بعطيني f(x,y)
سواء كان cont or Discrete

Scanned with CamScanner

Scanned with CamScanner

8) If X,y are indep. then

$$F(x,y) = P\{X=x, Y=y\}$$

joint د

حاصل ضرب د

marginal.

$$= P\{X=x\} \cdot P(Y=y\}$$

$$= f_x(x) \cdot f_y(y).$$

9) If x,y are indep. then

$$EXy = \sum_{x \in R_x} \sum_{y \in R_y} xy\, f(x,y) = \sum_{x \in R_x} \sum_{y \in R_y} xy\, f_x(x)\, f_y(y)$$

$$= \sum_{x \in R_x} x f_x(x) \sum_{y \in R_y} y f_x(y)$$

$$= (Ex)(Ey)$$

اذا كان X وY
independent الـ Ex
بتوزع على X و Y دايما الـ
Co-Var

يساوي 0 لما يكون X و Y
indep فقط!!

Scanned with CamScanner

Scanned with CamScanner

لنه لما يكونو معتمدين بكم

Monday
23/10

No.

هاض القانون بتطبق على الحالتين وممكن يكون
Sigma in Disc
Integral in Cont.

10) $Var\,X = EX^2 - (EX)^2$
$Var\,Y = EY^2 - (EY)^2$

توقع الثابت بعطيني نفسه

11) $E(aX+b) = aEX + b$
$E(cY+d) = cEY + d$

12) The Co-Variance of X, Y is given by

التباين المشترك بجمع **X** و **Y**

$Cov(X,Y) = EXY - (EX)(EY)$

↳ **Notes:**

a) $Cov(X,X) = EXX - (EX)(EX)$
$= EX^2 - (EX)^2 = Var(X)$

b) $Cov(X,Y)$ could be (-ve)

c) $Cov(aX+b, cY+d)$ لحد هون
$= (a)(c)\,Cov(X,Y)$

d) If X, Y are indep then, **Co-0**
$Cov(X,Y) = EXY - (EX)(EY)$ **Var**
$= (EX)(EY) - (EX)(EY) = 0$

13) The correlation coefficient: ← بيربط العلاقة
of X, Y is given by

$Corr(X,Y) = \rho(X,Y) = \frac{Cov(X,Y)}{\sigma_X \sigma_Y}$; $\sigma_X = \sqrt{Var}$; $\sigma_Y = \sqrt{Var}$

Independent = Uncorrelated
dependent = Correlation

Scanned with CamScanner

Scanned with CamScanner

#Hint
Correlation = Corr

*Notes: (about correlation coefficient):

اذا كانت X وY موجب 1 بتكون علاقةطرديه
واذا كان سالب بتكون علاقة قوية عكسية
وكل ما اقتربت من ال0 بتضعف العلاقة

a) $-1 \leq corr(x,y) \leq 1$

b) $corr(aX+b, cY+d)$

$$= \begin{cases} corr(x,y) & \text{if } a,c \text{ have same sign} \\ -corr(x,y) & \text{if } a,c \text{ have diff. Signs.} \end{cases}$$

اذا كانو معاملات X , Y نفس
الإشارة بنحط
موجب واذا عكس بنحط سالب

c) If x, y are indep. then

$$corr(x,y) = \frac{cov(x,y)}{\sigma x \sigma y} = 0$$

d) If $corr(x,y) \approx 1$, then



→ strong relation between x, y
(+ve direction) علاقة طردية

e) If $corr(x,y) \approx -1$, then



→ strong relation
(-ve direction) علاقة عكسية

Scanned with CamScanner

Scanned with CamScanner

f) If corr(x,y) = 0, then

g)



Scanned with CamScanner

Scanned with CamScanner

No.

Ex) Three of ~~five~~ items are defective 8-t
1 item is defective in size and 2 items
are defective in design. (DD) (Ds)
Two items are drawn randomly.

let X: no. of Ds items
Y: no. of DD items.

Find every thing:



$X = \{0, 1\}$
$Y = \{0, 1, 2\}$
$R_{X,Y} = \{(0,0), (0,1), (0,2), (1,0), (1,1), (1,2)\}$

Joint p.d.f جدول

Y \ X	0	1	Σ
0	0.1	0.2	0.3
1	0.4	0.2	0.6
2	0.1	0	0.1
Σ	0.6	0.4 →	1

$P(0,0) = \frac{\binom{1}{0}\binom{2}{0}\binom{2}{2}}{\binom{5}{2}} =$

$P(1,0) = \frac{\binom{1}{1}\binom{2}{0}\binom{2}{1}}{\binom{5}{2}} =$

$P(0,1) = \frac{\binom{1}{0}\binom{2}{1}\binom{2}{1}}{\binom{5}{2}} =$

$f(1,1) = \frac{\binom{1}{1}\binom{2}{1}\binom{2}{0}}{\binom{5}{2}} =$

$P(0,2) = \frac{\binom{1}{0}\binom{2}{2}\binom{2}{0}}{\binom{5}{2}} =$

$P(1,2) = 0$

2-way table
- joint p.d.f

مجموعهم كلهم ١
حتى يكون p.d.f

لو ما كان المجموع ١ يكون في خطأ بالحل

Scanned with CamScanner

Scanned with CamScanner

NO.

Wensday
25/10

The marginal p.d.f of X is:

X	$f_X(x)$	$x f_X(x)$	$x^2 f_X(x)$
0	0.6	0	0
1	0.4	0.4	0.4
	1	0.4	0.4

$\rightsquigarrow EX = \sum_{x \in R_X} x f_X(x) = 0.4$

$\rightsquigarrow EX^2 = \sum_{x \in R_X} x^2 f_X(x) = 0.4$

$\rightsquigarrow E(3-2x) = 3 - 2EX = 3 - (2)(0.4) = 2.2$

$\rightsquigarrow E(2x^2 + 5x - 15) = 2EX^2 + 5EX - 15 = (2)(0.4) + (5)(0.4) - 15 =$

$\rightsquigarrow VarX = EX^2 - (EX)^2 = 0.4 - (0.4)^2 = 0.24$

$\rightsquigarrow Var(2-3x) = (-3)^2 VarX = (9)(0.24) = --$

Scanned with CamScanner

Scanned with CamScanner

y	$f_Y(y)$	$y f_Y(y)$	$y^2 f_Y(y)$
0	0.3	0	0
1	0.6	0.6	0.6
2	0.1	0.2	0.4
Σ	1	0.8	1

$$EY = \sum_{y \in R_Y} y f_Y(y) = 0.8$$

$$EY^2 = \sum_{y \in R_Y} y^2 f_Y(y) = 1$$

$$\text{Var}Y = EY^2 - (EY^2) = 1 - (0.8)^2 = 0.36$$

$$EXY = \sum_{x \in R_X} \sum_{y \in R_Y} xy f(x,y)$$

$$= (0)(0) f(0,0) + (0)(1) f(0,1) + (0)(2) f(0,2) + (1)(0) f(1,0) + (1)(1) f(1,1) + (1)(2) f(1,2)$$

$$= 0.2$$

Are X, Y indep.?

$$EXY \stackrel{?}{=} (EX)(EY)$$

$$0.2 \neq (0.4)(0.8)$$

So, X, Y are not indep.

مهم

لو كانت المسلواة متحققه بكونو indep

Scanned with CamScanner

Scanned with CamScanner

(c)

$$\text{Cov}(X,Y) = EXY - (EX)(EY)$$
$$= 0.2 - (0.4)(0.8) = -0.12$$

$$\text{cov}(2X-3, 3-4Y)$$
$$= (2)(-4)\ \text{cov}(X,Y)$$
$$= (-8)(-0.12) = 0.96$$

$$\text{Corr}(X,Y) = \frac{\text{Cov}(X,Y)}{\sigma_X\ \sigma_Y} = \frac{-0.12}{\sqrt{0.24}\sqrt{0.36}} = -0.41$$

$$\text{Corr}(2X-3, 3-4Y) = -\text{Corr}(X,Y) = 0.41$$

Scanned with CamScanner

Scanned with CamScanner

Monday
30/10/17

* Joint distribution function :- (Joint cdf) :

في حالة Cont. ا. فقط

.. The joint cdf or simply the joint d.f. is given by

$$F_{XY}(x,y) = P\{X \leq x , Y \leq y\}$$

and can be calculated as:

$$F_{XY}(x,y) = \int_{-\infty}^{y}\int_{-\infty}^{x} f(u,v)\, du\, dv$$

تعويض مباشر

$$\int_0^1 e^{x^2} dx \quad \leftarrow \text{لا يمكن التكامل}$$

$$\int_0^1\int_0^x e^{x^2} dy\, dx$$

$$\int_0^1 y e^{x^2} \Big]_0^x dx$$

$$\int_0^1 x e^{x^2} dx$$

• Properties :-

1) $0 \leq F_{XY}(x,y) \leq 1$

2) $F_{XY}(-\infty,-\infty) = 0$ ← اول قيمة بالفترة

$F_{XY}(\infty,\infty) = 1$ ← اخر قيمة بالفترة

3) $F_{XY}(-\infty,y) = 0$, $F_{XY}(x,-\infty) = 0$

لو كان في سالب ما لانهاية بتصفر

4) $f(x,y) = \dfrac{\partial^2}{\partial x\, \partial y} F(x,y)$

5) $f_X(x) = \int_{-\infty}^{\infty} f(x,y)\, dy$

$f_Y(y) = \int_{-\infty}^{\infty} f(x,y)\, dx$

مهم جدا

smile for life

Scanned with CamScanner

Scanned with CamScanner

6) $F_X(x) = \int_{-\infty}^{x} f_X(u)\,du$

- $F_Y(y) = \int_{-\infty}^{y} f_Y(v)\,dv$

7) $F_X(x) = \int_{-\infty}^{x}\int_{-\infty}^{\infty} f(u,y)\,dy\,du$ → marginal p.d.f

مشطوب

- $F_Y(y) = \int_{-\infty}^{y}\int_{-\infty}^{\infty} f(x,v)\,dx\,dv$

8) $f_X(x) = \frac{d}{dx} F_X(x)$

$f_Y(y) = \frac{d}{dy} F_Y(y)$

9) $P(X \in A, Y \in B) = \iint_{B}\int_{A} f(x,y)\,dx\,dy$ هام جدا

$P(x_1 \le X \le x_2, y_1 \le y \le y_2) = \int_{y_1}^{y_2}\int_{x_1}^{x_2} f(x,y)\,dx\,dy$

10) $\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(x,y)\,dx\,dy = 1$

Scanned with CamScanner

Scanned with CamScanner

Ex: $\int_0^1 \left(\int_0^2 xy \, dy \right) dx$ "$x \in [0,1]$ & $y \in [0,2]$"

$$= \int_0^1 x \left. \frac{y^2}{2} \right|_0^2 dx$$

$$= \int_0^1 x \left(\frac{4}{2} - 0 \right) dx$$

$$= 2 \int_0^1 x \, dx = \left. x^2 \right|_0^1 = 1^2 - 0^2 = 1$$

$$\int_0^1 \int_0^2 xy \, dy \, dx = \int_0^2 \int_0^1 xy \, dx \, dy$$ note

Scanned with CamScanner

Scanned with CamScanner

Ex) let

$$f(x,y) = \begin{cases} b e^{-x} \cos y, & 0 \le x \le 2,\ 0 \le y \le \pi/2 \\ 0, & \text{O.W} \end{cases}$$

a) Find b b) Find $F_{xy}(x,y)$

c) Find $f_x(x)$, $f_y(y)$

d) Find $F_x(x)$, $F_y(y)$

e) Find $P\{0 \le x \le 1,\ 0 \le y \le \pi/4\}$

f) ~~Find $P\{0 \le x \le 1,\ y = \pi/4\}$~~

Sol:

a) $\int_0^{\pi/2}\int_0^{2} b e^{-x} \cos y \, dx\, dy = 1$

$\int_0^{\pi/2} \left(-b e^{-x} \cos y\right]_0^{2} dy = 1$

$\int_0^{\pi/2} \left(-b \cos y (e^{-2} - e^{0})\right) dy = 1$

$-b \sin y (e^{-2} - 1) \Big|_0^{\pi/2} = 1$

$-b(1)(e^{-2} - 1) - 0 = 1 \rightarrow b = \frac{-1}{e^{-2} - 1}$

$= \frac{1}{1 - e^{-2}} = \frac{1}{1 - \frac{1}{e^2}} = \frac{1}{\frac{e^2}{e^2} - \frac{1}{e^2}} = \boxed{\frac{e^2}{e^2 - 1}}$

Scanned with CamScanner

Scanned with CamScanner

No.

$$f(x,y) = \frac{e^2}{e^2-1} e^{-x} \cos y, \quad 0 \le x \le 2, \quad 0 \le y \le \pi/2$$

b) $$F_{XY}(x,y) = \int_0^y \int_0^x \frac{e^2}{e^2-1} e^{-u} \cos v \, du \, dv$$

$$= \frac{e^2}{e^2-1} \int_0^y -e^{-u} \cos v \Big]_0^x dv$$

$$= \frac{e^2}{e^2-1} \int_0^y \cos v \, (1 - e^{-x}) \, dv$$

$$= \frac{e^2}{e^2-1} (1 - e^{-x}) \sin v \Big]_0^y = \frac{e^2}{e^2-1} (1 - e^{-x}) \sin y$$

c) $$f_X(x) = \int_0^{\pi/2} \frac{e^2}{e^2-1} e^{-x} \cos y \, dy$$

$$= \frac{e^2}{e^2-1} e^{-x} \sin y \Big]_0^{\pi/2}$$

$$= \frac{e^2}{e^2-1} e^{-x}$$

$$f_Y(y) = \int_0^2 \frac{e^2}{e^2-1} e^{-x} \cos y \, dx$$

$$= \frac{-e^2}{e^2-1} \cos y \, e^{-x} \Big]_0^2 = \frac{-e^2}{e^2-1} \cdot \cos y \, (e^{-2} - 1)$$

Scanned with CamScanner

Scanned with CamScanner

No.

d) $F_X(x) = \int_0^x f_X(u)\,du$

$$= \frac{e^2}{e^2-1}\int_0^x e^{-u}\,du$$

$$= \frac{-e^2}{e^2-1}\, e^{-u}\Big]_0^x$$

$$= \frac{-e^2}{e^2-1}\left(e^{-x}-1\right)$$

$$F_Y(y) = \int_0^y \frac{-e^2}{e^2-1}\left(e^{-2}-1\right)\cos v\,dv$$

$$= \frac{-e^2}{e^2-1}\left(e^{-2}-1\right)\sin v\Big]_0^y$$

$$= \frac{-e^2}{e^2-1}\left(e^{-2}-1\right)\sin y$$

Scanned with CamScanner

Scanned with CamScanner

Unit step function

$$u(x) = \begin{cases} 1, & x \geq 0 \\ 0, & x < 0 \end{cases}$$

$$u(x - x_0) = \begin{cases} 1, & x \geq x_0 \\ 0, & x < x_0 \end{cases}$$

Note:

$$f(x)\,u(x) = \begin{cases} f(x), & x \geq 0 \\ 0, & x < 0 \end{cases}$$

$$f(x)\,u(x-1) = \begin{cases} f(x), & x \geq 1 \\ 0, & x < 1 \end{cases}$$

Impulse fn.

$\delta(x)$ satisfies impulse fn.

The

$$\int_{-\infty}^{\infty} \varphi(x)\,\delta(x - x_0)\,dx = \varphi(x_0)$$

Scanned with CamScanner

Scanned with CamScanner

Ex. $\int_{-\infty}^{\infty} x^2 \delta(x-2)\,dx = x^2\Big|_{x=2} = 4$

$\int_{-\infty}^{\infty} \delta(x)\,dx = 1$

MID

Scanned with CamScanner

Scanned with CamScanner

unit step fn.

$$u(x) = \begin{cases} 1 , & x \geq 0 \\ 0 , & x < 0 \end{cases}$$



$$f(x)\,u(x) = \begin{cases} f(x), & x \geq 0 \\ 0 , & x < 0 \end{cases}$$

$$u(x-c) = \begin{cases} 1 , & x \geq c \\ 0 , & x < c \end{cases}$$

$$f(x)\,u(x-c) = \begin{cases} f(x) , & x \geq c \\ 0 , & x < c \end{cases}$$

$$u(c-x) = \begin{cases} 0 , & x > c \\ 1 , & x \leq c \end{cases}$$



هوم نكمم أيس نيبطخ ابا نيبطخ...

1. Find the generalize p.d.f
2. Find the p.d.f imbalance

Scanned with CamScanner

Scanned with CamScanner

unit impulse function

$$\int_{-\infty}^{\infty} \varphi(x)\,\delta(x-c)\,dx = \varphi(c)$$

$$\int_{-\infty}^{\infty} \delta(x-c)\,dx = 1$$

Note: $\dfrac{du(x)}{dx} = \delta(x)$

Note:- Let $x \in \{x_1, x_2, \cdots, x_n\}$ be

Discrete r.v.

→ The p.d.f

$$f(x) = p_r(x_1)\,\delta(x-x_1) + p_r(x_2)\,\delta(x-x_2) + \cdots + p_r(x_n)\,\delta(x-x_n)$$

& the c.d.f

$$F(x) = p_r(x_1)\,u(x-x_1) + p_r(x_2)\,u(x-x_2) + \cdots + p_r(x_n)\,u(x-x_n)$$

Scanned with CamScanner

Scanned with CamScanner

الدكتور بده يعيده وبدي اعدله..

Ex. X discrete r.v.

$S = \{1, -1\}$

$f(1) = \frac{1}{2}$ & $f(-1) = \frac{1}{2}$

$\rightarrow f(x) = \frac{1}{2}\delta(x-1) + \frac{1}{2}\delta(x+1)$

$F(x) = \frac{1}{2}u(x-1) + \frac{1}{2}u(x+1)$



Ex. X cont. r.v. with c.d.f

$F(x) = u(x)\left(1 - e^{-x^2/b}\right)$

. Then

Scanned with CamScanner

Scanned with CamScanner

Ex.

$$f(x,y) = u(x)\,u(y)\,x\,e^{-x(y+1)}$$

find $f_x(x)$, $f_y(y)$, $F_x(x)$, $F_y(y)$

$F_{xy}(x,y)$,

Sol.

$$f_x(x) = \int_{-\infty}^{\infty} f(x,y)\,dy$$

$$= \int_{-\infty}^{\infty} u(x)\,u(y)\,x\,e^{-x(y+1)}\,dy$$

$$= \int_{0}^{\infty} u(x)\,x\,e^{-x(y+1)}\,dy$$

$$= u(x)\,x\,e^{-x}\int_{0}^{\infty} e^{-xy}\,dy$$

$$= u(x)\,x\,e^{-x}\left(\frac{1}{-x}e^{-xy}\Big|_0^{\infty}\right)$$

$$= u(x)\,e^{-x}\,(1-0)$$

$$= u(x)\,e^{-x}$$

Scanned with CamScanner

Scanned with CamScanner

$$f_y(y) = \int_{-\infty}^{\infty} u(x)\,u(y)\,x\,e^{-x(y+1)}\,dx$$

$$= \int_{0}^{\infty} u(y)\,x\,e^{-x(y+1)}\,dx$$

$$= u(y)\left[\frac{-x}{y+1}e^{-x(y+1)} - \frac{1}{(y+1)^2}e^{-x(y+1)}\right]_0^{\infty}$$

$$= u(y)\left[0 - 0 - 0 + \frac{1}{(y+1)^2}\right)$$

$$= \frac{u(y)}{(y+1)^2}$$

x		$e^{-x(y+1)}$	
	$+$		$-\frac{1}{y+1}e^{-x(y+1)}$
1		$e^{-x(y+1)}$	
	$-$		$\frac{1}{(y+1)^2}e^{-x(y+1)}$
0			

Scanned with CamScanner

Scanned with CamScanner

No.

- $F_X(x) = \int_0^x f_X(u)\, du$

$= \int u(x)\, e^{-x}\, dx$

$= \int_0^x e^{-u}\, du \longrightarrow -e^{-u}\Big]_0^x = -(e^{-x} - 1) = 1 - e^{-x}$

- $F_Y(y) = \int u(y) \frac{1}{(y+1)^2}\, dy$

$= \int_0^y \frac{1}{(v+1)^2}\, dv = \int_0^y (v+1)^{-2}\, dv$

$= \frac{(v+1)^{-1}}{-1}\Big]_0^y \longrightarrow \frac{(y+1)^{-1}}{-1} - \frac{1}{-1}$

$= 1 - \frac{1}{y+1}$

- $F_{XY}(x,y) = \int_0^y \int_0^x u\, e^{-u(v+1)}\, du\, dv.$ ⋮

Scanned with CamScanner

Scanned with CamScanner

* Prob. Distribution:-

عدد مرات التجربة
بنسميها N
نجاح التجربة P
فشل التجربة Q

* Discrete Dist. :-

- Binomial dist. :-
- ~~Bois~~ Binomial experement:-

خواص التجربة حتى تكون Binomial :

① n indep. and identical trails.

n ← من المحاولات المستقلة المتماثلة

② X: no. of successes out of n trails.

$X = \{0, 1, 2, \ldots, n\} = R_X$

← Range ال تكون محدد من الأدنى لرقم الأعلى

Ex

X no. of heads in 5 trails

$X = \{0, 1, 2, 3, 4, 5\}$

e.g: Y: no. of student who pass an exam out of

$Y = \{0, 1, \ldots, 10\}$

③ output → Success
→ fail

لها وجهتان فقط

④ $P(\text{success}) = p$ (fixed) الاحتمال كل مرة يكون ثابت

$P(\text{fail}) = q = 1 - p$

⑤ The Binomial r.v X is denoted by

$X \sim B(n, p)$

is ←

Scanned with CamScanner

Scanned with CamScanner

(6) If $X \sim B(n, p)$ then

اذا اجتمعت صفت الايدنتكل والاندبندنت بنستعمل البايناوميل دستريبيوشن

a) $EX = mean = np$

b) $Var\ X = (n * p * q)$

c) $p(X=x) = \binom{n}{x} p^x q^{n-x}$, $x = 0, 1, \dots n$

Ex) In a multiple Choice exam with 10 question, suppose that each question has 5 answer with only one true answer. If you answer all question with pure guess

a) Find the prob. that you will answer only 5 true

b) Find the prob. that you will get a full mark.

c) Find the prob. that you will answer at most 2 questions correctly.

d) Find the mean of true answered questions.

e) Find the variance of true answered questions

Scanned with CamScanner

Scanned with CamScanner

No.

Sol:

let X: no. of true answered question.

$X = \{0, 1, 2, \dots, 10\}$

$n = 10 \qquad p = \frac{1}{5} = 0.2$

$X \sim B(10, 0.2)$

دائما في Bio الاكس تبدأ من 0

a. $p(X=5) = \binom{10}{5} (0.2)^5 (0.8)^{10-5} = 0.026$

b. $p(X=10) = \binom{10}{10} (0.2)^{10} (0.8)^0 = 1.02 \times 10^{-7} = 0.0000001$

c. $p(X \leq 2) = p(X=0) + p(X=1) + p(X=2)$

$= \binom{10}{0} (0.2)^0 (0.8)^{10} + \binom{10}{1} (0.2)^1 (0.8)^9 + \binom{10}{2} (0.2)^2 (0.8)^8 =$

$=$

d. $EX = np$

$= (10)(0.2) = 2$ ← الوسط الحسابي لعدد الاجابات الصحيحة

e. $Var\ X = n.p.q \qquad = (10)(0.2)(0.8) = 1.6$

f. $std(x) = \sqrt{1.6} = 1.26$

دائما في Bio الاكس تبدأ من 0

Bio Dist الـ
فيها 3 براميتر
-N
-P
-Q

Scanned with CamScanner

Scanned with CamScanner

اذا كان في وقت بالسؤال
بكون Poisson dist

بواسون

* Poisson dist :-

- Poisson experiment :-

1) X : No. of occurences per unit of time.

e.g : X : No. of patients entering the ER per hour

$X = \{0, 1, 2, \dots\}$

↳ محدد من الاسفل وغير محدد من الأعلى .

↳ محدد في وقت معين

2) m : average of occurences per unit of time.

3) The poisson r.v. X denoted by

$X \sim p(m)$

4) If $X \sim p(m)$, then

↳ a) $p(X=x) = \frac{e^{-m} m^x}{x!}$, $x = 0, 1, 2, \dots$

↳ b) $EX = \text{Var } X = m$

Scanned with CamScanner

Scanned with CamScanner

No. 13/11

Assume that Average no. of patients entering the emergency room is 1.2 per hour.

a) Find the prob. that only 2 patients will enter per hour.

b) Find the prob. that at least 2 patients will enter per hour.

c) Find the prob. that no. of patients is between 2 and 3 per hour (included) ← بتشمل included القيمة المقفلة

d) Find $P(1.1 < X < 2.9)$

e) Find $P(X = 2.5)$

f) Find EX, $Var\,X$

g) Find the prob. that only 3 patients will enter in 4 hours.

h) Find the average no. of patients that will enter in a day.

الحل:

$X \sim P(1.2)$

a) $P(X=2) = \frac{e^{-1.2}(1.2)^2}{2!} = \dots$

b) $P(X \geq 2) = 1 - P(X < 2)$

$= 1 - [P(X=0) + P(X=1)]$

$= 1 - \left[\frac{e^{-1.2}(1.2)^0}{0!} + \frac{e^{-1.2}(1.2)^1}{1!}\right] = \dots$

اذا كان في وقت بالسؤال بكوّن Poisson dist

كم برامتر بكون في Poisson dist

1 الي هو الافرج اما الاكس لا تعتبر برامتر

Scanned with CamScanner

24
12
48
240
28.8

c) $P(2 \le X \le 3) = P(X=2) + P(X=3)$

$= \frac{e^{-1.2}(1.2)^2}{2!} + \frac{e^{-1.2}(1.2)^3}{3!} = \dots$

d) $P(1.1 < X < 2.9) = P(X=2) = \frac{e^{-1.2}(1.2)^2}{2!} = \dots$

e) $P(X=2.5) =$ Zero

f) $EX = \text{Var}\, X = 1.2$

g) let $Z = 4X$ (2.)

$\rightarrow Z \sim P((1.2)(4))$

$Z \sim P(4.8)$

$P(Z=3) = \frac{e^{-4.8}(4.8)^3}{3!} = \dots$

G & H cancellations

h) let $Y = 24X$

$\rightarrow Y \sim P((24)(1.2))$

$Y \sim P(28.8)$

$\rightarrow EY = 28.8$ patients per day

Scanned with CamScanner

Scanned with CamScanner

* Normal distribution :-
(Gaussian dist)
(Bell-shaped dist)

متماثل يمين ويسار



خواص هذه النقطة : يتساوى الوسط مع المنوال مع

1) mean = EX = median = mode = μ

median ← القيمة التي تتساوى فيها الأعلى مع الأدنى

له القيمة الأكثر تكراراً .

الرسمة بتكون متماثلة يمين ويسار

2) symmetric dist. نقطة التماثل

3) The normal r.v X depends only on μ and σ^2. So, we denoted it by:

$$X \sim N(\mu, \sigma^2)$$

μ ← الوسط الحسابي ، σ^2 ← Var

4) If $X \sim N(\mu, \sigma^2)$. then

a) $EX = \mu$, $\text{Var } X = \sigma^2$

p.d.f ←

b) $$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(x-\mu)^2} \quad , -\infty < x < \infty$$

c) $P(a \le x \le b) = P(a < x < b)$

$$= \int_a^b f(x)\,dx = \text{area under } f(x) \text{ between a and b}$$

$$= F(b) - F(a)$$

$$= \int_{-\infty}^{b} f(x)\,dx - \int_{-\infty}^{a} f(x)\,dx$$

= area to the left of b − area to the left of a

ال cdf لعدد هي المساحة على يسار العدد

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

1.

d) $P(X=c) = 0$

e) $P(X \leq b) = \int_{-\infty}^{b} f(x)\,dx = F(b)$ = area to the left of

* Standard normal distribution:-

let $X \sim N(\mu, \sigma^2)$, then the r.v.

$Z = \frac{X-\mu}{\sigma} \sim N(0,1) \equiv$ standard Normal

Properties:-

1) 

2) Symmetric about 0.

3) $EZ = 0$, $Var\ Z = 1$

4) $f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$, $-\infty < z < \infty$

5) $P(a \leq z \leq b) = \int_{a}^{b} f(z)\,dz$

= area under $f(z)$ between a and

= $T(b) - T(a)$

= table at b - table at a

= area to the left of b - area to the left of a

ال cdf بشكل عام
F الا بالستاندر
بتكون T

smile

Scanned with CamScanner

Scanned with CamScanner

6) $\int_{-\infty}^{\infty} f(z)\,dz = 1$ = total area

7) $\int_{-\infty}^{0} f(z)\,dz = 0.5 = \int_{0}^{\infty} f(z)\,dz$ لذة متماثل حول النقطة.

8) $p(z \le b) = T(b)$

9) $p(z > b) = 1 - p(z < b) = 1 - T(b)$

10) $p(z = b) = 0$

دايما 0 لنه كونتينيوس راندم فاريبل

لو بدي المساحة على يمين عدد بطلع المساحة على اليسار وبطرحها من 1

Scanned with CamScanner

Scanned with CamScanner

No.

15/11/2017

Ex) Find $p(Z<1.15) = T(1.15) = 0.87493$

Ex) $p(-1.25 < Z < 2.15)$

$= T(2.15) - T(-1.25)$

$= 0.98422 - 0.10565$

Ex) Find a s.t. $p(Z<a) = 0.9$

Ex) Find a s.t

$p(Z>a) = 0.95$



Ex) Find a s.t. $p(-a<Z<a) = 0.95$



we want a s.t $p(Z<a) = 0.975$

$a = 1.96$

Ex) Find a s.t $p(Z>a) = 0.9772$



$a = -2$

$1 - 0.9772$

$= 0.0228$

Scanned with CamScanner

Scanned with CamScanner

Ex) Suppose that the ranges of some signals are normally distributed with mean 15 Kms (μ) and std 3 Kms (σ).

15~ 19

3^2 ←

a) Find the proportion of the signals that exceeds 17 Kms.

[If a signal is chosen at random, find the prob. that its range will exceed 17 Kms

b) Find the proportion of signals with ranges less than 10 Kms

c) Find the proportion of signals with ranges between 10 and 17 Kms.

d) Find the range of signal that are best than 90% of other signals.

e) Find the range of signal that is considred with max range of the worst 5% of signals.

f) $P(X=16)$

العثور على نطاقه الإشارة التي تعتبر ذات إشارات أضعف بـ 5٪

Scanned with CamScanner

Scanned with CamScanner

No.

8

g) If two signals are chosen at random, find the prob. that their ranges will exceed 17 Kms.

h) If 5 signals are chosen at random, find the prob. that the ranges of two of them will exceed 17 Kms.

$$Z=(X-\mu)/\sigma$$

Sol:-

$$X \sim N(15, 9)$$

a) $P(X>17) = P\left(\frac{X-\mu}{\sigma} > \frac{17-\mu}{\sigma}\right)$

$= P\left(Z > \frac{17-15}{3}\right)$ $= P(Z > 0.67)$

$= 1 - P(Z \leq 0.67)$

$= 1 - T(0.67)$

$21 - 0.74857$

20.25143

b) $P(X<10) = P\left(\frac{X-\mu}{\sigma} < \frac{10-\mu}{\sigma}\right)$

$= P\left(Z < \frac{10-15}{3}\right)$

$= P(Z < -1.67)$

$= T(-1.67)$ $= 0.04746$

Scanned with CamScanner

Scanned with CamScanner

c) $P(10 < X < 17)$

$= P\left(\frac{10-\mu}{\sigma} < \frac{X-\mu}{\sigma} < \frac{17-\mu}{\sigma}\right)$

$= P(-1.67 < Z < 0.67)$

$= T(0.67) - T(-1.67)$

$= 0.74857 - 0.04746 = \ldots\ldots$

d)



but $Z = \frac{X-\mu}{\sigma}$

$1.28 = \frac{X-15}{3} \rightarrow X = (3)(1.28) + 15$

$= 18.84$ Kms

e)



$X = \sigma Z + \mu$

$= (3)(-1.64) + 15 = 10.08$ Kms

f) $P(X=16) = 0$

Scanned with CamScanner

Scanned with CamScanner

(15)

g) $p(X_1 > 17) = 0.25143$ (من الفرع a)

$p(X_2 > 17) = 0.25143$

we want

$\rightarrow p(X_1 > 17 \cap X_2 > 17)$

$= p(X_1 > 17) \cdot p(X_2 > 17)$

$= (0.25143)^2$

هون مختار 2 سقنل وبده 2 يكونو صح

h) $p(X_1 > 17) = 0.25143$

$p(X_2 > 17) = 0.25143$

$\vdots$

$p(X_5 > 17) = 0.25143$

هون مختار 5 سقنل وبده 2 يكونو صح عشان هيك حله بختلف عن الي فوق لو طلب 5 صح بنحله زي الي فوق

All signals are indep. and identical $n=5$

$\rightarrow Y \sim B(5, 0.25143)$

$\leftarrow P(y=2) = \binom{5}{2} (0.25143)^2 (1-0.25143)^{5-2}$ binomial

$= \dots$

Scanned with CamScanner

Scanned with CamScanner

STANDARD NORMAL DISTRIBUTION: Table Values Represent AREA to the LEFT of the Z score.

Z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.9	.00005	.00005	.00004	.00004	.00004	.00004	.00004	.00004	.00003	.00003
-3.8	.00007	.00007	.00007	.00006	.00006	.00006	.00006	.00005	.00005	.00005
-3.7	.00011	.00010	.00010	.00010	.00009	.00009	.00008	.00008	.00008	.00008
-3.6	.00016	.00015	.00015	.00014	.00014	.00013	.00013	.00012	.00012	.00011
-3.5	.00023	.00022	.00022	.00021	.00020	.00019	.00019	.00018	.00017	.00017
-3.4	.00034	.00032	.00031	.00030	.00029	.00028	.00027	.00026	.00025	.00024
-3.3	.00048	.00047	.00045	.00043	.00042	.00040	.00039	.00038	.00036	.00035
-3.2	.00069	.00066	.00064	.00062	.00060	.00058	.00056	.00054	.00052	.00050
-3.1	.00097	.00094	.00090	.00087	.00084	.00082	.00079	.00076	.00074	.00071
-3.0	.00135	.00131	.00126	.00122	.00118	.00114	.00111	.00107	.00104	.00100
-2.9	.00187	.00181	.00175	.00169	.00164	.00159	.00154	.00149	.00144	.00139
-2.8	.00256	.00248	.00240	.00233	.00226	.00219	.00212	.00205	.00199	.00193
-2.7	.00347	.00336	.00326	.00317	.00307	.00298	.00289	.00280	.00272	.00264
-2.6	.00466	.00453	.00440	.00427	.00415	.00402	.00391	.00379	.00368	.00357
-2.5	.00621	.00604	.00587	.00570	.00554	.00539	.00523	.00508	.00494	.00480
-2.4	.00820	.00798	.00776	.00755	.00734	.00714	.00695	.00676	.00657	.00639
-2.3	.01072	.01044	.01017	.00990	.00964	.00939	.00914	.00889	.00866	.00842
-2.2	.01390	.01355	.01321	.01287	.01255	.01222	.01191	.01160	.01130	.01101
-2.1	.01786	.01743	.01700	.01659	.01618	.01578	.01539	.01500	.01463	.01426
-2.0	.02275	.02222	.02169	.02118	.02068	.02018	.01970	.01923	.01876	.01831
-1.9	.02872	.02807	.02743	.02680	.02619	.02559	.02500	.02442	.02385	.02330
-1.8	.03593	.03515	.03438	.03362	.03288	.03216	.03144	.03074	.03005	.02938
-1.7	.04457	.04363	.04272	.04182	.04093	.04006	.03920	.03836	.03754	.03673
-1.6	.05480	.05370	.05262	.05155	.05050	.04947	.04846	.04746	.04648	.04551
-1.5	.06681	.06552	.06426	.06301	.06178	.06057	.05938	.05821	.05705	.05592
-1.4	.08076	.07927	.07780	.07636	.07493	.07353	.07215	.07078	.06944	.06811
-1.3	.09680	.09510	.09342	.09176	.09012	.08851	.08691	.08534	.08379	.08226
-1.2	.11507	.11314	.11123	.10935	.10749	.10565	.10383	.10204	.10027	.09853
-1.1	.13567	.13350	.13136	.12924	.12714	.12507	.12302	.12100	.11900	.11702
-1.0	.15866	.15625	.15386	.15151	.14917	.14686	.14457	.14231	.14007	.13786
-0.9	.18406	.18141	.17879	.17619	.17361	.17106	.16853	.16602	.16354	.16109
-0.8	.21186	.20897	.20611	.20327	.20045	.19766	.19489	.19215	.18943	.18673
-0.7	.24196	.23885	.23576	.23270	.22965	.22663	.22363	.22065	.21770	.21476
-0.6	.27425	.27093	.26763	.26435	.26109	.25785	.25463	.25143	.24825	.24510
-0.5	.30854	.30503	.30153	.29806	.29460	.29116	.28774	.28434	.28096	.27760
-0.4	.34458	.34090	.33724	.33360	.32997	.32636	.32276	.31918	.31561	.31207
-0.3	.38209	.37828	.37448	.37070	.36693	.36317	.35942	.35569	.35197	.34827
-0.2	.42074	.41683	.41294	.40905	.40517	.40129	.39743	.39358	.38974	.38591
-0.1	.46017	.45620	.45224	.44828	.44433	.44038	.43644	.43251	.42858	.42465
-0.0	.50000	.49601	.49202	.48803	.48405	.48006	.47608	.47210	.46812	.46414

STANDARD NORMAL DISTRIBUTION: Table Values Represent AREA to the LEFT of the Z score.

Z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.50000	.50399	.50798	.51197	.51595	.51994	.52392	.52790	.53188	.53586
0.1	.53983	.54380	.54776	.55172	.55567	.55962	.56356	.56749	.57142	.57535
0.2	.57926	.58317	.58706	.59095	.59483	.59871	.60257	.60642	.61026	.61409

Scanned with CamScanner

0.3	.61791	.62172	.62552	.62930	.63307	.63683	.64058	.64431	.64803	.65173
0.4	.65542	.65910	.66276	.66640	.67003	.67364	.67724	.68082	.68439	.68793
0.5	.69146	.69497	.69847	.70194	.70540	.70884	.71226	.71566	.71904	.72240
0.6	.72575	.72907	.73237	.73565	.73891	.74215	.74537	.74857	.75175	.75490
0.7	.75804	.76115	.76424	.76730	.77035	.77337	.77637	.77935	.78230	.78524
0.8	.78814	.79103	.79389	.79673	.79955	.80234	.80511	.80785	.81057	.81327
0.9	.81594	.81859	.82121	.82381	.82639	.82894	.83147	.83398	.83646	.83891
1.0	.84134	.84375	.84614	.84849	.85083	.85314	.85543	.85769	.85993	.86214
1.1	.86433	.86650	.86864	.87076	.87286	.87493	.87698	.87900	.88100	.88298
1.2	.88493	.88686	.88877	.89065	.89251	.89435	.89617	.89796	.89973	.90147
1.3	.90320	.90490	.90658	.90824	.90988	.91149	.91309	.91466	.91621	.91774
1.4	.91924	.92073	.92220	.92364	.92507	.92647	.92785	.92922	.93056	.93189
1.5	.93319	.93448	.93574	.93699	.93822	.93943	.94062	.94179	.94295	.94408
1.6	.94520	.94630	.94738	.94845	.94950	.95053	.95154	.95254	.95352	.95449
1.7	.95543	.95637	.95728	.95818	.95907	.95994	.96080	.96164	.96246	.96327
1.8	.96407	.96485	.96562	.96638	.96712	.96784	.96856	.96926	.96995	.97062
1.9	.97128	.97193	.97257	.97320	.97381	.97441	.97500	.97558	.97615	.97670
2.0	.97725	.97778	.97831	.97882	.97932	.97982	.98030	.98077	.98124	.98169
2.1	.98214	.98257	.98300	.98341	.98382	.98422	.98461	.98500	.98537	.98574
2.2	.98610	.98645	.98679	.98713	.98745	.98778	.98809	.98840	.98870	.98899
2.3	.98928	.98956	.98983	.99010	.99036	.99061	.99086	.99111	.99134	.99158
2.4	.99180	.99202	.99224	.99245	.99266	.99286	.99305	.99324	.99343	.99361
2.5	.99379	.99396	.99413	.99430	.99446	.99461	.99477	.99492	.99506	.99520
2.6	.99534	.99547	.99560	.99573	.99585	.99598	.99609	.99621	.99632	.99643
2.7	.99653	.99664	.99674	.99683	.99693	.99702	.99711	.99720	.99728	.99736
2.8	.99744	.99752	.99760	.99767	.99774	.99781	.99788	.99795	.99801	.99807
2.9	.99813	.99819	.99825	.99831	.99836	.99841	.99846	.99851	.99856	.99861
3.0	.99865	.99869	.99874	.99878	.99882	.99886	.99889	.99893	.99896	.99900
3.1	.99903	.99906	.99910	.99913	.99916	.99918	.99921	.99924	.99926	.99929
3.2	.99931	.99934	.99936	.99938	.99940	.99942	.99944	.99946	.99948	.99950
3.3	.99952	.99953	.99955	.99957	.99958	.99960	.99961	.99962	.99964	.99965
3.4	.99966	.99968	.99969	.99970	.99971	.99972	.99973	.99974	.99975	.99976
3.5	.99977	.99978	.99978	.99979	.99980	.99981	.99981	.99982	.99983	.99983
3.6	.99984	.99985	.99985	.99986	.99986	.99987	.99987	.99988	.99988	.99989
3.7	.99989	.99990	.99990	.99990	.99991	.99991	.99992	.99992	.99992	.99992
3.8	.99993	.99993	.99993	.99994	.99994	.99994	.99994	.99995	.99995	.99995
3.9	.99995	.99995	.99996	.99996	.99996	.99996	.99996	.99996	.99997	.99997

Scanned with CamScanner

Transformation of random Variable:-

If $f_X(x)$ is the p.d.f of a r.v. X and $\boxed{Y = T(X)}$ is a function of the r.v. X, we want to find the distribution of a r.v. Y

$$F_Y(y_0) = P\{Y \leq y_0\}$$

ال3 اسطر اثبات مش معنا

$$= P\{T(X) \leq y_0\}$$

$$= P\{X \leq T^{-1}(y_0)\} = F_X(x_0)$$

Hence,

$$f_Y(y) = f_X(T^{-1}(y)) \left| \frac{d\,T^{-1}(y)}{dy} \right| \rightarrow \text{Jacobian}$$

ex) Id $Y = aX + b$, then

$$X = \frac{Y-b}{a} \longrightarrow \frac{dx}{dy} = \frac{1}{a}$$

$$f_Y(y) = f_X\left(\frac{Y-b}{a}\right) \left|\frac{1}{a}\right|$$

ex) If $Y = aX^2 + b$, then

$$X = \pm\sqrt{\frac{Y-b}{a}} \longrightarrow \frac{dx}{dy} = \frac{\frac{1}{a}}{\pm 2\sqrt{\frac{Y-b}{a}}}$$

$$f_Y(y) = F_X\left(\pm\sqrt{\frac{Y-b}{a}}\right) \left| \frac{\frac{1}{a}}{\pm 2\sqrt{\frac{Y-b}{a}}} \right|$$

smile for life

Scanned with CamScanner

Scanned with CamScanner

Ex] Consider the dist. with p.d.f

$$f(x) = \frac{\theta^2}{\theta+1}(1+x)e^{-\theta x}, \quad x>0$$

Find the density of Y s.t.

$$Y = e^X - 1, \quad y>0.$$

بنخلي x موضع قانون

$$y+1 = e^x \longrightarrow x = \ln(y+1)$$

$$\frac{dx}{dy} = \frac{1}{y+1}$$

$$f_Y(y) = f_X(\ln(y+1)) \left|\frac{1}{y+1}\right|$$

$$= \frac{\theta^2}{\theta+1}\left(1+\ln(y+1)\right) e^{-\theta(\ln(y+1))} \left|\frac{1}{y+1}\right|$$

$$= \frac{\theta^2}{\theta+1}\left(1+\ln(y+1)\right)(y+1)^{-\theta}\frac{1}{y+1}$$

$$= \frac{\theta^2}{\theta+1}\left(1+\ln(y+1)\right)\frac{1}{(y+1)^{\theta+1}}$$

Scanned with CamScanner

Scanned with CamScanner

$$y = \frac{1}{x} \longrightarrow x = \frac{1}{y} \longrightarrow \frac{dx}{dy} = \frac{-1}{y^2}$$

$$f_Y(y) = f_X\left(\frac{1}{y}\right)\left|\frac{-1}{y^2}\right|$$

(*) Distribution of sum of two random variables:-

Let X, Y be two indep. r.v's and $\boxed{W = X+Y}$

$$F_W(w) = P\{W \le w\} = P\{X+Y \le w\}$$

هون بدنا نوجد pdf للـ w



$$= \int_{-\infty}^{\infty}\int_{-\infty}^{w-y} f(x,y)\,dx\,dy$$

$$f_W(w) = \int_{-\infty}^{\infty} f_Y(y) \int_{-\infty}^{w-y} f_X(x)\,dx\,dy$$

by diff.

$$f_W(w) = \int_{-\infty}^{\infty} f_Y(y)\, f_X(w-y)\,dy .$$

Scanned with CamScanner

Scanned with CamScanner

No.

Ex] Find the density of $W = X + Y$ where

$$f_X(x) = \frac{1}{a}\left[u(x) - u(x-a)\right]$$

$$f_Y(y) = \frac{1}{b}\left[u(y) - u(y-b)\right]$$

$x = w - y$ نعوض بدل x

يو اف واي ناقص عدد هو واحد على يمين العدد وصفر على يساره

$$* f_W(w) = \int_{-\infty}^{\infty} \frac{1}{b}\left[u(y) - u(y-b)\right] \cdot \frac{1}{a}\left[u(w-y) - u(w-y-a)\right]$$

$$= \frac{1}{ab}\int_{0}^{\infty}\left[1 - u(y-b)\right]\left[u(w-y) - u(w-y-a)\right] dy$$

ضرب ← القوسين

يو اف عدد ناقص واي صفر على يمين العدد وواحد على يسارة

تكامل ← طرح

$$= \frac{1}{ab}\int_{0}^{\infty} \underbrace{\left(u(w-y)\right)}_{①} - \underbrace{u(w-y-a)}_{②} - \underbrace{u(y-b)\,u(w-y)}_{③} + \underbrace{u(y-b)\,u(w-y-a)}_{④}\, dy$$

الـ A < B والـ2 موجبات

$$① = \frac{1}{ab}\int_{0}^{\infty} u(w-y)\,dy = \frac{1}{ab}\int_{0}^{w} 1_o\, dy = \frac{w}{ab}$$

$$② = \frac{-1}{ab}\int_{0}^{\infty} u(w-y-a)\,dy = \frac{-1}{ab}\int_{0}^{w-a} 1 \cdot dy = -\left(\frac{w-a}{ab}\right)$$

IF W>=A

If w-a >= 0 , and 0 o.w

$$③ = \frac{-1}{ab}\int_{0}^{\infty} u(y-b)\,u(w-y)\,dy = \frac{1}{ab}\int_{b}^{w} 1 \cdot dy = -\left(\frac{w-b}{ab}\right)$$

$w > b$:ف

Scanned with CamScanner

Scanned with CamScanner

④ $= \frac{1}{ab} \int_0^{\infty} u(y-b)\, u(w-a-y)\, dy$

$= \frac{1}{ab} \int_b^{w-a} 1\, dy \quad = \frac{w-a-b}{ab}$

If w-a >= b , and 0 o.w

- $f_W(w) = \begin{cases} ① & , \ w < a \\ ①+② & , \ a \le w < b \\ ①+②+③ & , \ b \le w < a+b \\ ①+②+③+⓪ & , \ w \ge a+b \end{cases}$

smile for life

Scanned with CamScanner

Scanned with CamScanner

$$\textcircled{1} = \frac{w}{ab} \qquad \textcircled{2} = 0 \qquad \textcircled{3} = 0 \qquad \textcircled{4} = 0$$

$$\textcircled{1} = \frac{w}{ab} \qquad \textcircled{2} = -\frac{(w-a)}{ab} \qquad \textcircled{3} = 0 \qquad \textcircled{4} = 0$$

$$\textcircled{1} = \frac{w}{ab} \qquad \textcircled{2} = -\left(\frac{w-a}{ab}\right) \qquad \textcircled{3} = -\left(\frac{w\,a-b}{ab}\right) \qquad \textcircled{4} = 0$$

$$\textcircled{1} = \frac{w}{ab} \qquad \textcircled{2} = \frac{-(w-a)}{ab} \qquad \textcircled{3} = -\left(\frac{w\,a-b}{ab}\right) \qquad \textcircled{4} = \frac{w-a-b}{ab}$$

Scanned with CamScanner

Scanned with CamScanner

$$f_W(w) = \int_{-\infty}^{\infty} f_y(y)\, f_x(w-y)\, dy$$

Ex) $f_x(x) = \frac{1}{a}[u(x) - u(x-a)]$

$$f_y(y) = b\, u(y)\, e^{-by}$$

Find the density of $W = X+Y$
s.t. X, Y are indep.

$$f_W(w) = \int_{-\infty}^{\infty} f_y(y)\, f_x(w-y)\, dy$$

$$= \int_{-\infty}^{\infty} b\, u(y)\, e^{-by} \frac{1}{a}[u(w-y) - u(w-y-a)]\, dy$$

$$= \frac{b}{a}\int_{0}^{\infty} e^{-by}[u(w-y) - u(w-a-y)]\, dy$$

smile for life

Scanned with CamScanner

Scanned with CamScanner

No.

$$= \frac{b}{a} \int_0^{\infty} \left(\underbrace{e^{-by}\, u(w-y)}_{①} - \underbrace{e^{-by}\, u(w-a-y)}_{②} \right) dy$$

$$① = \frac{b}{a} \int_0^{\infty} e^{-by}\, u(w-y)\, dy$$

$$= \frac{b}{a} \int_0^{w} e^{-by} \cdot 1\, dy = \frac{b}{a} \cdot \frac{-1}{b} e^{-by} \Big]_0^{w}$$

$$= \frac{-1}{a} \left(e^{-bw} - 1 \right)$$

$$② = \frac{-b}{a} \int_0^{\infty} e^{-by}\, u(w-a-y)\, dy$$

$$= \frac{-b}{a} \int_0^{w-a} e^{-by}\, dy = \frac{-b}{a} \cdot \frac{-1}{b} e^{-by} \Big]_0^{w-a}$$

$$= \frac{1}{a} \left(e^{-b(w-a)} - 1 \right)$$

$$f_W(w) = \begin{cases} ① & , w < a \\ ① + ② & , a \leq w \end{cases}$$

Scanned with CamScanner

Scanned with CamScanner

No.

⊗ Expected value of functions of r.v's :-

- Recall; $$EX = \int_{-\infty}^{\infty} x f_X(x)\, dx$$

$$EY = \int_{-\infty}^{\infty} y f_Y(y)\, dy$$

$$EXY = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x y\, f(x,y)\, dx\, dy$$

$$Eg(x) = \int_{-\infty}^{\infty} g(x) f_X(x)\, dx$$

$$Eg(y) = \int_{-\infty}^{\infty} g(y) f_Y(y)\, dy$$

$$Eg(x,y) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} g(x,y) f(x,y)\, dx\, dy$$

* Special Case :-

⇝ The joint moment about the origin is given by:

$$m_{nk} = E(X^n Y^k) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x^n y^k f(x,y)\, dx\, dy$$

$$m_{n0} = E(X^n) = \int_{-\infty}^{\infty} x^n f_X(x)\, dx$$

is called the n^{th} moment of X

$$m_{0k} = E(Y^k) = \int_{-\infty}^{\infty} y^k f_Y(y)\, dy$$

is called the k^{th} moment of Y

smile for life

Scanned with CamScanner

Scanned with CamScanner

⊗ Expected value of functions of r.v's :-

- Recall; $EX = \int_{-\infty}^{\infty} x f_X(x)\, dx$

$$EY = \int_{-\infty}^{\infty} y f_Y(y)\, dy$$

$$EXY = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} xy\, f(x,y)\, dx\, dy$$

$$Eg(x) = \int_{-\infty}^{\infty} g(x) f_X(x)\, dx$$

$$Eg(y) = \int_{-\infty}^{\infty} g(y) f_Y(y)\, dy$$

$$Eg(x,y) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} g(x,y) f(x,y)\, dx\, dy$$

* Special case :-

↝ The joint moment about the origin is given by:

$$m_{nk} = E(X^n Y^k) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x^n y^k f(x,y)\, dx\, dy$$

$$m_{n0} = E(X^n) = \int_{-\infty}^{\infty} x^n f_X(x)\, dx$$

is called the n^{th} moment of X

$$m_{0k} = E(Y^k) = \int_{-\infty}^{\infty} y^k f_Y(y)\, dy$$

is called the K^{th} moment of Y

smile for life
Scanned with CamScanner
Scanned with CamScanner

The 1^{st} moment of X is EX_1 = mean = μ,

The 2^{nd} moment of X is EX^2

$\text{Var } X = EX^2 - (EX)^2$

$= 2^{nd}$ moment $- (1^{st}$ moment$)^2$

* $n+K$ is called the order of the moment.

* $m_{11} = E(XY)$ is called the correlation.

$$m_{11} = E(XY) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} xy\, f(x,y)\, dx\, dy$$

= Correlation of X_1Y

$= R_{XY}$.

Scanned with CamScanner

Scanned with CamScanner

(*) If X, y are indep., then

$$EXy = (EX)(EY)$$

$$m_{11} = (m_{10})(m_{01}).$$

* If $R_{xy} = 0 = Exy$,
then X, y are orthogonal.

Ex) let X be a r.v. with $EX = 3$, $6x^2 = 2$
and $Y = -6X + 22$.

a) Find the correlation of X, y
b) Are X, y correlated.

sol:

$$m_{11} = EXY$$

$$= E(X(-6X + 22))$$

$$= E(-6X^2 + 22X)$$

$$= -6EX^2 + 22EX$$

But $6x^2 = EX^2 - (EX)^2$

$$2 = EX^2 - (3)^2 \qquad EX^2 = 11$$

So, $EXy = -6EX^2 + 22EX$

$= (-6)(111 + (22)(3)) = 0$ مهم

So, X, y are orthogonal

Scanned with CamScanner

Scanned with CamScanner

b) $EXY \stackrel{?}{=} (EX)(EY)$

Now, $EY = E(-6X + 22)$

$= -6EX + 22$

$= (-6)(3) + 22 = 4$

So, $EXY \neq (EX)(EY)$

$0 \neq (3)(4)$

So, X, Y are not indep.

X, Y are depent.

X, Y are correlated.

* If $EXY = (EX)(EY)$, then

X, Y are indep.

X, Y are uncorrelated

* If $Y = aX + b$, $a \neq 0$, then

X, Y are correlated.

Scanned with CamScanner

Scanned with CamScanner

⊛ Joint Central Moment :-

It is defined as:-

$$\mu_{nk} = E\left((x-\bar{x})^n (y-\bar{y})^k\right)$$

$$= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} (x-\bar{x})^n (y-\bar{y})^k f(x,y)\, dx\, dy$$

Notice that:

$$\mu_{20} = E(x-\bar{x})^2 = \int_{-\infty}^{\infty} (x-\bar{x})^2 f_x(x)\, dx = \sigma_x^2$$

$$\mu_{02} = E(y-\bar{y})^2 = \sigma_x^2$$

and

$$\mu_{11} = E\left((x-\bar{x})(y-\bar{y})\right)$$

$$= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} (x-\bar{x})(y-\bar{y}) f(x,y)\, dx\, dy$$

$$= C_{xy} = Cov(x,y)$$

= is the covariance of x, y

$$= Exy - (Ex)(Ey)$$

$$= Rxy - (m_{10})(m_{01})$$

Scanned with CamScanner

Scanned with CamScanner

~ If X,Y are orthogonal, then

$$C_{XY} = Cov(X,Y) = -(EX)(EY)$$

~ If X,Y are indep. (uncorrelation) then

$$C_{XY} = Cov(X,Y) = 0$$

~ Correlation Coefficient

$$= Corr(X,Y) = \frac{Cov(X,Y)}{\sigma_X \sigma_Y} = \frac{C_{XY}}{\sigma_X \sigma_Y}$$

$$= 0$$

* The Uniform distribution:-

The Uniform r.v. is defined as

$X \sim U(a,b)$ with p.d.f.

$$f(x) = \begin{cases} \frac{1}{b-a} & , a \le x \le b \\ 0 & , o.w \end{cases}$$

and c.d.f.

$$F(x) = \begin{cases} 0 & , x < a \\ \frac{x-a}{b-a} & , a \le x < b \\ 1 & , x \ge b \end{cases}$$

Scanned with CamScanner

Scanned with CamScanner

No.





Scanned with CamScanner

Random process

Def A random process (RP) [or stochastic process] is an infinite indexed family $X(t,s) = \{x(t,s)\}$

"$s \in \mathcal{S}$ is a possible outcome of an experiment"

note $x(t,s_0)$ (s_0 fixed) is called sample function or ensemble function or realization of a process.

note $x(t_0,s)$ (t_0 fixed) is a random variable on $\mathcal{S}$

note For frozen time $t = t_1$

$$X_1 := X(t_1,s) = X(t_1)$$

describes statistically the properties of the R.V at time t_1

Scanned with CamScanner

Scanned with CamScanner

$E(X_1)$ = mean value of the process

is called ensemble average



note

In general

$X_i = X(t_i, s) = X(t_i)$

$i = 1, 2, \ldots$

Scanned with CamScanner

Scanned with CamScanner

classification of process

① X is cont & t is cont

X(t) is called Continuous r.p



② X is discrete & t is cont. (discrete random process)



The sample function has only two values

③ X is cont, & t is discrete

(Cont. random sequence)

(discrete time "DT" R.P)



Scanned with CamScanner

Scanned with CamScanner

A DT random process is a set of random variables $\{X(t_n)\}$, for sample times $t_n = nT_s$, $n = 0, \pm 1, \pm 2, \ldots$

T_s := sampling interval

$1/T_s$ = sampling rate "sample per second"

④ X & t discrete

(discrete r.p)



Scanned with CamScanner

Scanned with CamScanner

Def In a deterministic R.P / any future values of any sample fn can be predicted from the past ones :

ex : $X(t) = A \cos(\omega t + \theta)$

where A & ω are fixed

θ r.v

θ is determined,

note : Once θ is ... $X(t)$ as a fn of time is entirely specified.

Scanned with CamScanner

Scanned with CamScanner

No.

Def: The random process is called stationary if all of its statistical properties do not change with time.

i.e. At time t_1, the c.d.f. of the random variable $X_1 = x(t_1)$

is

$$F_X(x_1, t_1) = P\{X(t_1) \leq x_1\}, \quad \forall x_1$$

And, for two random variables $X_1 = x(t_1)$, $X_2 = x(t_2)$, the joint c.d.f.

$$F(x_1, x_2, t_1, t_2) = P\{X(t_1) \leq x_1, X(t_2) \leq x_2\}, \quad \forall x_1, x_2$$



بكون wide
لما يتحققو ال2

Scanned with CamScanner

Scanned with CamScanner

Remember that:-

- x, y indep. if $f(x,y) = f_x(x) \cdot f_y(y)$

→ $EXY = EX\,EY$

$Cov(x,y) = C_{xy} = 0$

$Corr(x,y) = 0$

⑤ Independence:-

Two random processes $x(t)$, $y(t)$ are statistically indep. if

$$f(x_1, x_2 \dots x_n, y_1, y_2, \dots, y_m, t_1, t_2, \dots, t_n, t_1', t_2', \dots, t_m')$$

$$= f_X(x_1, x_2 \dots x_n, t_1, t_2, \dots t_n) \cdot f_Y(y_1, y_2, \dots y_m, t_1', t_2'$$

$$(f(\tilde{x}, \tilde{y}, \tilde{t}, \tilde{t}') = f_x(\tilde{x}, \tilde{t}) \cdot f_y(\tilde{y}, \tilde{t}')$$

$$\tilde{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \tilde{y} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix}, \tilde{t} = \begin{pmatrix} t_1 \\ \vdots \\ t_n \end{pmatrix}, \tilde{t}' = \begin{pmatrix} \tilde{t_1'} \\ \vdots \\ \tilde{t_m'} \end{pmatrix}$$

⑥ First order stationary:

A random process is called 1st order-stationary if its first-order density function (p.d.f) do not change with time shift.

i.e: $f_X(x, t_1) = f_X(x, t_1 + \Delta)$

Scanned with CamScanner

Scanned with CamScanner

First order stationary:

A random process is called 1st order-stationary if its first-order density (p.d.f) function do not change with time shift.

i.e. $f_X(x, t_1) = f_X(x, t_1 + \Delta)$

Scanned with CamScanner

Scanned with CamScanner

This mean that $f_X(x, t_1)$ is indep. of t_1
and so $E(X(t)) = \overline{X} =$ constant

↳ ليس اقتران بدلالة t
(فيه ثابت)

⊛ Second order stationary :-

A process is called 2nd order stationary if its 2nd order density satisfies

$$f_X(x_1, x_2, t_1, t_2) = f_X(x_1, x_2, t_1 + \Delta, t_2 + \Delta)$$ ⊛

Remarks:

(1) The correlation $E(x_1 x_2) = E(X(t_1)\, X(t_2))$
can be denoted by $R_{XX}(t_1, t_2)$
and it is called the auto-correlation function.
of the random process $X(t)$.

Hense $R_{XX}(t_1, t_2) = E(X(t_1)\, X(t_2))$
let $t_1 = t$, $t_2 = t_1 + \tau$ $(\tau = t_2 - t_1)$

In case of 2nd order stationary, the random process satisfies ⊛ Consequently

$$R_{XX}(t, t+\tau) = E(X(t_1)\, X(t_2 + \tau))$$
$$= E(X(t)\, X(t+\tau))$$
$$= R_{XX}(\tau)$$ Auto-correlation

smile for life

Scanned with CamScanner

Scanned with CamScanner

(*) $x(t)$ is called 2nd order stationary if the auto-correlation of $x(t)$

$$R_{xx}(t, t+\tau) = R_{xx}(\tau)$$

The random process is called wide-sense stationary if $E(x(t)) = \bar{x} = \text{Constant}$

$$R_{xx}(t, t+\tau) = R_{xx}(\tau)$$

(*) Ex] Show that the process $x(t) = A\cos(\omega_0 t + \Theta)$ is wide-sense stationary if A, ω_0 are constants and θ is Uniformly distributed r.v. on $(0, 2\pi)$

Now,

$$E(x(t)) = \int_{-\infty}^{\infty} x(t)\, f(\theta)\, d\theta$$

$$= \int_0^{2\pi} A\cos(\omega_0 t + \Theta) \cdot \frac{1}{2\pi}\, d\theta$$

$$= \frac{A}{2\pi} \int_0^{2\pi} \cos(\omega_0 t + \Theta)\, d\theta$$

$= 0 \equiv$ Constant.

$\rightarrow$ $x(t)$ is 1st order stationary

Remember that

If $X \sim U(a, b)$

$$f(x) = \begin{cases} \frac{1}{b-a}, & a \le x \le b \\ 0, & \text{o.w} \end{cases}$$

$$F(x) = \begin{cases} \frac{x-a}{b-a}, & a \le x < b \\ 0, & \text{o.w} \end{cases}$$

$\Theta \sim U(0, 2\pi)$

$$f(\theta) = \begin{cases} \frac{1}{2\pi - 0}, & 0 \le \theta \le 2\pi \\ 0, & \text{o.w.} \end{cases}$$

Scanned with CamScanner

Scanned with CamScanner

Now, the auto-correlation with

$t_1 = t$, $t_2 = t + \tau$ is

$$R_{xx}(t, t+\tau) = E(x(t) * x(t+\tau))$$

$$= E(A\cos(\omega_0 t + \theta) \cdot A\cos(\omega_0 t + \omega_0 \tau + \theta))$$

$$= A^2 E(\cos(\omega_0 t + \theta) \cdot \cos(\omega_0 t + \omega_0 \tau + \theta))$$

$$= \frac{A^2}{2} E(\cos(2\omega_0 t + \omega_0 \tau + 2\theta) + \cos(\omega_0 \tau))$$

$$= \frac{A^2}{2} E(\cos(2\omega_0 t + \omega_0 \tau + 2\theta)) + \frac{A^2}{2} E(\cos(\omega_0 \tau)) \rightarrow \text{free of } \theta$$

$$\frac{A^2}{2} \int_0^{2\pi} \cos(2\omega_0 t + \omega_0 \tau + 2\theta) \cdot \frac{1}{2\pi} \, d\theta + \frac{A^2}{2} \cos(\omega_0 \tau)$$

$$= \frac{A^2}{2} \cos(\omega_0 \tau) = R_{xx}(\tau)$$

So, $x(t)$ is 2nd order stationary.

$\rightarrow$ $x(t)$ is wide-sense stationary.

Remember that:-

- $\cos(a+b) = \cos a \cos b - \sin a \sin b$
- $\cos(a-b) = \cos a \cos b + \sin a \sin b$
- $\frac{\cos(a+b) + \cos(a-b)}{2} = \cos a \cos b$
- $\frac{\cos(a-b) - \cos(a+b)}{2} = \sin a \sin b$
- $\cos(-x) = \cos x$

  $\sin(-x) = -\sin x$

Scanned with CamScanner

Scanned with CamScanner

NO.

* If we have two random processes $x(t)$, $y(t)$. then,
they are jointly wide-sense stationary if each one of them is wide-sense stationary and their cross-correlation is a function of $\tau = t_2 - t_1$,

$$\text{i.e. } R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau))$$
$$= R_{xy}(\tau).$$

* Correlation function:-

* Auto-correlation function:-

The auto-correlation function of $x(t)$ is denoted by

$$R_{xx}(t, t+\tau) = E(x(t)\, x(t+\tau))$$

If $x(t)$ is wide-sense stationary, then

$$R_{xx}(t, t+\tau) = R_{xx}(\tau).$$

(*) For any random process, the auto-correlation function satisfies the following:-

1) $|R_{xx}(\tau)| \leq R_{xx}(0)$.

2) $R_{xx}(-\tau) = R_{xx}(\tau)$

Scanned with CamScanner

Scanned with CamScanner

3) $R_{xx}(\infty) = E(X^2(t))$

مهم 4 + 3

4) $\lim_{\tau \to \infty} R_{xx}(\tau) = \bar{X}^2$

5) If $x(t)$ is periodic, then $R_{xx}(\tau)$ is periodic with the same period.

if $E(x(t)) = \bar{X}$ & $x(t) = \bar{X}$ + متغير عشوائي

Def: A process $x(t)$ with constant mean value $\bar{X}$ is called mean-ergodic

i.e. $E(x(t)) = \bar{X} = A(x(t)) = \bar{x}(t)$

$$A(\star) = \lim_{T \to \infty} \frac{1}{2T} \int_{-T}^{T} \star \, dt$$

Ex) Given the auto-correlation function for a stationary ergodic process with (no) periodic component is

$$R_{xx}(\tau) = 25 + \frac{4}{1+6\tau^2}$$

- Find the mean and variance of the process.

Sol:

By (4);

$$\lim_{\tau \to \infty} R_{xx}(\tau) = \lim_{\tau \to \infty} \left(25 + \frac{4}{1+6\tau^2}\right) = 25 = \bar{X}^2 \longrightarrow \bar{X} = \pm 5$$

By (3); $R_{xx}(0) = 25 + \frac{4}{1+6(0)^2} = 29 = EX^2_{(t)}$

$\sigma^2_{x(t)} = EX^2_{(t)} - (EX_{(t)})^2$

$= 29 - (\pm 5)^2 = 4$

smile

Scanned with CamScanner

Scanned with CamScanner

* Cross-correlation function:-

The cross-correlation of $x(t)$, $y(t)$ is given by

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau))$$

* If $x(t)$, $y(t)$ are jointly wide-sense stationary, then

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau)) = R_{xy}(\tau)$$

* If $R_{xy}(\tau) = 0$, then $x(t)$, $y(t)$ are orthogonal processes.

* If $x(t)$, $y(t)$ are indep., then

$$R_{xy}(t, t+\tau) = E(x(t))\, E(y(t+\tau))$$

If $x(t)$, $y(t)$ are indep and at least wide-sense stationary, then

$$R_{xy}(\tau) = \bar{x}\,\bar{y} \text{ which is constant.}$$

⊛ properties:

1) $R_{xy}(-\tau) = R_{xy}(\tau)$.

2) $|R_{xy}(\tau)| \leq \sqrt{R_{xx}(0)\, R_{yy}(0)}$

مهم

3) $|R_{xy}(\tau)| \leq \frac{1}{2}[R_{xx}(0) + R_{yy}(0)]$

smile for life
smile for life
Scanned with CamScanner
Scanned with CamScanner

Ex Assume the two random process

$$x(t) = A\cos(\omega_0 t) + B\sin(\omega_0 t)$$
$$y(t) = B\cos(\omega_0 t) - A\sin(\omega_0 t),$$

where A, B are r.v's, ω_0 constant, A, B are uncorrelated, Zero-mean with same variance.

Find the cross-correlation $R_{xy}(t, t+\tau)$ and show that $x(t), y(t)$ are jointly wide-sense stationary.

$$E(A) = E(B) = 0$$
$$E(AB) \overset{\text{indep}}{=} E(A)E(B) = 0$$

$$\sigma_A^2 = \sigma_B^2 = \sigma^2$$

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau))$$
$$= E(\{A\cos(\omega_0 t) + B\sin(\omega_0 t)\}.$$
$$\{B\cos(\omega_0 t + \omega_0\tau) - A\sin(\omega_0 t + \omega_0\tau)\})$$

$$= E(AB\cos(\omega_0 t)\cos(\omega_0 t + \omega_0\tau) - A^2\cos(\omega_0 t)\sin(\omega_0 t + \omega_0\tau)$$
$$+ B^2\sin(\omega_0 t)\cos(\omega_0 t + \omega_0\tau) - AB\sin(\omega_0 t)\sin(\omega_0 t + \omega_0\tau))$$

$$= E(AB)E(\cos(\omega_0 t)\cos(\omega_0 t + \omega_0\tau)) - E(A^2)E(\cos(\omega_0 t)\sin(\omega_0 t + \omega_0\tau))$$
$$+ E(B^2)E(\sin(\omega_0 t)\cos(\omega_0 t + \omega_0\tau)) - E(AB)E(\sin(\omega_0 t)\sin(\omega_0 t + \omega_0\tau))$$

Scanned with CamScanner

Scanned with CamScanner

No.

$$\sigma_A^2 = E(A^2) - (E(A))^2 \to 0$$

$$\sigma_B^2 = E(B^2) - (E(B))^2 \to 0$$

But $\sigma_A^2 = \sigma_B^2 = \sigma^2 = E(A^2) = E(B^2)$

$$= -\sigma^2 E(\cos(\omega_0 t)\sin(\omega_0 t + \omega_0 \tau)) + \sigma^2 E(\sin(\omega_0 t)\cos(\omega_0 t + \omega_0 \tau))$$

$\sin(a-b) = \sin a \cos b - \cos a \sin b$

$$= \sigma^2 E(\sin(\omega_0 t - \omega_0 t - \omega_0 \tau))$$

$$= -\sigma^2 E(\sin(\omega_0 \tau))$$

$$= -\sigma^2 \sin(\omega_0 \tau)$$

$$= R_{xy}(\tau)$$

So, $X(t)$, $Y(t)$ are jointly wide-sense stationary. (#)

Scanned with CamScanner

Scanned with CamScanner

No.

Ex) let $X(t)$ be a wide-sense stationary with auto-correlation $R_{xx}(\tau) = e^{-a|\tau|}$, (وجب a)

let $Y(t) = X(t)\cos(\omega_0 t + \Theta)$, where Θ is uniform r.v. on $(-\pi, \pi)$, ω_0 is constant and Θ is indep. of $X(t)$, find the auto-correlation of $y(t)$.

$\cos(\omega_0 t + \Theta)$ ←

Sol:- $R_{yy}(t, t+\tau) = E(y(t)\, y(t+\tau))$

$= E(X(t)\cos(\omega_0 t + \Theta)\, X(t+\tau)\cos(\omega_0 t + \omega_0 \tau + \Theta))$

$= E(X(t)X(t+\tau))\, E(\cos(\omega_0 t + \Theta)\cos(\omega_0 t + \omega_0 \tau + \Theta))$

$= R_{xx}(t, t+\tau) \int_{-\pi}^{\pi} \left(\cos\left(\frac{2\omega_0 t + 2\Theta + \omega_0\tau}{2}\right) + \cos(\omega_0\tau)\right) \frac{1}{2\pi}\, d\theta$

$= R_{xx}(\tau) \cdot \frac{1}{2\pi}\left[0 + \cos(\omega_0\tau)(2\pi)\right]$

$= R_{xx}(\tau)\cos(\omega_0\tau)$

$= e^{-a|\tau|}\cos(\omega_0\tau) = R_{yy}(\tau)$.

So, it is wide-sense stationary.

Scanned with CamScanner

Scanned with CamScanner

* Cross-correlation function:-

The cross-correlation of $x(t)$, $y(t)$ is given by

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau))$$

* If $x(t)$, $y(t)$ are jointly wide-sense stationary, then

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau)) = R_{xy}(\tau)$$

* If $R_{xy}(\tau) = 0$, then $x(t)$, $y(t)$ are orthogonal processes.

* If $x(t)$, $y(t)$ are indep., then

$$R_{xy}(t, t+\tau) = E(x(t))\, E(y(t+\tau))$$

If $x(t)$, $y(t)$ are indep and at least wide-sense stationary, then

$$R_{xy}(\tau) = \bar{x}\,\bar{y} \text{ which is constant.}$$

Ref page w.s

⊛ properties:

1) $R_{xy}(-\tau) = R_{xy}(\tau)$.

2) $|R_{xy}(\tau)| \leq \sqrt{R_{xx}(0)\, R_{yy}(0)}$

3) $|R_{xy}(\tau)| \leq \frac{1}{2}[R_{xx}(0) + R_{yy}(0)]$

smile for life
smile for life
Scanned with CamScanner
Scanned with CamScanner

Ex Assume the two random process

$$x(t) = A\cos(\omega_0 t) + B\sin(\omega_0 t)$$
$$y(t) = B\cos(\omega_0 t) - A\sin(\omega_0 t),$$

where A, B are r.v's, ω_0 constant, A, B are uncorrelated, Zero-mean with same variance.

Find the cross-correlation $R_{xy}(t, t+\tau)$ and show that $x(t), y(t)$ are jointly wide-sense stationary.

$$E(A) = \underset{\text{indep}}{E(B)} = 0$$
$$E(AB) = E(A)E(B) = 0$$

$$\sigma_A^2 = \sigma_B^2 = \sigma^2$$

$$R_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau))$$
$$= E(\{A\cos(\omega_0 t) + B\sin(\omega_0 t)\}.$$
$$\{B\cos(\omega_0 t + \omega_0\tau) - A\sin(\omega_0 t + \omega_0\tau)\})$$

$$= E(AB\cos(\omega_0 t)\cos(\omega_0 t + \omega_0\tau) - A^2\cos(\omega_0 t)\sin(\omega_0 t + \omega_0\tau)$$
$$+ B^2\sin(\omega_0 t)\cos(\omega_0 t + \omega_0\tau) - AB\sin(\omega_0 t)$$
$$\sin(\omega_0 t + \omega_0\tau))$$

$$= E(AB)E(\cos(\omega_0 t)\cos(\omega_0 t + \omega_0\tau)) - E(A^2)E(\cos(\omega_0 t)\sin(\omega_0 t$$
$$+ E(B^2)E(\sin(\omega_0 t)\cos(\omega_0 t + \omega_0\tau)) - E(AB)E(\sin(\omega_0 t)\sin($$
$$(\omega_0 t$$

Scanned with CamScanner

Scanned with CamScanner

No.

$$\sigma_A^2 = E(A^2) - (E(A))^2 \to 0$$

$$\sigma_B^2 = E(B^2) - (E(B))^2 \to 0$$

But $\sigma_A^2 = \sigma_B^2 = \sigma^2 = E(A^2) = E(B^2)$

$$= -\sigma^2 E(\cos(\omega_0 t)\sin(\omega_0 t + \omega_0 \tau)) + \sigma^2 E(\sin(\omega_0 t)\cos(\omega_0 t + \omega_0 \tau))$$

$\sin(a-b) = \sin a \cos b - \cos a \sin b$

$$= \sigma^2 E(\sin(\omega_0 t - \omega_0 t - \omega_0 \tau))$$

$$= -\sigma^2 E(\sin(\omega_0 \tau))$$

$$= -\sigma^2 \sin(\omega_0 \tau)$$

$$= R_{xy}(\tau)$$

So, $X(t)$, $Y(t)$ are jointly wide-sense stationary. (#)

Scanned with CamScanner

Scanned with CamScanner

No.

(*) Covariance function :-

- The definition of covariance function can be extended to random process.

~ The auto-covariance function is

$$C_{xx}(t, t+\tau) = E(x(t)\, x(t+\tau)) \overset{*}{=} E(x(t))\, E(x(t+\tau))$$
$$= R_{xx}(t, t+\tau) - E(x(t))\, E(x(t+\tau)).$$

~ The cross covariance function is :-

$$C_{xy}(t, t+\tau) = E(x(t)\, y(t+\tau)) - E(x(t))\, E(y(t+\tau))$$
$$= \underbrace{R_{xy}(t, t+\tau) - E(x(t))\, E(y(t+\tau))}_{\text{Cross-correlation.}}$$

(*) If $x(t)$, $y(t)$ are wide-sense stationary, then

$$C_{xx}(t, t+\tau) = R_{xx}(t, t+\tau) - \bar{x}^2 = R_{xx}(\tau) - \bar{x}^2$$
$$C_{xy}(t, t+\tau) = R_{xy}(t, t+\tau) - \bar{x}\bar{y}$$

smile

Scanned with CamScanner

Scanned with CamScanner

No.

* If $X(t), Y(t)$ are jointly wide sense stationary

$$C_{xx}(t, t+\tau) = R_{xx}(\tau) - \bar{X}^2$$
$$C_{yy}(t, t+\tau) = R_{yy}(\tau) - \bar{Y}^2$$
$$C_{xy}(t, t+\tau) = R_{xy}(\tau) - \bar{X}\bar{Y}$$

* For any random process, the variance can be calculated by

$$C_{xx}(t, t+\tau)\Big|_{\tau=0}$$

مهم

* Recall:-

$$Cov(X, Y) = E(XY) - E(X)E(Y)$$

→ Properties:-

→ $Cov(X, X) = E(XX) - (EX)(EX)$

$$= EX^2 - (EX)^2$$

$$= VarX = \sigma_X^2$$

Now:

$$C_{xx}(t, t+\tau) = E(x(t)\,x(t+\tau)) - E(x(t))\,E(x(t+\tau))$$

$$C_{xx}(t, t+\tau)\Big|_{\tau=0} = E(x(t)\,x(t)) - E(x(t))\,E(x(t))$$

$$= E(x^2(t)) - (E(x(t)))^2$$

$$= Var(x(t)).$$

Scanned with CamScanner

Scanned with CamScanner

No.

(*) For any random process $x(t)$, $y(t)$,
if $x(t)$, $y(t)$ are indep,
then $C_{xy}(t, t+\tau) = 0$

(*) Discrete-time random process:-

It is defined as $x(t) = x(nT_s)$; $n = 0, \pm 1, \pm 2$

~> If $t_1 = n_1 T_s$, $t_2 = n_2 T_s$

$\rightarrow \tau = t_2 - t_1$,

$= n_2 T_s - n_1 T_s = (n_2 - n_1) T_s = \underline{K t_s}$

Here, the differences τ can only have discrete values KT_s.

So, for a DT-random process, we have mean
$= E(x(nT_s))$

auto-correlation $= R_{xx}(t, t+\tau)$

$= R_{xx}(nT_s, nT_s + KT_s)$

$= E(x(nT_s) X(nT_s + KT_s))$

Cross-correlation =

$R_{xy}(t, t+\tau) = R_{xy}(nT_s, nT_s + KT_s)$

$= E(x(nT_s)\, y(nT_s + KT_s))$

Scanned with CamScanner

Scanned with CamScanner

No.

* auto-Covarian

* auto-Covariance $= C_{xx}(t, t+\tau)$
$= R_{xx}(nT_s, nT_s + kT_s) - E(x(nT_s))\, E(x(nT_s + kT$

- Cross-Corariance $= C_{xy}(t, t+\tau)$
$= R_{xy}(nT_s, nT_s + kT_s)$
$- E(x(nT_s))\, E(y(nT_s + kT_s))$

* For process that are wide-sense stationary

$E(x(nT_s)) = \bar{x}$, $E(y(nT_s)) = \bar{y}$

$R_{xx}(nT_s, nT_s + \underbrace{kT_s}_{\tau}) = R_{xx}(\underbrace{kT_s}_{\tau})$

$R_{yy}(nT_s, nT_s + kT_s) = R_{yy}(kT_s)$

$C_{xx}(nT_s, nT_s + kT_s) = R_{xx}(kT_s) - \bar{x}^2$

$C_{yy}(nT_s, nT_s + kT_s) = R_{yy}(kT_s) - \bar{y}^2$

$C_{xy}(nT_s, nT_s + kT_s) = R_{xy}(kT_s) - \bar{x}\bar{y}$

Joint

Scanned with CamScanner

Scanned with CamScanner