Numerical Techniques
+ Applied Numerical Analysis.

Introduction:

$$\int_1^2 \sin x^2 dx \quad , \quad \int_0^1 e^{x^3} dx \ldots$$

$$2x - 6 = 0 \rightarrow x = 3$$

$$x^3 + x - 3 = 0$$

$$x = e^{-x}$$

Modeling

Parachutist Problem.



m : mass (Kg)

V: velocity

$Fu \propto V \rightarrow Fu = cV$

c: drag coefficient Kg/s

$$F_{resultant} = ma \qquad a = \frac{dv}{dt}$$

$$mg - cV = m\frac{dv}{dt}$$ Differential Equation.

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$m\frac{dv}{dt} = mg - cv$$

$$m\frac{dv}{dt} + cv = mg$$

$\div m \longrightarrow$ $\boxed{\frac{dv}{dt} + \frac{c}{m}v = g}$ First order LDE

$\times \longrightarrow e^{\int \frac{c}{m} dt} = e^{\frac{c}{m}t}$

$$e^{\frac{c}{m}t}\frac{dv}{dt} + \frac{c}{m}e^{\frac{c}{m}t}v = g e^{\frac{c}{m}t}$$

$y' + P(x)y = g(x)$

$\rightarrow * \; e^{\int P(x)dx}$

$$\frac{d}{dt}\left(e^{\frac{c}{m}t}v\right) = g e^{\frac{c}{m}t}$$

Integrate $\longrightarrow$ $e^{\frac{c}{m}t}v = g \cdot \frac{m}{c} e^{\frac{c}{m}t} + K$

$\div e^{\frac{c}{m}t} \longrightarrow$ $v = \frac{gm}{c} + K e^{-\frac{c}{m}t}$

But $v(0) = 0 \Longrightarrow \frac{gm}{c} + K = 0$

$$\Rightarrow K = -\frac{gm}{c}$$

$$\therefore v = \frac{gm}{c} - \frac{gm}{c} e^{-\frac{c}{m}t}$$

$$\boxed{v = \frac{gm}{c}\left(1 - e^{-\frac{c}{m}t}\right)}$$

as $t \longrightarrow \infty \Rightarrow v \longrightarrow \frac{gm}{c}$



KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Taylor and Maclaurin series:

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \frac{f'''(a)(x-a)^3}{3!} + \cdots$$

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)(x-a)^k}{k!}$$

Taylor series of $f(x)$ about $x=a$.

$a = 0 \longrightarrow$ $f(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \cdots$

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)\,x^k}{k!}$$

Maclaurin series

---

e.g: Find Taylor Expansion of $f(x) = e^x$ about $x = 1$.

$$f(x) = f(1) + f'(1)(x-1) + \frac{f''(1)(x-1)^2}{2!} + \frac{f'''(1)(x-1)^3}{3!} + \cdots$$

$f(x) = e^x \rightarrow f(1) = e$

$f'(x) = e^x \rightarrow f'(1) = e$

$f''(x) = e^x \rightarrow f''(1) = e$

$$\therefore f(x) = e + e(x-1) + \frac{e(x-1)^2}{2!} + \frac{e(x-1)^3}{3!} + \cdots$$

$$e^x = \sum_{k=0}^{\infty} \frac{e(x-1)^k}{k!}$$

KAMEL FEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Find Maclaurin series of the given function.

1) $f(x) = e^x$

$$f(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \cdots$$

$f(x) = e^x \longrightarrow f(0) = e^0 = 1$

$f'(x) = e^x \longrightarrow f'(0) = 1$

$f''(x) = e^x \longrightarrow f''(0) = 1$

$\vdots$

$$\therefore f(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

$$e^x = \sum_{K=0}^{\infty} \frac{x^K}{K!}$$

2) $f(x) = \sin x$

$$f(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2!} + \frac{f'''(0)x^3}{3!} + \cdots$$

$f(x) = \sin x \longrightarrow f(0) = 0$

$f'(x) = \cos x \longrightarrow f'(0) = 1$

$f''(x) = -\sin x \longrightarrow f''(0) = 0$

$f'''(x) = -\cos x \longrightarrow f'''(0) = -1$

$f^{(4)}(x) = \sin x \longrightarrow f^{(4)}(0) = 0$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$f(x) = 0 + 1x + \frac{(0)x^2}{2!} + \frac{(-1)x^3}{3!} + \frac{0x^4}{4!} + \frac{1x^5}{5!} + \cdots$$

$$f(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$$

$$\sin x = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!}$$

Rules:

1) $$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} \qquad x \in \mathbb{R}$$

2) $$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!}$$

3) $$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!}$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

4) $\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \cdots$

$$= \sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!}$$

5) $\cosh x = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \cdots$

$$= \sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!}$$

$a + ar + ar^2 + \cdots = \frac{a}{1-r} \quad |r|<1$

6) $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots$

$|x|<1$

$\Leftrightarrow -1 < x < 1$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k \qquad |x|<1$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

(7)

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots \qquad x \in \mathbb{R}$$

$$x = 1 \rightarrow e = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots$$

$$x = \frac{1}{2} \rightarrow e^{\frac{1}{2}} = \sqrt{e} = 1 + \left(\frac{1}{2}\right) + \frac{\left(\frac{1}{2}\right)^2}{2!} + \frac{\left(\frac{1}{2}\right)^3}{3!} + \cdots$$

$$\int_0^1 e^{x^2} dx$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

$$e^{x^2} = 1 + x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} + \cdots$$

$$e^{x^2} \sim 1 + x^2$$

$$\int_0^1 e^{x^2} dx = \int_0^1 (1 + x^2) dx = \cdots$$

END

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Recall

1) $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$

$= \sum_{k=0}^{\infty} \frac{x^k}{k!}$

2) $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$

$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!}$

3) $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$

$= \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!}$

~~4) $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots$~~

4) $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots$, $|x| < 1$

$= \sum_{k=0}^{\infty} x^k$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Set of Numbers

$\mathbb{Z} = \{\ldots, -2, -1, 0, 1, 2, 3, \ldots\}$ set of integers

$\mathbb{Q} = \{\frac{a}{b} \mid a, b \in \mathbb{Z}, b \neq 0\}$ set of rationals

$\hookrightarrow \frac{2}{5}, -\frac{1}{4}, \frac{22}{7}, 8, 0,$

$2.5, 0.\overline{3} = 0.3333\ldots$



repeating decimal

$0.4444\ldots = 0.\overline{4}$

$0.9999\ldots = 0.\overline{9}$

$2.3275275275\ldots = 2.3\overline{275}$

terminating decimal

1.42, 7.6,

3, 2.31750

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$\bar{Q}$: irrational numbers

- $\sqrt{2}, \sqrt{3}, \sqrt{5}, \cdots$
- $\pi \approx \frac{22}{7}$
- $e \approx 2.7$
- $5.141441444144441\cdots$

$\mathbb{R} = Q \cup \bar{Q}$ = real numbers

$Q \cap \bar{Q} = \phi$

e.g.: Show that $0.\bar{9} = 1$?

$$0.\bar{9} = 0.9999999\cdots$$

$$= 0.9 + 0.09 + 0.009 + \cdots$$

$$\frac{0.09}{0.9} = 0.1 \quad , \quad \frac{0.009}{0.09} = 0.1, \cdots$$

G.S $\quad r = 0.1$

$a = 0.9$

$$0.\bar{9} = \frac{0.9}{1 - 0.1} = \frac{0.9}{0.9} = 1$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots \quad |x| < 1$$

$$a + ar + ar^2 + \cdots = \frac{a}{1-r}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Write $0.\overline{4}$ in the form $\frac{a}{b}$, $a, b \in \mathbb{Z}$ (4)

$0.\overline{4} = 0.4 + 0.04 + 0.004 + \cdots$

g.s $a = 0.4$, $r = 0.1$

$$0.\overline{4} = \frac{0.4}{1 - 0.1} = \frac{0.4}{0.9} = \frac{4}{9}$$

e.g: write $0.\overline{23}$ in the form $\frac{a}{b}, a, b \in \mathbb{Z}$?

$0.\overline{23} = 0.23 + 0.0023 + 0.000023 + \cdots$

g.s $r = 0.01$, $a = 0.23$

$$\therefore 0.\overline{23} = \frac{0.23}{1 - 0.01}$$

$$= \frac{0.23}{0.99}$$

$$= \frac{23}{99}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Accuracy and Precision:

Accuracy: The closeness of computed or measured values to the true value

Precision: The closeness of computed or measured value to each other.

True length of the pen = 10 cm

10.1, 9.9, 10.01, 9.95 high accuracy and high precision

20.1, 20.2, 19.9, 20.01 Low ~~high~~ accuracy and high precision



KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Error Definitions:

If X is the true value and $\hat{X}$ is the approximation, then

1) True Error: $E_t = X - \hat{X}$

2) Absolute True Error: $|E_t| = |X - \hat{X}|$

3) True Relative Error: $\epsilon_t = \frac{X - \hat{X}}{X}$

4) Absoule True Relative Error: $|\epsilon_t| = \left|\frac{X - \hat{X}}{X}\right|$

5) $|\epsilon_t| = \left|\frac{X - \hat{X}}{X}\right| * 100\%$

Absolute true percent relative error.

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: If the true weight of car is 1000 kgs and the approximation was 999 kgs.

Also, the true weight of a boy 20 kgs and the approximation was 19 kgs.

Find the true error and the true percent relative error for the car and the boy.

Sol:

Car

True value = 1000 kg = X

Approximated value = 999 kgs = $\tilde{X}$

$$E_t = X - \hat{X} = 1000 - 999 = 1 kg$$

$$\epsilon_t = \frac{X - \hat{X}}{X} * 100\% = \frac{1}{1000} * 100\% = 0.1\%$$

Boy:

True value: $Y = 20$ kgs

Approximated value $\hat{Y} = 19$ kg.

$$E_t = Y - \hat{Y} = 20 - 19 = 1 kg$$

$$\epsilon_t = \frac{Y - \hat{Y}}{Y} * 100\% = \frac{1}{20} * 100\% = 5\%$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Def: If X^{New} and X^{old} are two approximations for a quantity then

1) The Approximate Error : $E_a = X^{New} - X^{old}$

2) The Relative Approximate Error

$$\epsilon_a = \frac{X^{New} - X^{old}}{X^{New}}$$

$$= \frac{X^{New} - X^{old}}{X^{New}} * 100\%$$

e.g: $X^{current} = 5.5$, $X^{previous} = 3.5$

find ϵ_a

$$\epsilon_a = \frac{5.5 - 3.5}{5.5} = \frac{2}{5.5} \simeq 0.36 = 0.36 * 100\%$$

$$= 36\%$$

END

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Errors (Part 2)

ϵ : Tolerance

ϵ_s : Tolerance related to significant figures.

Scarborough criterion:

To get a ~~beta~~ result to be correct to at least n significant figures, do iterations till you get $|\epsilon_a| < \epsilon_s$ (i.e stop).

where $\epsilon_s = (0.5 \times 10^{-n}) * 100\%$

$$\epsilon_s = (0.5 \times 10^{2-n})\%$$

Recall $\epsilon_a\% = \left| \frac{\text{current app.} - \text{previous app.}}{\text{current approximation}} \right| * 100\%$

$$\epsilon_t\% = \left| \frac{\text{true value} - \text{approximation}}{\text{true value}} \right| * 100\%$$

KAMEL ELEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Approximate the value of $e^{0.3}$ by using (2) Maclaurin Series of e^x, start by using 2 terms and on of the series.

Use Scarborough Criterion so that your answer to be correct to at least 3 significant figures.

Sol:

The answer will be correct to at least 3 significant figure, $\epsilon_a\% \leq (0.5 \times 10^{2-3})\% = \epsilon_s$

i.e stop when $\epsilon_a\% < (0.05)\%$ ✓

Now $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$

if $x = 0.3 \Rightarrow$ $e^{0.3} = 1 + 0.3 + \frac{(0.3)^2}{2!} + \frac{(0.3)^3}{3!} + \frac{(0.3)^4}{4!} + \cdots$

1st iteration (2 terms):

$e^{0.3} \simeq 1 + 0.3$

$e^{0.3} \simeq 1.3$

we cant compute ϵ_a here.

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2nd iteration (3 terms)

$$e^{0.3} \simeq \underbrace{1 + 0.3}_{\downarrow} + \frac{(0.3)^2}{2!}$$

$$e^{0.3} \simeq 1.3 + \frac{(0.3)^2}{2!}$$

$$\therefore \boxed{e^{0.3} \simeq 1.345}$$

$$\epsilon_a \% = \left|\frac{1.345 - 1.3}{1.345}\right| * 100\%$$

$$\simeq 3.3457\% \not< 0.05\% = \epsilon_s$$

3rd iteration (4 terms):

$$e^{0.3} \simeq \underbrace{1 + 0.3 + \frac{(0.3)^2}{2!}} + \frac{(0.3)^3}{3!}$$

$$e^{0.3} \simeq 1.345 + \frac{(0.3)^3}{3!}$$

$$\boxed{e^{0.3} \simeq 1.3495}$$

$$\epsilon_a = \left|\frac{1.3495 - 1.345}{1.3495}\right| * 100\%$$

$$\simeq 0.333 \not< 0.05\% = \epsilon_s$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

4th iteration (5 terms):

$$e^{0.3} \cong 1 + 0.3 + \frac{(0.3)^2}{2!} + \frac{(0.3)^3}{3!} + \frac{(0.3)^4}{4!}$$

$$\therefore e^{0.3} \cong 1.34495 + \frac{(0.3)^4}{4!}$$

$$\boxed{e^{0.3} \cong 1.3498375}$$

$$\epsilon_a = \left| \frac{1.3498375 - 1.3495}{1.3498375} \right| * 100\%$$

$$\epsilon_a \cong 0.02\% < 0.05\% \rightarrow \text{Stop}$$

$$\therefore \boxed{e^{0.3} \cong 1.3498375}$$

To find the (true value)? use ~~calculator~~ Calculator

Could be used as true value: 1.349858808

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Using the first three non-zero terms of Maclaurin series for $\sin x$, approximate the value of the integral

$$\int_1^{1.2} \frac{\sin x}{x}\,dx$$ (round off your answer to 5 decimal places).

And then compute the relative true error ($\epsilon_t\%$)

Solution

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$$

$$\sin x \cong x - \frac{1}{6}x^3 + \frac{1}{120}x^5$$

$$\Rightarrow \frac{\sin x}{x} \cong 1 - \frac{1}{6}x^2 + \frac{1}{120}x^4$$

$$\Rightarrow \int_1^{1.2} \frac{\sin x}{x}\,dx \cong \int_1^{1.2} \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4\right) dx$$

$$\Rightarrow \int_1^{1.2} \frac{\sin x}{x}\,dx \cong 0.16204 \quad \text{(to 5 decimal places)}$$

True value: $\int_1^{1.2} \frac{\sin x}{x}\,dx = 0.161964128\,6$

$$\therefore \epsilon_t \cong \left|\frac{0.16196 - 0.16204}{0.16196}\right| * 100\% = 0.04\%$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Chopping and Rounding off:

Def: A number X is said to be in normalized decimal form (decimal floating point form) if

$$X = (0.d_1 d_2 d_3 \cdots) \times 10^{n}$$

(fraction part (mantissa)) (exponential part)

where $d_1, d_2, \cdots \in \{0, 1, 2, \cdots, 9\}$

and $d_1 \neq 0$ (non-zero), $n \in \mathbb{Z}$ (n ← integer)

i.e $X = 0.\square\square\square\cdots \times 10^{n}$

e.g: Normalize the following numbers.

1) 145.7632×10^{0}

$= 0.1457632 \times 10^{3}$

2) 12.3×10^{-10}

$= 0.123 \times 10^{-10+2}$ $= 0.123 \times 10^{-8}$

3) 0.00027×10^{-15}

$= 0.27 \times 10^{-15-3}$

$= 0.27 \times 10^{-18}$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: After normalizing, chop the following numbers to 4 significant digits. [7]

1) $0.0007247783 = 0.7247783 \times 10^{-3}$

$= 0.7247 \times 10^{-3}$

$= 0.0007247$

2) 7.8135×10^{5}

$= 0.78135 \times 10^{5+1}$

$= 0.7813|5 \times 10^{6}$

$= 0.7813 \times 10^{6}$

3) 0.0085482

$= 0.8548|2 \times 10^{-2}$

$= 0.854 \times 10^{-2}$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Round off the following numbers to 4 significant digits.

1) 0.624783×10^{6}

$\simeq 0.6248 \times 10^{6}$

2) 0.179231×10^{-7}

$\simeq 0.1792 \times 10^{-7}$

3) 0.854851×10^{20}

$= 0.8549 \times 10^{20}$

4) 0.85485×10^{-5}

$= 0.8548 \times 10^{-5}$.

- END -

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Truncation Errors and Taylor series

Truncation error occurs if we approximate mathematical procedure.

examples:

1) $e^{0.3} = \underbrace{1 + 0.3 + \frac{(0.3)^2}{2!}}_{\text{if approximation of } e^{0.3}} + \underbrace{\frac{(0.3)^3}{3!} + \cdots}_{\text{Truncation error}}$

2) If $f(x) = x^2$

$f'(1) = 2$ ← $f'(1) = \lim_{h \to 0} \frac{f(1+h) - f(1)}{h}$

if $h = 0.1 \Rightarrow f'(1) \simeq \frac{f(1.1) - f(1)}{0.1}$

$\simeq \frac{1.21 - 1}{0.1}$

$\simeq \frac{0.21}{0.1}$.

$\simeq 2.1$

Truncation error

3) $\int_a^b f(x)dx \simeq$ area of trapezoid



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Taylor series (Taylor Expansion)

If f is a smooth function ($f^{(n)}$ exists for any n)

$a \in \mathbb{R}$ then

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \frac{f'''(a)(x-a)^3}{3!} + \cdots$$

$$+ \frac{f^{(n)}(a)(x-a)^n}{n!} + R_n$$

where $R_n = \frac{f^{(n+1)}(\xi)(x-a)^{n+1}}{(n+1)!}$ (Lagrange remainder form)

where ξ between a and x.

Now ~~let~~ if $a = x_i$ and $x = x_{i+1}$

and $x_{i+1} - x_i = h$

we get

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \frac{f'''(x_i)h^3}{3!}$$

$$+ \frac{f^{(4)}(x_i)h^4}{4!} + \cdots + \frac{f^{(n)}(x_i)h^n}{n!} + R_n$$

or

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \cdots + \frac{f^{(n)}(x_i)h^n}{n!} + O(h^{n+1})$$

$R_n = O(h^{n+1})$ ≡ truncation error

↗ order of error

Scanned with CamScanner

CamScanner الممسوحة ضوئيا بـ

$$* \; f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \frac{f'''(x_i)h^3}{3!} + \cdots$$

$P_0(x)$ — Taylor polynomial of order 0.

$R_0 = O(h)$

$$* \; f(x_{i+1}) = \underbrace{f(x_i) + f'(x_i)h}_{P_1(x)} + \underbrace{\frac{f''(x_i)h^2}{2!} + \frac{f'''(x_i)h^3}{3!} + \cdots}_{R_1 = O(h^2)}$$

$P_1(x)$ — Taylor polynomial of order 1

$$* \; f(x_{i+1}) = \underbrace{f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!}}_{P_2(x)} + \underbrace{\frac{f'''(x_i)h^3}{3!} + \cdots}_{R_2 = O(h^3)}$$

$P_2(x)$ — Taylor polynomial of order 2

⋮

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g.: Use zero-through second order Taylor series expansions to approximate $f(2.1)$ and where $f(x) = 2.5x^2 + 3x + 7$ take $x_i = 2$. Compute ϵ_a % in each step (if possible).

Sol:

$x_i = 2$, $x_{i+1} = 2.1$ want to predict $f(x_{i+1})$

$h = x_{i+1} - x_i = 2.1 - 2$

$\Rightarrow h = 0.1$

$\therefore f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \cdots$

$\Rightarrow f(2.1) = f(2) + f'(2)(0.1) + \frac{f''(2)(0.1)^2}{2!} + \cdots$

1st iteration: (order 0 $\equiv$ 1 term)

$f(2.1) \cong f(2)$

$f(2.1) \cong 23$

$f(x) = 2.5x^2 + 3x + 7$

2nd iteration (order 1 $\equiv$ 2 terms)

$f(2.1) \cong f(2) + f'(2)(0.1)$

$\cong 23 + 13(0.1)$

$\cong 23 + 1.3$

$f(2.1) \cong 24.3$

$f'(x) = 5x + 3$

$\epsilon_a\% = \left| \frac{24.3 - 23}{24.3} \right| * 100\% \cong 5.35\%$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

3rd iteration (order 2 ≡ 3 terms)

$$f(2.1) \simeq f(2) + f'(2)(0.1) + \frac{f''(2)(0.1)^2}{2!}$$

$f''(x) = 5$

$$\simeq 24.3 + \frac{(5)(0.1)^2}{2}$$

$$\simeq 24.325$$

$$\epsilon_a\% = \left|\frac{24.325 - 24.3}{24.325}\right| * 100\% \simeq 0.1\%$$

$f(x) = 2.5x^2 + 3x + 7$

$f(2.1) = 24.325$ i.e $\epsilon_t = 0\%$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Using Taylor polynomial of order (2), approximate $\sin(\frac{\pi}{6})$ taking $x_i = \frac{\pi}{4}$.

Then compute ϵ_t %.

$x_{i+1} = \frac{\pi}{6}$, $x_i = \frac{\pi}{4} \Rightarrow h = x_{i+1} - x_i$

$h = \frac{\pi}{6} - \frac{\pi}{4}$

$h = -\frac{\pi}{12}$, $f(x) = \sin x$

$$f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \dots$$

$f'(x) = \cos x$

$f''(x) = -\sin x$

$$\therefore \sin\frac{\pi}{6} \simeq \sin\frac{\pi}{4} + (\cos\frac{\pi}{4})(\frac{-\pi}{12}) + \frac{(-\sin\frac{\pi}{4})(\frac{-\pi}{12})^2}{2!}$$

$$\sin\frac{\pi}{6} \simeq 0.497854491-4$$

$$\epsilon_t = \frac{0.5 - 0.497854491-4}{0.5} * 100\ \%$$

$$\epsilon_t = 0.45\ \%$$

END

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Bracketing Methods

Introduction:

* The equation $f(x)=0$, has a root (C) if

$f(c)=0$.

Graphically the graph of $y=f(x)$ meets the x-axis at c. (c: real root)









Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* $f(x) = g(x)$ this equation has root(s) if any at x-coordinates of the point(s) of intersection.



$x - e^{-x} = 0 \longrightarrow x = e^{-x}$



$x - \cos x = 0 \longrightarrow x = \cos x$



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Bolzano Theorem:

If $f(x)$ is continuous on $[a,b]$, (1)

$$f(a)\,f(b) < 0 \quad (2)$$

then $\exists\, c \in (a,b)$ such that $f(c)=0$

i.e c is a root of $f(x)=0$





3-root





Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Bolzano Theorem:

If $f(x)$ is continuous on $[a,b]$,

(1)

$$f(a)\,f(b) < 0$$

(2)

then $\exists\, c \in (a,b)$ such that $f(c)=0$

i.e c is a root of $f(x)=0$





3-root





Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g.: Show that $x^3+x^2-3=0$ has a root in the interval $(1,2)$

Sol:

Let $f(x) = x^3+x^2-3$, $x \in [1,2]$

f is continuous on $[1,2]$

$f(1) = -1$

$f(2) = 9$

$f(1)\,f(2) = -9 < 0$

$\therefore$ By Bolzano theorem

$\exists\ c \in (1,2)$ such that $f(c)=0$

$\Rightarrow c^3+c^2-3=0$, $c \in (1,2)$

i.e there exists a root $c \in (1,2)$ for the equation $x^3+x^2-3=0$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

1) Bisection Method

If $f(x)$ satisfies Bolzano conditions

i.e f is continuous on $[a,b]$ and $f(a)f(b)<0$

then f has a root between a and b

Let $x_L = a$ and $x_u = b$

1st iteration:

$$x_r = \frac{x_L + x_u}{2}$$



$$|x_r - \text{true root}| < \frac{b-a}{2}$$

assume $f(x_r) < 0$

2nd iteration



$$x_r = \frac{x_L + x_u}{2}$$

$$|x_r - \text{true root}| < \frac{b-a}{2^2}$$

assume $f(x_r) > 0$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

3rd iteration :

$$x_r = \frac{x_L + x_u}{2}$$



$$|\text{true root} - x_r| < \frac{b-a}{2^3}$$

⋮

nth iteration

$$|\text{true root} - x_r| < \frac{b-a}{2^n}$$

if ϵ ~~the~~ is the Tolerance for true error

and $\frac{b-a}{2^n} \leq \epsilon \Rightarrow |\text{true root} - x_r| \leq \epsilon$

$$\Rightarrow \ln\left(\frac{b-a}{2^n}\right) \leq \ln \epsilon$$

$$\Rightarrow \ln(b-a) - n\ln 2 \leq \ln \epsilon$$

$$n \ln 2 \geq \ln(b-a) - \ln \epsilon \Rightarrow \boxed{n \geq \frac{\ln(b-a) - \ln \epsilon}{\ln 2}}$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2] False-Position Method:

If f is continuous on $[a,b]$

and $f(a)f(b)<0 \Rightarrow \exists\, c \in (a,b) : f(c)=0$

According to the figure

$f(x_L) < 0$

$f(x_u) > 0$] $f(x_L)\,f(x_u) < 0$

f is continuous.

to find x_r (the approximation of the true root of $f(x)=0$)



slope AB : $\dfrac{f(x_L)-f(x_u)}{x_L - x_u}$ ----①

Also slope AB : $\dfrac{f(x_u)-0}{x_u - x_r} = \dfrac{f(x_u)}{x_u - x_r}$ --②

from ① and ②

$$\frac{f(x_L)-f(x_u)}{x_L - x_u} = \frac{f(x_u)}{x_u - x_r}$$

KAMEL FLEIFEL

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

$$\Rightarrow \quad \frac{x_L - x_u}{f(x_L) - f(x_u)} = \frac{x_u - x_r}{f(x_u)}$$

$$\xrightarrow{* f(x_u)} \quad \frac{f(x_u)(x_L - x_u)}{f(x_L) - f(x_u)} = x_u - x_r$$

$$\Rightarrow \quad \boxed{x_r = x_u - \frac{f(x_u)(x_L - x_u)}{f(x_L) - f(x_u)}}$$

2nd iteration

$x_L = x_r \rightarrow f(x_L) < 0$

$x_u = x_u \rightarrow f(x_u) > 0$



KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Show that the equation $x^3 + x^2 - 3 = 0$ has a root in the interval $(1, 1.5)$, then use false position method to approximate the root of this equation by rounding off the answer of second iteration to (4) decimal places.

Sol:

Let $f(x) = x^3 + x^2 - 3$, $x \in [1, 1.5]$

f is continuous on $[1, 1.5]$

$f(1) = -1$

$f(1.5) = 2.625$

$\Big]\ f(1)f(1.5) < 0$

$\Rightarrow \exists\ c \in (1, 1.5)$ such that $f(c) = 0$

$\Rightarrow c^3 + c^2 - 3 = 0$ (i.e has root in $(1, 1.5)$)

1st iteration:

$x_L = 1 \longrightarrow f(x_L) = -1$

$x_u = 1.5 \longrightarrow f(x_u) = 2.625$

$\Big]\ f(x_L)f(x_u) < 0$

$$x_r = x_u - \frac{f(x_u)(x_L - x_u)}{f(x_L) - f(x_u)}$$

$$x_r = 1.5 - \frac{2.625(1 - 1.5)}{-1 - 2.625}$$

$\Rightarrow x_r = 1.137931$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$f(x) = x^3 + x^2 - 3$$

$x_r = 1.137931$

$f(x_r) = -0.231621$

2nd iteration:

$x_L = 1.137931 \Longrightarrow f(x_L) = -0.231621$

$x_u = 1.5 \longrightarrow f(x_u) = 2.625$

$f(x_L)\, f(x_u) < 0$

$$x_r = 1.5 - \frac{2.625(1.137931 - 1.5)}{-0.231621 - 2.625}$$

$x_r \cong 1.1673$

True root 1.174559411

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Use false position method, approximate [5]
the root of $x + \sin x = 1$ $\quad x \in [0, \frac{\pi}{2}]$
by doing one iteration.

Sol:

$x + \sin x - 1 = 0$

$f(x) = x + \sin x - 1$

f is Cont. on $[0, \frac{\pi}{2}]$

$f(0) = -1$

$f(\frac{\pi}{2}) = 1.5707963$] $f(0) f(\frac{\pi}{2}) < 0$

$\Rightarrow \exists\, c \in (0, \frac{\pi}{2}) : f(c) = 0$

1st iteration:

$x_L = 0 \longrightarrow f(x_L) = -1$

$x_u = \frac{\pi}{2} \longrightarrow f(x_u) = 1.5707963$ } $\Rightarrow f(x_L) f(x_u) < 0$

$$x_r = \frac{\pi}{2} - \frac{1.5707963\,(0 - \frac{\pi}{2})}{0 - \frac{\pi}{2}}$$

$x_r = 0.61101559$

True value
0.5109734

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Open Methods

I) Simple fixed point method:

$f(x) = 0$

Rearrange ↓

$x = g(x)$

initial guess given. (x_0)

For example

$x^2 - 2x - 3 = 0$

$x = \sqrt{2x+3}$ ← $x = g_1(x)$

$x = \frac{x^2-3}{2}$ ← $x = g_2(x)$

$x^2 = 2x + 3$

$\div x \rightarrow x = \frac{2}{x} + \frac{3}{x}$ ← $x = g_3(x)$

$x_{i+1} = g(x_i)$ (x_0 given) (initial guess)

$x_0, x_1, x_2, \ldots$ → converges / diverges

this sequence will converge if

$|g'(\text{near the root})| < 1$.

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: To, find a root of $x^2-2x-3=0$

Use $x_{i+1} = \sqrt{2x_i+3}$ $\quad x_0 = 0$

$x_1 = \sqrt{2x_0+3}$

$x_2 = \sqrt{2x_1+3}$

$x_3 = \sqrt{2x_2+3}$

$\left|\frac{x_i^{New} - x_i^{old}}{x_i^{New}}\right| * 100\%$

$\epsilon_t = \left|\frac{3 - x_i}{3}\right| * 100\%$

i	x_i	ϵ_a %	ϵ_t %
0	0	-	100 %
1	1.732051	100 %	42.3 %
2	2.542460		
3	2.843399		
4	2.947338		
5	2.982394		

true root 3

3

Notice that $g(x) = \sqrt{2x+3}$

and $g'(x) = \frac{2}{2\sqrt{2x+3}}$

$g'(x) = \frac{1}{\sqrt{2x+3}} < 1$

KAMEL FLEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

e.g: To find a root of the equation $x^3 - x + 1 = 0$. by using simple fixed point method and

using $x_{i+1} = x_i^3 + 1$, $x_0 = 1$

find $x_0, x_1, x_2, \ldots$ (is this sequence convergent).

sol:

$$x_{i+1} = x_i^3 + 1$$

$x_0 = 1$, $x_1 = 2$, $x_2 = 9$, $x_3 = \ldots$

$x_4 =$, x_5 $= x_6 =$

diverges

KAMEL PLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2 Newton - Raphson Method:

If f is differentiable function

to find the root of $f(x)=0$ (if it exists)

taking x_0 as initial guess

$$\frac{f'(x_i)}{1} = \frac{f(x_i) - 0}{x_i - x_{i+1}}$$

~~$f(x_i) = f'$~~

$$\frac{1}{f'(x_i)} = \frac{x_i - x_{i+1}}{f(x_i)}$$

$f(x_i)$

$$\Longrightarrow \quad \frac{f(x_i)}{f'(x_i)} = x_i - x_{i+1}$$

$$\boxed{x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}}$$



x_0

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$\vdots$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

E.g Use Newton - Raphson method, to solve the equation $x = e^{-x}$ take the initial guess $x_0 = 0$, do 4 iterations and compute $\epsilon_a\%$, $\epsilon_t\%$ where it is possible.

Solution

$$f(x) = x - e^{-x} = 0$$

$$f'(x) = 1 + e^{-x}$$

$$x_{i+1} = x_i - \frac{x_i - e^{-x_i}}{1 + e^{-x_i}} \qquad x_0 = 0$$

$x_0 = 0$

$$x_1 = 0 - \frac{0 - e^{0}}{1 + e^{0}} = 0.5$$

x_2

x_3

x_4

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

i	x_r	$\epsilon_t\%$	$\epsilon_a\%$
0	0	100%	—
1	0.5	$\frac{\text{true} - 0.5}{\text{true}} \times 100$	$\frac{0.5 - 0}{0.5} \times 100$
2	0.566311		
3	0.567143		
4	0.567143		

True value,

0.567143290 4

END -

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

3) secant - Method

x_{i-1} , x_{i-2} two initial guesses

$A(x_i, f(x_i))$, $B(x_{i-1}, f(x_{i-1}))$



$$\text{slope } AB = \frac{f(x_{i-1}) - f(x_i)}{x_{i-1} - x_i} \quad \text{--- (1)}$$

$$\text{Also slope } AB = \frac{f(x_i) - 0}{x_i - x_{i+1}} \quad \text{--- (2)}$$

from (1) and (2)

$$\frac{f(x_{i-1}) - f(x_i)}{x_{i-1} - x_i} = \frac{f(x_i)}{x_i - x_{i+1}}$$

$$\Rightarrow \frac{x_{i-1} - x_i}{f(x_{i-1}) - f(x_i)} = \frac{x_i - x_{i+1}}{f(x_i)}$$

$$\xrightarrow{* f(x_i)} \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)} = x_i - x_{i+1}$$

$$\Rightarrow \boxed{x_{i+1} = x_i - \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)}} \quad i = 0, 1, 2, 3, \ldots$$

x_{-1}, x_0 are given

x_{-1} , x_0 , x_1 , x_2 , x_3 , - - - → Conv. / div.

KAMEL FLEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

e.g: Use secant method to estimate the root of $e^{-x} - x = 0$. Start with initial guesses $x_{-1} = 0$ and $x_0 = 1$, do 3 iterations

Sol:

Let $f(x) = e^{-x} - x$

$x_{i-1} = 0$

$x_i = 1$

$$x_{i+1} = x_i - \frac{f(x_i)(x_{i-1} - x_i)}{f(x_{i-1}) - f(x_i)}$$

$$x_{i+1} = x_i - \frac{(e^{-x_i} - x_i)(x_{i-1} - x_i)}{(e^{-x_{i-1}} - x_{i-1}) - (e^{-x_i} - x_i)}$$

$x_{i-1} = x$
$x_i = y$ لاستخدام الآلة

$$x_{i+1} = y - \frac{(e^{-y} - y)(x - y)}{(e^{-x} - x) - (e^{-y} - y)}$$

1st iteration

$x = x_{i-1} = 0$

$y = x_i = 1$

$x_{i+1} = 0.6126998$

KAMEL ELFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

2nd iteration

$x = x_{i-1} = 1$

$y = x_i \cong 0.6126998$

$x_{i+1} \cong 0.5638384$

3rd iteration

$x = x_{i-1} = 0.6126998$

$y = x_i = 0.5638384$

$x_{i+1} \cong 0.5671704$

The answer

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

4 Modified - Newton Raphson Method:

Introduction:



Multiplicity of the root = 1
(simple root)



Multiplicity = 2



Multiplicity = 3



Multiplicity = 4

Def: If $f(x)$ is a polynomial and $f(a) = 0$ then the ~~root~~ multiplicity of the root a is (n) if

$$f(x) = (x-a)^n g(x) \quad \text{such that } g(a) \neq 0$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

If a is a root with odd multiplicity then the curve $f(x)$ intersects the x-axis (crosses) at $x = a$.

But if a with even multiplicity then f touches the x-axis at $x = a$.

has odd. multiplicity



has even multiplicity.

e.g.: If $f(x) = (x-a)^2 g(x)$ where $g(a) \neq 0$ (the multiplicity of a is 2) and g is twice differentiable, find $f(a)$, $f'(a)$, $f''(a)$.

Sol: $f(a) = 0$

$$f'(x) = (x-a)^2 g'(x) + 2(x-a) g(x)$$

$$f'(a) = 0$$

$$f''(x) = (x-a)^2 g''(x) + 2(x-a) g'(x) + 2(x-a) g'(x) + 2 g(x)$$

$$f''(a) = 2g(a) \neq 0$$

KAMEL FLEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

In genal If f is a polynomial,

a is root of $f(x) = 0$

then the multiplicity of (a) is (n) iff

$$f(a) = f'(a) = f''(a) = \cdots = f^{(n-1)}(a) = 0 \text{ but}$$

$$f^{(n)}(a) \neq 0 .$$

e.g: Find the roots of $f(x) = (x^2-1)^2(x^2-x)^3$. and determine the multiplicity of each.

Sol:

$$f'(x) = \left((x-1)(x+1)\right)^2 \left(x(x-1)\right)^3$$

$$= (x-1)^2 (x+1)^2 x^3 (x-1)^3$$

$$= x^3 (x+1)^2 (x-1)^5$$

The roots are:

0 and its multiplicity = 3

-1 " " " = 2

1 " " = 5

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Given that 1 is a root of $x^4 - 2x^3 + 2x - 1 = 0$

find its multiplicity.

Sol:

Let $f(x) = x^4 - 2x^3 + 2x - 1$

then $f(1) = f'(1) = f''(1) = 0$

but $f'''(1) = 12 \neq 0$ (check that)

So the multiplicity of 1 is (3).

Recal (Newton - Raphson Method)

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$



Modified Newton - Raphson Method

$$x_{i+1} = x_i - \frac{f(x_i)\, f'(x_i)}{(f'(x_i))^2 - f(x_i)\, f''(x_i)}$$

x_0 = initial guess.

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Using Modified Newton-Raphson method approximate the root of the equation: $x^3 - 4x^2 + 5x - 2 = 0$

take $x_0 = 0.5$ (initial guess).

Do 3 iterations (i.e find x_3) and round off your answer to 6 d.p's.

Sol

Let $f(x) = x^3 - 4x^2 + 5x - 2$

$f'(x) = 3x^2 - 8x + 5$

$f''(x) = 6x - 8$

$$x_{i+1} = x_i - \frac{f'(x_i)\, f(x_i)}{(f'(x_i))^2 - f(x_i) f''(x_i)}$$

$$x_{i+1} = x_i - \frac{(3x_i^2 - 8x_i + 5)(x_i^3 - 4x_i^2 + 5x_i - 2)}{(3x_i^2 - 8x_i + 5)^2 - (x_i^3 - 4x_i^2 + 5x_i - 2)(6x_i - 8)}$$

$x_0 = 0.5$

$x_1 = 1.052631579$

$x_2 = 1.001540832$

$x_3 = 1.000001191$

$x_3 \cong 1.000001$ to 6 d.p's.

- END -

KAMEL FEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

Muller Method:

Aim: $f(x) = 0$

x_0, x_1, x_2 are guesses want to find x_3

Introduction.

To solve: $ax^2 + bx + c = 0$

we can use $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

or we could use ??

$$x = \frac{-2c}{b \pm \sqrt{b^2 - 4ac}}$$

Check: $x^2 - 2x - 3 = 0$

$a = 1, b = -2, c = -3$

$$x = \frac{-2(-3)}{-2 \pm \sqrt{4 - 4(1)(-3)}}$$

$$= \frac{6}{-2 \pm 4}$$

$x_1 = \frac{6}{-2+4}$, $x_2 = \frac{6}{-2-4}$

$x_1 = 3$, $x_2 = -1$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$f(x) = 0$

x_0, x_1, x_2 are initial guesses



Let $\boxed{q(x) = a(x-x_2)^2 + b(x-x_2) + c}$

such that

$q(x_0) = f(x_0)$

$q(x_1) = f(x_1)$

$q(x_2) = f(x_2)$

Now $q(x_2) = f(x_2) \longrightarrow \boxed{c = f(x_2)}$

$q(x_0) = f(x_0) \longrightarrow \boxed{a(x_0-x_2)^2 + b(x_0-x_2) + c = f(x_0)}$

$q(x_1) = f(x_1) \longrightarrow \boxed{a(x_1-x_2)^2 + b(x_1-x_2) + c = f(x_1)}$

So we can find a, b and c

Now solve $a(x-x_2)^2 + b(x-x_2) + c = 0$

$$\Rightarrow x - x_2 = \frac{-2c}{b \pm \sqrt{b^2 - 4ac}}$$

$$x = x_2 - \frac{2c}{b \pm \sqrt{b^2 - 4ac}}$$

$$\Rightarrow \boxed{x_3 = x_2 - \frac{2c}{b + sgn(b)\sqrt{b^2 - 4ac}}}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Also we can find a, b and c using the following formulas

$$h_0 = x_1 - x_0 \qquad h_1 = x_2 - x_1$$

$$\delta_0 = \frac{f(x_1) - f(x_0)}{x_1 - x_0} \qquad \delta_1 = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

$$a = \frac{\delta_1 - \delta_0}{h_1 + h_0}, \quad b = a h_1 + \delta_1, \quad c = f(x_2)$$

then use

$$x_3 = x_2 - \frac{2c}{b + \operatorname{sgn}(b)\sqrt{b^2 - 4ac}}$$

First iteration

$x_0, x_1, x_2 \longrightarrow x_3$

2nd iteration

$x_0, x_1, x_2 \longrightarrow x_3$

3rd iteration

$x_0 \quad x_1 \quad x_2 \longrightarrow x_3$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g (From the Text)

Use Muller's method

to solve $x^3 - 13x - 12 = 0$ take $x_0 = 4.5$,

$x_1 = 5.5$, $x_2 = 5$

Do 2 iterations.

Solution

Let $\boxed{f(x) = x^3 - 13x - 12}$.

$x_0 = 4.5 \longrightarrow f(x_0) = 20.625$

$x_1 = 5.5 \longrightarrow f(x_1) = 82.875$

$x_2 = 5 \longrightarrow f(x_2) = 48$

Let $\boxed{q(x) = a(x-5)^2 + b(x-5) + c}$

$f(x_2) = q(x_2) \Longrightarrow c = f(x_2) \longrightarrow \boxed{c = 48}$

$f(x_0) = q(x_0) \Longrightarrow 0.25a + 0.5b + c = 20.625$

$\boxed{0.25a - 0.5b = 27.375}$ — ①

$f(x_1) = q(x_1) \Longrightarrow 0.25a + 0.5b + c = 82.875$

$\Longrightarrow \boxed{0.25a + 0.5b = 34.875}$ — ②

use calculator or any method else,

find a, b

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$\Rightarrow$ $a = 15$, $b = 62.25$, $c = 48$

$$x_3 = x_2 - \frac{2c}{b + \text{sgn}(b)\sqrt{b^2 - 4ac}}$$

$$x_3 = 5 - \frac{2(48)}{62.25 + \sqrt{62.25^2 - 4(15)(48)}}$$

$$x_3 = 3.976487042$$

$$x_3 \tilde{=} 3.976487$$

2nd iteration

$x_0 = 5.5 \longrightarrow f(x_0) = 82.875$

$x_1 = 5 \longrightarrow f(x_1) = 48$

$x_2 = 3.976487 \Longrightarrow f(x_2) = -0.816334$

$\vdots$

$x_3 \tilde{=} 4.00105$

- END -

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Solving systems of Non-linear Equations:

(by fixed point-iteration approach):

$u(x,y) = 0$
$v(x,y) = 0$
] 2-equations with 2 unknows | initial guesses x_0, y_0

$u(x,y) = 0 \longrightarrow x = g_1(x,y)$

$v(x,y) = 0 \longrightarrow y = g_2(x,y)$

1st iteration: x_0, y_0

$x_1 = g_1(x_0, y_0)$

$y_1 = g_2(x_1, y_0)$

2nd iteration

$x_2 = g_1(x_1, y_1)$

$y_2 = g_2(x_2, y_1)$

3rd iteration

$x_3 = g_1(x_2, y_2)$

$y_3 = g_2(x_3, y_2)$

⋮

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Solve the non-linear system

$$x^2 + xy = 10$$

$$y + 3xy^2 = 57$$

by using

$$x = \sqrt{10 - xy}$$

$$y = \sqrt{\frac{57 - y}{3x}}$$

take $x_0 = 1.5$ and $y_0 = 3.5$

Do 3 iterations give your answer to 5 d.p's.

$$x = \sqrt{10 - xy} \leftarrow g_1(x,y)$$

$$y = \sqrt{\frac{57 - y}{3x}} \leftarrow g_2(x,y)$$

1st iteration:

$x_0 = 1.5$, $y_0 = 3.5$

$$x_1 = \sqrt{10 - (1.5)(3.5)} = 2.179449$$

$$y_1 = \sqrt{\frac{57 - 3.5}{3(2.179449)}} = 2.860506$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

2nd iteration

$x_1 = 2.179449$, $y_1 = 2.860506$

$$x_2 = \sqrt{10 - (2.179449)(2.860506)}$$

$$x_2 = 1.940534$$

$$y_2 = \sqrt{\frac{57 - 2.860506}{3(1.940534)}}$$

$$y_2 = 3.049551$$

3rd iteration

$x_2 = 1.940534$, $y_2 = 3.049551$

$$x_3 = \sqrt{10 - (1.940534)(3.049551)}$$

$$x_3 = 2.020456$$

$$y_3 = \sqrt{\frac{57 - 3.049551}{3(2.020456)}}$$

$$y_3 = 2.983405$$

$\therefore x \cong 2.02046$, $y = 2.98340$ to 5 d.p's

→ Given that the true solution $x = 2$, $y = 3$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Approximate the solution of simultaneous system of non-linear equations

$$y = x^2 + 1$$
$$y = 2\cos x$$

by using fixed-point method use the equations

$$x = \sqrt{y-1} \quad , y = 2\cos x$$

taking $x_0 = 1$ and $y_0 = 1.5$. Do 2 iterations

Sol:

$$x = \sqrt{y-1}$$
$$y = 2\cos x$$

Note
set up your calculator to radian mod

1st iteration:

$x_0 = 1 \quad , y_0 = 1.5$

$$x_1 = \sqrt{1.5-1} = 0.707167812$$

$$y_1 = 2\cos(0.707167812) = 1.520489194$$

KAMEL ELEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2nd iteration:

$x_1 = 0.707167812$

$y_1 = 1.520489194$

$x_2 = \sqrt{1.520489194 - 1} = 0.721449704$

$y_2 = 2\cos(0.721449704)$

$y_2 = 1.501698494$

$\therefore x \cong 0.721449704$

$y \cong 1.501698494$

END

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Matrices and determinants:

Notation $A = [A] \longleftrightarrow$ matrix A

$$A = \begin{bmatrix} 2 & -1 & 3 \\ 5 & 0 & 7 \end{bmatrix} = \begin{pmatrix} 2 & -1 & 3 \\ 5 & 0 & 7 \end{pmatrix}$$



$a_{11} = 2$ $\quad a_{12} = -1$ $\quad a_{13} = 3$

$a_{21} = 5$ $\quad a_{22} = 0$ $\quad a_{23} = 7$

entries (elements) of the matrix A

$$B = \begin{bmatrix} 5 & 8 \\ 4 & -2 \\ 0 & 10 \end{bmatrix} \longrightarrow 3 \times 2 \text{ - matrix}$$

$b_{12} = 8$, $b_{32} = 10$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Def: If A and B are two matrices then $A = B$ if

1) A and B have the same size

2) $a_{ij} = b_{ij}$ for any i and j.

e.g: Given that

$$\begin{bmatrix} x^2 & 5 \\ 7 & 2Z \end{bmatrix} = \begin{bmatrix} 25 & y \\ 7 & -2 \end{bmatrix}$$

find the values of x, y and Z ?

$x^2 = 25 \longrightarrow x = \pm 5$

$5 = y \longrightarrow y = 5$

$2Z = -2 \longrightarrow Z = -1$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

If $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1m} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2m} \\ \vdots & \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nm} \end{bmatrix}$ is

$n \times m$ matrix then A could be written as

$$A = [a_{ij}] \qquad 1 \le i \le n \;,\; 1 \le j \le m$$

and A is called a square matrix if $n = m$, otherwise it is called a rectangular matrix

Notice that

If A is a square matrix

$$A = \begin{bmatrix} a_{11} & & & & \\ & a_{22} & & & \\ & & \ddots & & \\ & & & & a_{nn} \end{bmatrix}$$

$i < j$ (above the diagonal), $i > j$ (below the diagonal), $i = j$ (the diagonal)

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$i > j$ $\quad$ $i = j$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & a_{3n} \\ \vdots & & & \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix}$$

→ upper triangular matrix

all the elements are zeros

upper triangular matrix ≡ square matrix such that all the elements below the diagonal are Zeros

For any $i > j$, $a_{ij} = 0$

---

all elements are zeros

$$A = \begin{bmatrix} a_{11} & 0 & 0 & \cdots & 0 \\ a_{21} & a_{22} & 0 & \cdots & 0 \\ \vdots & & \ddots & & \vdots \\ a_{n1} & a_{n2} & \cdots & & a_{nn} \end{bmatrix}$$

→ Lower triangular matrix

Lower triangular matrix ≡ square matrix such that all elements above the diagonal are Zeros.

For any $i < j$, $a_{ij} = 0$

KAMEL ELEIEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$\begin{bmatrix} 1 & 2 & 5 \\ 0 & -1 & 3 \\ 0 & 0 & 7 \end{bmatrix}, \quad \begin{bmatrix} 4 & 8 \\ 0 & -10 \end{bmatrix}, \quad \begin{bmatrix} 2 & 3 & 1 & 8 \\ 0 & 0 & -1 & 3 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & -2 \end{bmatrix}$$

are upper triangular matrices

$$\begin{bmatrix} 2 & 0 & 0 \\ 5 & -3 & 0 \\ 0 & -1 & 8 \end{bmatrix}, \quad \begin{bmatrix} 5 & 0 \\ -1 & 6 \end{bmatrix}, \quad \begin{bmatrix} 4 & 0 & 0 & 0 \\ -1 & 10 & 0 & 0 \\ -4 & 5 & 8 & 0 \\ 0 & -7 & 3 & 0 \end{bmatrix}$$

are lower triangular matrix

Any matrix (with any size) such that all its elements are Zeros is called a Zero matrix

$$0 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & 0 & 0 \end{bmatrix}$$

are Zero matrices.

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Identity (unit) matrix (square matrix)

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

$$I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \ldots$$ are identity (unit) matrices

$I_n = I \cong$ square matrix such that all the elements on the diagonal are 1's, and all the elements outside the diagonal are 0's.

KAMEL FLEIFEL

Def:

If $A = [a_{ij}]$ is $n \times m$-matrix then the transpose of A is $A^T = [a_{ji}]$ ($m \times n$-matrix)

Def: A matrix is called symmetric if $A^T = A$.

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Find the transpose of each matrix then determine whether the matrix is symmetric.

1) $A = \begin{bmatrix} 2 & -1 & 3 \\ 4 & 0 & 8 \end{bmatrix}$ 2×3

$\rightarrow A^T = \begin{bmatrix} 2 & 4 \\ -1 & 0 \\ 3 & 8 \end{bmatrix}$ 3×2

2) $B = \begin{bmatrix} 4 & 5 \\ 1 & -2 \end{bmatrix}$ 2×2

$B^T = \begin{bmatrix} 4 & 1 \\ 5 & -2 \end{bmatrix}$ 2×2 $\quad B^T \neq B$

$\rightarrow$ B is not symmetric

3) $C = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}$ 3×1

$C^T = \begin{bmatrix} 2 & -1 & 4 \end{bmatrix}$ 1×3

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

4) $A = \begin{bmatrix} 2 & 3 & 1 \\ 3 & 7 & -5 \\ 1 & -5 & 0 \end{bmatrix}$ 3×3

$$A^T = \begin{bmatrix} 2 & 3 & 1 \\ 3 & 7 & -5 \\ 1 & -5 & 0 \end{bmatrix}$$

$\Rightarrow A^T = A \rightarrow A$ is symmetric

5) $B = \begin{bmatrix} 4 & -1 \\ -1 & 5 \end{bmatrix}$

$$B^T = \begin{bmatrix} 4 & -1 \\ -1 & 5 \end{bmatrix}$$

$\rightarrow B^T = B \rightarrow B$ is symmetric

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Operation on Matrices:

Def: If $A=[a_{ij}]$, $B=[b_{ij}]$ are two $n\times m$-matrices then

1) $A \mp B = [a_{ij} \mp b_{ij}]$ $n\times m$-matrix

2) $kA = [ka_{ij}]$ $n\times m$-matrix k: scalar $k \in \mathbb{R}$

e.g: Find if possible

1) $\begin{bmatrix} 2 & -1 & 3 \\ 0 & 5 & 8 \end{bmatrix} + \begin{bmatrix} -7 & 5 & 1 \\ 3 & 2 & 1 \end{bmatrix} = \begin{bmatrix} -5 & 4 & 4 \\ 3 & 7 & 9 \end{bmatrix}$

2×3 — 2×3

2) $\begin{bmatrix} 2 \\ 1 \end{bmatrix} + \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix}$ (undefined)

2×1 3×1

3) $[-1 \quad 3] - [1 \quad 5] = [-2 \quad -2]$

1×2 1×2

4) $5\begin{bmatrix} -1 & 2 \\ 1 & -3 \end{bmatrix} = \begin{bmatrix} -5 & 10 \\ 5 & -15 \end{bmatrix}$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

5) $2\begin{bmatrix} -1 & 4 \\ 5 & 3 \end{bmatrix} + 7\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$= \begin{bmatrix} -2 & 8 \\ 10 & 6 \end{bmatrix} + \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$

$= \begin{bmatrix} 5 & 8 \\ 10 & 13 \end{bmatrix}$

Matrix - Multiplication:

If A is $n \times m$ - matrix,

B is $m \times p$ - matrix then

$AB = C$ is $n \times p$ - matrix.

$n \times m \quad m \times p$

where $c_{ij} = \sum_{k=1}^{m} a_{ik} b_{kj}$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Find if possible

1) $\begin{bmatrix} 2 & 1 & -1 \\ 3 & 0 & 2 \end{bmatrix} \begin{bmatrix} 5 & 1 \\ 0 & -3 \\ 4 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} \square & \square \\ \square & \square \end{bmatrix}$

$2\times 3 \qquad 3\times 2$ ✓

2×2-matrix

$$= \begin{bmatrix} (2)(5)+(1)(0)+(-1)(4) & (2)(1)+(1)(-3)+(-1)(-3) \\ (3)(5)+(0)(0)+(2)(4) & (3)(1)+(0)(-3)+(2)(1) \end{bmatrix}$$

$$= \begin{bmatrix} 6 & 2 \\ 23 & 5 \end{bmatrix}$$

2) $\begin{bmatrix} 5 & 1 \\ 0 & -3 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & -1 \\ 3 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 13 & 5 & -3 \\ -9 & 0 & -6 \\ 11 & 4 & -2 \end{bmatrix}$

$3\times 2 \qquad 2\times 3$ ✓

$AB \neq BA$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Given that $A = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 2 & -3 \\ -2 & 0 \end{bmatrix}$

Find

1) $AB = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}\begin{bmatrix} 2 & -3 \\ -2 & 0 \end{bmatrix} = \begin{bmatrix} -6 & 3 \\ -2 & -9 \end{bmatrix}$

$2\times 2 \quad 2\times 2$

2) $BA = \begin{bmatrix} 2 & -3 \\ -2 & 0 \end{bmatrix}\begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$

$= \begin{bmatrix} -11 & -8 \\ 2 & -4 \end{bmatrix}$

3) $A^2 = AA = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}\begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$

$= \begin{bmatrix} 7 & 6 \\ 9 & 22 \end{bmatrix}$

4) $A^T + B^T = \begin{bmatrix} -1 & 3 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} 2 & -2 \\ -3 & 0 \end{bmatrix}$

$= \begin{bmatrix} 1 & 1 \\ -1 & 4 \end{bmatrix}$ ✓

5) $(A+B)^T$

$A+B = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix} + \begin{bmatrix} 2 & -3 \\ -2 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 1 & 4 \end{bmatrix}$

$\Rightarrow (A+B)^T = \begin{bmatrix} 1 & 1 \\ -1 & 4 \end{bmatrix}$ ✓

$(A+B)^T = A^T + B^T$

KAMEL FLEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

6) $(A-B)^T$

7) $A^T - B^T$

8) $A^T B^T$

9) $B^T A^T$

10) $(AB)^T$

EX

$(AB)^T = B^T A^T$

$(ABC)^T = C^T B^T C^T$

KAMEL FLEIFEL

Answers

6) $\begin{bmatrix} -3 & 5 \\ 5 & 4 \end{bmatrix}$ 7) $\begin{bmatrix} -3 & 5 \\ 5 & 4 \end{bmatrix}$ 8) $\begin{bmatrix} -11 & 2 \\ -8 & -4 \end{bmatrix}$ 9) $\begin{bmatrix} -6 & -2 \\ 3 & -9 \end{bmatrix}$

10) $\begin{bmatrix} -6 & -2 \\ 3 & -9 \end{bmatrix}$

In general $(AB)^T = B^T A^T$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Determinants: (For square matrices)

Def: (For 2×2-matrix)

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then the determinant of A is denoted by

$$\det(A) = |A| = ad - bc$$

Notes For any square matrix (of any size)

* $\det(A) = 0 \equiv$ A is singular
  $\equiv$ A has no inverse (A^{-1})
  $\equiv$ A is not invertible

* $\det(A) \neq 0 \equiv$ A is not singular.
  $\equiv$ A has inverse (A^{-1})
  $\equiv$ A is invertible

---

* $IA = AI = A$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Find the determinante of the given matrix

1) $A = \begin{bmatrix} 2 & -1 \\ 3 & 5 \end{bmatrix} \longrightarrow .$

2) $B = \begin{bmatrix} 2 & 4 \\ 6 & 12 \end{bmatrix}$

3) $C = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$

sol: 1) $\det(A) = (2)(5) - (-1)(3) = 10 + 3 = 13$

2) $\det(B) = (2)(12) - (4)(6)$
$= 24 - 24 = 0$

3) $\det(C) = \cos^2\theta - (-\sin^2\theta)$
$= \cos^2\theta + \sin^2\theta$
$= 1 .$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

A is 3x3

$$\begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix}$$

$(-1)^{i+j}$

e.g: If $A = \begin{bmatrix} 2 & 3 & -1 \\ 0 & 1 & 2 \\ 4 & -2 & 1 \end{bmatrix}$ find determinante of A. (det(A) or |A|).

$$\det(A) = \begin{vmatrix} 2 & 3 & -1 \\ 0 & 1 & 2 \\ 4 & -2 & 1 \end{vmatrix}$$

$$= +(2)\begin{vmatrix} 1 & 2 \\ -2 & 1 \end{vmatrix} - (3)\begin{vmatrix} 0 & 2 \\ 4 & 1 \end{vmatrix} + (-1)\begin{vmatrix} 0 & 1 \\ 4 & -2 \end{vmatrix}$$

$$= 2(1+4) - 3(0-8) - 1(0-4)$$

$$= 10 + 24 + 4$$

$$= 38$$

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

حل آخر

$$\begin{vmatrix} 2 & 3 & -1 \\ 0 & 1 & 2 \\ 4 & -2 & 1 \end{vmatrix}$$

$$= +2(1+4) - 0(3-2) + 4(6+1)$$

$$= 10 - 0 + 28$$

$$= 38$$

- END -

KAMEL FLEIFEL

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Linear System of Equations

Def: The following set of linear equations

$$a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1$$
$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2$$
$$\vdots$$
$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m$$

m - equation
n - variables

is called Linear System of Equations

The above system could be written (using matrices) in two ways

① $$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & & & \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

$m \times n$ $\quad n \times 1$ $\quad m \times 1$

Simply $\boxed{AX = B}$

A: matrix of coefficients
X: matrix of variables (unknowns)
B: matrix of constants

② Augmented matrix $[A : B]$

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & & & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array}\right]$$

KAMEL FLEFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Write the following linear system of equations, using matrices in the two different ways.

$$2x_1 - x_2 + 3x_3 = 5$$
$$x_1 + x_2 + x_4 = -1$$
$$7x_1 + x_3 + x_4 = 8$$

Sol:

1) $AX = B$

$$\begin{bmatrix} 2 & -1 & 3 & 0 \\ 1 & 1 & 0 & 1 \\ 7 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 5 \\ -1 \\ 8 \end{bmatrix}$$

A: 3×4 — X: 4×1 — B: 3×1

2) $[A \,;\, B] \Rightarrow$

$$\left[\begin{array}{cccc|c} 2 & -1 & 3 & 0 & 5 \\ 1 & 1 & 0 & 1 & -1 \\ 7 & 0 & 1 & 1 & 8 \end{array}\right]$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Linear system



Cramer's Rules

$AX = B$

A: square matrix.

$\det(A) \neq 0$

e.g: Use Cramer's rule to solve the linear system?

$$x + 2y = 5$$
$$3x + 2y = 7.$$

Sol:

$AX = B \longrightarrow$

$$\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

A X B.

$\det(A) = (1)(2) - (2)(3)$

$= 2 - 6 = -4 \neq 0$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\det(A_x) = \begin{vmatrix} 5 & 2 \\ 7 & 2 \end{vmatrix} = 10 - 14 = -4$$

$$x = \frac{\det(A_x)}{\det(A)} = \frac{-4}{-4} = 1 \rightarrow \boxed{x = 1}$$

$$\det(A_y) = \begin{vmatrix} 1 & 5 \\ 3 & 7 \end{vmatrix} = 7 - 15 = -8$$

$$\therefore y = \frac{\det(A_y)}{\det(A)} = \frac{-8}{-4} = 2 \rightarrow \boxed{y = 2}$$

$$\therefore X = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \qquad X = (1, 2)$$

e.g: Using Cramer's rule, solve

$$x + y + z = 6$$
$$2x - y + z = 3$$
$$y + z = 5$$

Sol:

$$AX = B \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 2 & -1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix}$$

A X B

$$\det(A) = +0(1 - (-1)) - (1)(1 - 2) + (1)(-1 - 2)$$
$$= 0 + 1 - 3 = -2 \neq 0$$

$$\det(A) = -2 \neq 0$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\det(A_x) = \begin{vmatrix} 6 & -1 & 1 \\ 3 & -1 & 1 \\ 5 & 1 & 1 \end{vmatrix}$$

$$\begin{matrix} + & - & + \\ - & + & - \\ + & - & + \end{matrix}$$

5

$$= +(6)(-1-1) - (1)(3-5) + 1(3-(-5))$$

$$= -12 + 2 + 8$$

$$= -2$$

$$\therefore x = \frac{\det(A_x)}{\det(A)} = \frac{-2}{-2} = 1$$

$$x = 1$$

$$\det(A_y) = \begin{vmatrix} 1 & 6 & 1 \\ 2 & 3 & 1 \\ 0 & 5 & 1 \end{vmatrix} =$$

$$= +1(3-5) - (2)(6-5) + 0$$

$$= -2 - 2 = -4$$

$$\therefore y = \frac{\det(A_y)}{\det(A)} = \frac{-4}{-2} = 2 \rightarrow y = 2$$

$$\det(A_z) = \begin{vmatrix} 1 & 1 & 6 \\ 2 & -1 & 3 \\ 0 & 1 & 5 \end{vmatrix}$$

$$= 0 - 1(3-12) + 5(-1-2)$$

$$= 9 - 15 = -6$$

$$\therefore z = \frac{\det(A_z)}{\det(A)} = \frac{-6}{-2} = 3$$

$$\therefore X = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Inverse method:

A is n×n-matrix.

$$\det(A) \neq 0 \longrightarrow \exists A^{-1} \quad (A^{-1}A = I)$$

$$AX = B \longrightarrow A^{-1}AX = A^{-1}B$$

$$\longrightarrow IX = A^{-1}B$$

$$\therefore \boxed{X = A^{-1}B}$$

e.g.: Use the inverse method to solve, the linear system

$$x + 2y = 5$$
$$3x + 2y = 7$$

sol:

$$AX = B \longrightarrow \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$
$$A \quad X \quad B$$

$$\boxed{X = A^{-1}B}$$

$$\det(A) = 2 - 6 = -4 \neq 0$$

$$A^{-1} = \frac{1}{-4}\begin{bmatrix} 2 & -2 \\ -3 & 1 \end{bmatrix} \rightarrow A^{-1} = \begin{bmatrix} \frac{-2}{4} & \frac{-2}{-4} \\ \frac{+3}{4} & \frac{-1}{4} \end{bmatrix}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\therefore A^{-1} = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ +\frac{3}{4} & -\frac{1}{4} \end{bmatrix}$$

$$\therefore X = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ +\frac{3}{4} & -\frac{1}{4} \end{bmatrix} \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

$2 \times 2 \qquad 2 \times 1$

$$X = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \rightarrow \begin{array}{l} x = 1 \\ y = 2 \end{array}$$

Gauss - Jordan method

Def: A matrix A is called in the reduced row echelon form (RREF), if it satisfies the following conditions

1) Zero rows (if any) at the bottom.

2) the first non-zero element in any non-zero row is 1 (called leading 1)

3) For any two non-zero rows, the leading 1 of the lower row to the right of the leading 1 of the upper one.

4) the elements in the column of any leading 1 are zeros (except the leading 1)

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

If A satisfies ①, ② and ③ only it is called in row echelon form (REF)

e.g: which of the following in RREF, REF or neither.

1) $\begin{bmatrix} 1 & 0 & 1 & 0 & 3 \\ 0 & 1 & 0 & 0 & 3 \\ 0 & 0 & 0 & 1 & 5 \end{bmatrix}$ RREF.

2) $\begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 1 & -1 & 2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ REF but not in RREF

3) $\begin{bmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ Not REF
Not in RREF

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Elementary row operations

1. Multiplying any row by non-zero constant
2. Multiplying a row by a constant and adding the results to another row.
3. interchanging any two rows.

e.g: By Gauss-Jordan method, solve the linear system

$$x + y + z = 6$$
$$2x - y + z = 3$$
$$y + z = 5$$

Sol:

$$[A \mid B] \xrightarrow[\text{row operations}]{\text{elementary}} \text{RREF}$$

$[A \mid B]$

$$\begin{bmatrix} \textcircled{1} & 1 & 1 & 6 \\ 2 & -1 & 1 & 3 \\ 0 & 1 & 1 & 5 \end{bmatrix} \sim \begin{bmatrix} \textcircled{1} & 1 & 1 & 6 \\ 0 & -3 & -1 & -9 \\ 0 & 1 & 1 & 5 \end{bmatrix} -2R_1 + R_2$$

$$\sim \begin{bmatrix} \textcircled{1} & 1 & 1 & 6 \\ 0 & \textcircled{1} & 1 & 5 \\ 0 & -3 & -1 & -9 \end{bmatrix} \begin{matrix} \\ R_2 \\ R_3 \end{matrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & \textcircled{2} & 6 \end{bmatrix} \begin{matrix} -R_2 + R_1 \\ \\ 3R_2 + R_3 \end{matrix}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\sim \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & \textcircled{1} & 3 \end{bmatrix} \frac{1}{2}R_3$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 3 \end{bmatrix} -R_3 + R_2$$

$$\left.\begin{array}{l} x = 1 \\ y = 2 \\ z = 3 \end{array}\right\} \longmapsto X = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}.$$

Part 2

e.g: Solve, using Gauss-Jordan.

$$x + y + z = 6.$$
$$2x + y + z = 7$$
$$3x + 2y + 2z = 5$$

Sol:

$$\begin{bmatrix} 1 & 1 & 1 & 6 \\ 2 & 1 & 1 & 7 \\ 3 & 2 & 2 & 5 \end{bmatrix}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Solve, by Gauss-Jordan method

$$x + y + z = 6$$
$$2x + y + z = 7$$
$$3x + 2y + 2z = 5$$

sol:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 2 & 1 & 1 & 7 \\ 3 & 2 & 2 & 5 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & -1 & -1 & -5 \\ 0 & -1 & -1 & -13 \end{array}\right] \begin{array}{l} \\ -2R_1 + R_2 \\ -3R_1 + R_3 \end{array}$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & 1 & 5 \\ 0 & -1 & -1 & -13 \end{array}\right] -R_2 \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & 0 & -8 \end{array}\right] \begin{array}{l} -R_2 + R_1 \\ \\ 1R_2 + R_3 \end{array}$$

$$0x + 0y + 0z = -8$$
$$0 = -8 \quad \text{contradiction}$$

$\therefore$ The system has no solution

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e:9 : Using Gauss-Jordan, solve the system

$$2x_1 + x_2 + x_3 - 2x_4 = 4$$
$$x_1 + 2x_2 + 2x_3 - 4x_4 = 5$$
$$3x_1 + 3x_2 + 3x_3 - 6x_4 = 9$$

sol:

$$\left[\begin{array}{cccc|c} 2 & 1 & 1 & -2 & 4 \\ 1 & 2 & 2 & -4 & 5 \\ 3 & 3 & 3 & -6 & 9 \end{array}\right]$$

$$\sim \left[\begin{array}{cccc|c} \textcircled{1} & 2 & 2 & -4 & 5 \\ 2 & 1 & 1 & -2 & 4 \\ 3 & 3 & 3 & -6 & 9 \end{array}\right] \begin{matrix} R_1 \\ R_2 \\ \\ \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & 2 & -4 & 5 \\ 0 & \textcircled{-3} & -3 & 6 & -6 \\ 0 & -3 & -3 & 6 & -6 \end{array}\right] \begin{matrix} \\ -2R_1 + R_2 \\ -3R_1 + R_3 \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & 2 & -4 & 5 \\ 0 & \textcircled{1} & 1 & -2 & 2 \\ 0 & -3 & -3 & 6 & -6 \end{array}\right] \begin{matrix} \\ -\frac{1}{3}R_2 \\ \\ \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & -2 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right] \begin{matrix} -2R_2 + R_1 \\ \\ 3R_2 + R_3 \end{matrix}$$

→ Inifinitely many solutions

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$x_1 = 1$

$x_2 + x_3 - 2x_4 = 2$

Let $x_4 = r$, $r \in R$

$x_3 = s$ $s \in R$

$x_2 + s - 2r = 2$

$\rightarrow x_2 = 2 + 2r - s$

$$\therefore X = \begin{bmatrix} 1 \\ 2+2r-s \\ s \\ r \end{bmatrix} \qquad r, s \in R$$

$$\left.\begin{matrix} r = 0 \\ s = 1 \end{matrix}\right\} \longrightarrow X = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$$

$$\left.\begin{matrix} r = 1 \\ s = 1 \end{matrix}\right] \longrightarrow X = \begin{bmatrix} 1 \\ 3 \\ 1 \\ 1 \end{bmatrix} \begin{matrix} \rightarrow x_1 \\ \rightarrow x_2 \\ \rightarrow x_3 \\ \rightarrow x_4 \end{matrix}$$

$\vdots$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Gauss- Elimination Method

$AX = B$

$[A : B] \longrightarrow REF$

e.g: Using Gauss elimination method, solve

$$x + y + z = 4$$
$$3x + y + 2z = 8$$
$$x - y + 5z = 10$$

sol:

$$\begin{bmatrix} \textcircled{1} & 1 & 1 & 4 \\ 3 & 1 & 2 & 8 \\ 1 & -1 & 5 & 10 \end{bmatrix}$$ Pivot

$$\sim \begin{bmatrix} 1 & 1 & 1 & 4 \\ 0 & \textcircled{-2} & -1 & -4 \\ 0 & -2 & 4 & 6 \end{bmatrix} \begin{matrix} \\ -3R_1 + R_2 \\ -R_1 + R_3 \end{matrix}$$

$\frac{-(-2)}{-2} = -1$ ← Pivot

$$\sim \begin{bmatrix} 1 & 1 & 1 & 4 \\ 0 & -2 & -1 & -4 \\ 0 & 0 & 5 & 10 \end{bmatrix} \begin{matrix} \\ \\ -1R_2 + R_3 \end{matrix}$$

$5z = 10 \rightarrow \boxed{z = 2}$

$-2y - z = -4 \rightarrow -2y - 2 = -4 \rightarrow -2y = -2 \rightarrow \boxed{y = 1}$

$x + y + z = 4$

$x + 1 + 2 = 4 \rightarrow x = 1$

$\therefore X = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Find all the value(s) of (if any) for which, the system $AX = B$

where $[A:B] = \left[\begin{array}{ccc|c} 1 & 2 & 3 & 5 \\ 0 & 2 & 1 & 4 \\ 0 & 0 & a^2-4 & a-2 \end{array}\right]$

has, a) infinite solutions b) No solution

c) unique solution.

Sol:

$a^2 - 4 = 0 \longrightarrow a = \pm 2$

$a - 2 = 0 \longrightarrow a = 2$

a) $a = 2$

b) $a = -2$

c) $a \in \mathbb{R} \setminus \{-2, 2\}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Solve, by Gauss-Jordan method

$$x + y + z = 6$$
$$2x + y + z = 7$$
$$3x + 2y + 2z = 5$$

sol:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 2 & 1 & 1 & 7 \\ 3 & 2 & 2 & 5 \end{array}\right] \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & -1 & -1 & -5 \\ 0 & -1 & -1 & -13 \end{array}\right] \begin{array}{l} \\ -2R_1 + R_2 \\ -3R_1 + R_3 \end{array}$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & 1 & 5 \\ 0 & -1 & -1 & -13 \end{array}\right] -R_2 \sim \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 5 \\ 0 & 0 & 0 & -8 \end{array}\right] \begin{array}{l} -R_2 + R_1 \\ \\ 1R_2 + R_3 \end{array}$$

$$0x + 0y + 0z = -8$$
$$0 = -8 \quad \text{contradiction}$$

$\therefore$ The system has no solution

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e:9 : Using Gauss-Jordan, solve the system

$$2x_1 + x_2 + x_3 - 2x_4 = 4$$
$$x_1 + 2x_2 + 2x_3 - 4x_4 = 5$$
$$3x_1 + 3x_2 + 3x_3 - 6x_4 = 9$$

sol:

$$\left[\begin{array}{cccc|c} 2 & 1 & 1 & -2 & 4 \\ 1 & 2 & 2 & -4 & 5 \\ 3 & 3 & 3 & -6 & 9 \end{array}\right]$$

$$\sim \left[\begin{array}{cccc|c} \textcircled{1} & 2 & 2 & -4 & 5 \\ 2 & 1 & 1 & -2 & 4 \\ 3 & 3 & 3 & -6 & 9 \end{array}\right] \begin{matrix} R_1 \\ R_2 \\ \\ \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & 2 & -4 & 5 \\ 0 & \textcircled{-3} & -3 & 6 & -6 \\ 0 & -3 & -3 & 6 & -6 \end{array}\right] \begin{matrix} \\ -2R_1 + R_2 \\ -3R_1 + R_3 \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 2 & 2 & -4 & 5 \\ 0 & \textcircled{1} & 1 & -2 & 2 \\ 0 & -3 & -3 & 6 & -6 \end{array}\right] \begin{matrix} \\ -\frac{1}{3}R_2 \\ \\ \end{matrix}$$

$$\sim \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & -2 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right] \begin{matrix} -2R_2 + R_1 \\ \\ 3R_2 + R_3 \end{matrix}$$

→ Inifinitely many solutions

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$x_1 = 1$

$x_2 + x_3 - 2x_4 = 2$

Let $x_4 = r$, $r \in R$

$x_3 = s$ $s \in R$

$x_2 + s - 2r = 2$

$\rightarrow x_2 = 2 + 2r - s$

$$\therefore X = \begin{bmatrix} 1 \\ 2+2r-s \\ s \\ r \end{bmatrix} \qquad r, s \in R$$

$$\left.\begin{matrix} r = 0 \\ s = 1 \end{matrix}\right\} \longrightarrow X = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$$

$$\left.\begin{matrix} r = 1 \\ s = 1 \end{matrix}\right] \longrightarrow X = \begin{bmatrix} 1 \\ 3 \\ 1 \\ 1 \end{bmatrix} \begin{matrix} \rightarrow x_1 \\ \rightarrow x_2 \\ \rightarrow x_3 \\ \rightarrow x_4 \end{matrix}$$

$\vdots$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Gauss- Elimination Method

$AX = B$

$[A : B] \longrightarrow REF$

e.g: Using Gauss elimination method, solve

$$x + y + z = 4$$
$$3x + y + 2z = 8$$
$$x - y + 5z = 10$$

sol:

$$\begin{bmatrix} \textcircled{1} & 1 & 1 & 4 \\ 3 & 1 & 2 & 8 \\ 1 & -1 & 5 & 10 \end{bmatrix}$$ Pivot

$$\sim \begin{bmatrix} 1 & 1 & 1 & 4 \\ 0 & \textcircled{-2} & -1 & -4 \\ 0 & -2 & 4 & 6 \end{bmatrix} \begin{matrix} \\ -3R_1 + R_2 \\ -R_1 + R_3 \end{matrix}$$

$\frac{-(-2)}{-2} = -1$ ← Pivot

$$\sim \begin{bmatrix} 1 & 1 & 1 & 4 \\ 0 & -2 & -1 & -4 \\ 0 & 0 & 5 & 10 \end{bmatrix} \begin{matrix} \\ \\ -1R_2 + R_3 \end{matrix}$$

$5z = 10 \rightarrow \boxed{z = 2}$

$-2y - z = -4 \rightarrow -2y - 2 = -4 \rightarrow -2y = -2 \rightarrow \boxed{y = 1}$

$x + y + z = 4$

$x + 1 + 2 = 4 \rightarrow x = 1$

$\therefore X = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Find all the value(s) of (if any) for which, the system $AX = B$

where $[A:B] = \left[\begin{array}{ccc|c} 1 & 2 & 3 & 5 \\ 0 & 2 & 1 & 4 \\ 0 & 0 & a^2-4 & a-2 \end{array}\right]$

has, a) infinite solutions b) No solution

c) unique solution.

Sol:

$a^2 - 4 = 0 \longrightarrow a = \pm 2$

$a - 2 = 0 \longrightarrow a = 2$

a) $a = 2$

b) $a = -2$

c) $a \in \mathbb{R} \setminus \{-2, 2\}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Finding the inverse of a matrix by elementary row operations

Recall

$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ the multiplicative identity (unit) matrix for 2x2-matrices

$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ — — — — — — — for 3x3-matrices

$I = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ — — — — — — — for 4x4-matrices

⋮

To find A^{-1} by elementary row operations

$[A \mid I]$

↓ RREF

$[I \mid B]$

A has inverse (namely $A^{-1} = B$)

(invertable)

(non-singular)

$\left[\begin{array}{c|c} 0\ 0\ \cdots\ 0 & \cdots \end{array}\right]$

A has no inverse

(not invertable)

(singular)

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

Notice that:

If A has inverse, then

$$A^{-1}A = I = AA^{-1}$$

e.g: Find the inverse of the given matrix (if it exists) by using elementary row operations

1) $A = \begin{bmatrix} 1 & 2 \\ 3 & 5 \end{bmatrix}$

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 5 & 0 & 1 \end{array}\right] \sim \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & -1 & -3 & 1 \end{array}\right] -3R_1 + R_2$$

$$\sim \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & 1 & 3 & -1 \end{array}\right] -R_2$$

$$\sim \left[\begin{array}{cc|cc} 1 & 0 & -5 & 2 \\ 0 & 1 & 3 & -1 \end{array}\right] -2R_2 + R_1$$

$$\therefore A^{-1} = \begin{bmatrix} -5 & 2 \\ 3 & -1 \end{bmatrix}$$

Check: $AA^{-1} = \begin{bmatrix} 1 & 2 \\ 3 & 5 \end{bmatrix}\begin{bmatrix} -5 & 2 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ✓

$2\times 2 \quad 2\times 2$

$A^{-1}A = I$ Ex.

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2) $B = \begin{bmatrix} 1 & 3 \\ 4 & 12 \end{bmatrix}$

$$\left[\begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 4 & 12 & 0 & 1 \end{array}\right] \sim \left[\begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 0 & 0 & -4 & 1 \end{array}\right] -4R_1 + R_2$$

$\therefore$ B has no inverse (not invertable)
(singular)

e.g: Given that $A = \begin{bmatrix} 1 & 1 & -1 \\ 0 & 1 & -2 \\ 2 & 0 & 1 \end{bmatrix}$,

1) find the inverse of A by elementary row operations

2) and hence, solve the system

$$x + y - z = 4$$
$$y - 2z = -2$$
$$2x + z = 10$$

by using inverse method

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

4

$$\left[\begin{array}{ccc|ccc} 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 2 & 0 & 1 & 0 & 0 & 1 \end{array}\right]$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 0 & -2 & 3 & -2 & 0 & 1 \end{array}\right] -2R_1 + R_2$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & -1 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 0 & 0 & -1 & -2 & 2 & 1 \end{array}\right] \begin{array}{l} -R_2 + R_1 \\ \\ 2R_2 + R_3 \end{array}$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & -1 & 0 \\ 0 & 1 & -2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 2 & -2 & -1 \end{array}\right] -R_3$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & 1 & 1 \\ 0 & 1 & 0 & 4 & -3 & -2 \\ 0 & 0 & 1 & 2 & -2 & -1 \end{array}\right] \begin{array}{l} -R_3 + R_1 \\ 2R_3 + R_2 \end{array}$$

$$\therefore A^{-1} = \begin{bmatrix} -1 & 1 & 1 \\ 4 & -3 & -2 \\ 2 & -2 & -1 \end{bmatrix}$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2) $\begin{bmatrix} 1 & 1 & -1 \\ 0 & 1 & -2 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ -2 \\ 10 \end{bmatrix}$

$A \quad X \quad B$

$AX = B \rightarrow X = A^{-1}B$

$\therefore X = \begin{bmatrix} -1 & 1 & 1 \\ 4 & -3 & -2 \\ 2 & -2 & -1 \end{bmatrix} \begin{bmatrix} 4 \\ -2 \\ 10 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \\ 2 \end{bmatrix}$

$3\times3 \quad 3\times1$

$\rightarrow x = 4, \; y = 2, \; z = 2$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Solving linear systems by LU-decomposition method

$$AX = B \longrightarrow A = LU$$

$$Y$$

$$L(UX) = B$$

L: lower triangular matrix.

U: upper triangular matrix.

$$\boxed{LY = B} \quad -(1)$$

$$\boxed{UX = Y} \quad -(2)$$

* Doolittle's method

A 2×2 $\longrightarrow$

$$\begin{bmatrix} 1 & 0 \\ l_{21} & 1 \end{bmatrix}\begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix} = A$$

$$L \qquad U$$

A; 3×3 - matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix}\begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix} = A$$

$$L \qquad U$$

$\vdots$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Solve the following linear system using LU-decomposition method.

$$4x_1 + 5x_2 = 7$$
$$12x_1 + 14x_2 = 18$$

Sol:

$$\begin{bmatrix} 4 & 5 \\ 12 & 14 \end{bmatrix}\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 7 \\ 18 \end{bmatrix}$$

A X B

$AX = B$

$$\begin{bmatrix} 1 & 0 \\ l_{21} & 1 \end{bmatrix}\begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix} = \begin{bmatrix} 4 & 5 \\ 12 & 14 \end{bmatrix}$$

2x2 2x2

$u_{11} = 4$ $u_{12} = 5$

$l_{21}u_{11} = 12 \rightarrow 4l_{21} = 12 \longrightarrow l_{21} = 3$

$l_{21}u_{12} + u_{22} = 14$

$(3)(5) + u_{22} = 14 \longrightarrow u_{22} = -1$

check:

$$\begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}\begin{bmatrix} 4 & 5 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 4 & 5 \\ 12 & 14 \end{bmatrix}$$ ✓

L U A

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$LY = B$ — ①

$$\begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 7 \\ 18 \end{bmatrix}$$

$L \quad Y \quad B$

$y_1 = 7$

$3y_1 + y_2 = 18 \longrightarrow 3(7) + y_2 = 18$

$\rightarrow y_2 = -3$

$\therefore Y = \begin{bmatrix} 7 \\ -3 \end{bmatrix}$

$UX = Y$ — ②

$$\begin{bmatrix} 4 & 5 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 7 \\ -3 \end{bmatrix}$$

$-x_2 = -3 \rightarrow x_2 = 3$

$4x_1 + 5x_2 = 7$

$4x_1 + 5(3) = 7$

$4x_1 + 15 = 7 \longrightarrow 4x_1 = -8 \rightarrow x_1 = -2$

Sol. $X = \begin{bmatrix} -2 \\ 3 \end{bmatrix}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Using LU-decomposition method to solve, the linear system

$$3x_1 + 5x_2 + 2x_3 = 8$$
$$8x_2 + 2x_3 = -7$$
$$6x_1 + 2x_2 + 8x_3 = 26$$

sol:

$$\begin{bmatrix} 3 & 5 & 2 \\ 0 & 8 & 2 \\ 6 & 2 & 8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 8 \\ -7 \\ 26 \end{bmatrix}$$

A X B

$$\begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix} = \begin{bmatrix} 3 & 5 & 2 \\ 0 & 8 & 2 \\ 6 & 2 & 8 \end{bmatrix}$$

L U A

$u_{11} = 3$ $u_{12} = 5$ $u_{13} = 2$

$l_{21} u_{11} = 0 \rightarrow 3 l_{21} = 0 \rightarrow l_{21} = 0$

$l_{12} u_{12} + u_{22} = 8$

$(0)(5) + u_{22} = 8 \rightarrow u_{22} = 8$

$l_{21} u_{13} + u_{23} = 2 \rightarrow (0)(2) + u_{23} = 2$

$\rightarrow u_{23} = 2$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$l_{31} u_{11} = 6 \rightarrow 3 l_{31} = 6 \rightarrow l_{31} = 2$ (10)

$l_{31} u_{12} + l_{32} u_{22} = 2$

$(2)(5) + 8 l_{32} = 2 \rightarrow 8 l_{32} = -8$

$\rightarrow l_{32} = -1$

$l_{31} u_{13} + l_{32} u_{23} + u_{33} = 8$

$(2)(2) + (-1)(2) + u_{33} = 8$

$4 - 2 + u_{33} = 8 \rightarrow u_{33} = 6$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & -1 & 1 \end{bmatrix} \begin{bmatrix} 3 & 5 & 2 \\ 0 & 8 & 2 \\ 0 & 0 & 6 \end{bmatrix} = A$$

check!!

L U

$LY = B \rightarrow$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & -1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 8 \\ -7 \\ 26 \end{bmatrix}$$

L Y B

$y_1 = 8$

$y_2 = -7$

$2y_1 - y_2 + y_3 = 26$

$2(8) - (-7) + y_3 = 26$

$16 + 7 + y_3 = 26 \rightarrow y_3 = 3$

$$\therefore Y = \begin{bmatrix} 8 \\ -7 \\ 3 \end{bmatrix}$$

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

$UX = Y$ (2)

$$\begin{bmatrix} 3 & 5 & 2 \\ 0 & 8 & 2 \\ 0 & 0 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 8 \\ -7 \\ 3 \end{bmatrix}$$

$U \qquad X$

$6x_3 = 3 \rightarrow \boxed{x_3 = \frac{1}{2}}$

$8x_2 + 2x_3 = -7$

$8x_2 + 2(\frac{1}{2}) = -7 \rightarrow 8x_2 = -8$

$\rightarrow \boxed{x_2 = -1}$

$3x_1 + 5x_2 + 2x_3 = 8$

$3x_1 + 5(-1) + 2(\frac{1}{2}) = 8$

$3x_1 - 5 + 1 = 8$

$3x_1 = 12 \rightarrow \boxed{x_1 = 4}$

$\therefore \quad X = \begin{bmatrix} 4 \\ -1 \\ \frac{1}{2} \end{bmatrix}$

END .

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Finding the inverse of a matrix by using LU-decomposition method:

Recall

$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ is the identity for 2x2 - matrices

$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is the identity for 3x3 - matrix.

$AX = I$ $\qquad X = A^{-1}$

$L(\underbrace{UX}_{Y}) = I$

$LY = I$

$UX = Y$

For 2x2 - matrix

First Column of A^{-1}

$LY = e_1$ — ①

$UX = Y$ — ②

$e_1 \quad e_2$

$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$A^{-1} = \begin{bmatrix} x & \\ & \end{bmatrix}$

2nd column of A^{-1}

$LY = e_2$ — ①

$UX = Y$ — ②

$A^{-1} = \begin{bmatrix} x & x \\ & \end{bmatrix}$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

A is

* If 3x3 - matrix

e_1 e_2 e_3

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

1st column

$LY = e_1$ — ①

$UX = Y$ — ②

$$A^{-1} = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$ 3x3 - matrix

2nd column

$LY = e_2$

$UX = Y$

3rd column

$LY = e_3$

$UX = Y$

---

e.g: Using LU-decomposition, find A^{-1} ?

1) $A = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}$

$$\begin{bmatrix} 1 & 0 \\ l_{21} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}$$

$\rightarrow u_{11} = 1$ $u_{12} = 3$

$l_{21} u_{11} = 1 \rightarrow l_{21} = 1$

$l_{21} u_{12} + u_{22} = 2 \rightarrow (1)(3) + u_{22} = 2$

$\rightarrow u_{22} = -1$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

$$\therefore \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}\begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}$$

$L \quad U \quad A$

$e_1 \quad e_2$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} x_1 & x_2 \\ & \end{bmatrix}$$

First column of the inverse

$$LY = e_1 \quad -①$$

$$UX = Y \quad -②$$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$y_1 = 1$

$y_1 + y_2 = 0 \rightarrow y_2 = -1$

$$\therefore Y = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$UX = Y \rightarrow \begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix}\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$-x_2 = -1 \rightarrow x_2 = 1$

$x_1 + 3x_2 = 1$

$x_1 + 3(1) = 1 \rightarrow x_1 = -2$

$$\therefore X = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

2nd ~~row~~ Column

$$A^{-1} = \begin{bmatrix} -2 & \\ 1 & 0 \end{bmatrix}$$

$$LY = e_2$$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$y_1 = 0$

$y_1 + y_2 = 1$

$\rightarrow \quad 0 + y_2 = 1$

$\therefore y_2 = 1$

$\therefore \quad Y = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

$$UX = Y \rightarrow \begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix}\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$-x_2 = 1 \rightarrow x_2 = -1$

$x_1 + 3x_2 = 0$

$x_1 + 3(-1) = 0 \rightarrow x_1 = 3$

$\therefore X = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$

$$\therefore A^{-1} = \begin{bmatrix} -2 & 3 \\ 1 & -1 \end{bmatrix}$$

Check that $AA^{-1} = I = A^{-1}A$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

2) $A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 1 \\ 0 & 2 & 0 \end{bmatrix}$ $\xrightarrow{-1}$

$LU = A$

$$\begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix} = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 1 \\ 0 & 2 & 0 \end{bmatrix} \xrightarrow{-1}$$

$L \qquad U \qquad A$

$\therefore$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & -\frac{1}{3} & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 1 & 2 & 0 \\ 0 & -3 & 1 \\ 0 & 0 & \frac{1}{3} \end{bmatrix}$$

1st column of A^{-1}

$e_1 \quad e_2 \quad e_3$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$LY = e_1 \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & -\frac{1}{3} & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$A^{-1} = [\quad\quad]$

$y_1 = 1$

$2y_1 + y_2 = 0 \rightarrow y_2 = -2$

$-\frac{1}{3} y_2 + y_3 = 0 \rightarrow \frac{2}{3} + y_3 = 0 \rightarrow y_3 = -\frac{2}{3}$

$$\therefore Y = \begin{bmatrix} 1 \\ -2 \\ -\frac{2}{3} \end{bmatrix}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$UX = Y$

$$\begin{bmatrix} 1 & 2 & 0 \\ 0 & -3 & 1 \\ 0 & 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ -\frac{2}{3} \end{bmatrix}$$

$\frac{1}{3}x_3 = \frac{-2}{3} \rightarrow (x_3 = -2)$

$-3x_2 + x_3 = -2$

$-3x_2 - 2 = -2 \rightarrow -3x_2 = 0$

$\rightarrow (x_2 = 0)$

$x_1 + 2x_2 = 1$

$x_1 + 2(0) = 1 \rightarrow (x_1 = 1)$

$\therefore X = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}$

2nd Column ~~row~~ of A^{-1}

$$A^{-1} = \begin{bmatrix} 1 & (0) & \\ 0 & & \\ -2 & & \end{bmatrix}$$

$LY = e_2$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & -\frac{1}{3} & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \rightarrow Y = \begin{bmatrix} 0 \\ 1 \\ \frac{1}{3} \end{bmatrix}$$

$$UX = Y \longrightarrow X = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

3rd ~~row~~ Column of the inverse

$$A^{-1} = \begin{bmatrix} 1 & 0 & \\ 0 & 0 & \\ -2 & 1 & \end{bmatrix}$$

$$LY = e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\longrightarrow \quad Y = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$UX = Y \longrightarrow X = \begin{bmatrix} -2 \\ 1 \\ 3 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 1 & 0 & -2 \\ 0 & 0 & 1 \\ -2 & 1 & 3 \end{bmatrix}$$

Check! $AA^{-1} = I = A^{-1}A$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

LU-decomposition by Gaussian elimination method.

e.g: Find the LU-decomposition of the given matrix, by using Gaussian elimination method

$$A = \begin{bmatrix} 2 & -1 & 4 \\ 4 & 1 & 9 \\ -2 & 10 & -3 \end{bmatrix}$$

$$A \sim \begin{bmatrix} 2 & -1 & 4 \\ 0 & 3 & 1 \\ 0 & 9 & 1 \end{bmatrix} \begin{matrix} -2R_1 + R_2 \\ 1R_1 + R_3 \end{matrix} \qquad \frac{-4}{2} R_1 + R_2 \quad \frac{2}{2} R_1 + R_3$$

$$\sim \begin{bmatrix} 2 & -1 & 4 \\ 0 & 3 & 1 \\ 0 & 0 & -2 \end{bmatrix} -3R_2 + R_3 \qquad \frac{-9}{3} R_2 + R_3$$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 3 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 2 & -1 & 4 \\ 0 & 3 & 1 \\ 0 & 0 & -2 \end{bmatrix}$$

check

$$LU = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 3 & 1 \end{bmatrix} \begin{bmatrix} 2 & -1 & 4 \\ 0 & 3 & 1 \\ 0 & 0 & -2 \end{bmatrix} = \begin{bmatrix} 2 & -1 & 4 \\ 4 & 1 & 9 \\ -2 & 10 & -3 \end{bmatrix} \checkmark$$

END.

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Special Matrices

1) Tridiagonal Matrix (Thomas method)

$$\begin{bmatrix} f_1 & g_1 & 0 \\ e_2 & f_2 & g_2 \\ 0 & e_3 & f_3 \end{bmatrix}$$

3x3 – tridiagonal matrix

$$\begin{bmatrix} f_1 & g_1 & 0 & 0 \\ e_2 & f_2 & g_2 & 0 \\ 0 & e_3 & f_3 & g_3 \\ 0 & 0 & e_4 & f_4 \end{bmatrix}$$

4x4 – tridiagonal matrix

$$\begin{bmatrix} f_1 & g_1 & 0 & 0 & 0 \\ e_2 & f_2 & g_2 & 0 & 0 \\ 0 & e_3 & f_3 & g_3 & 0 \\ 0 & 0 & e_4 & f_4 & g_4 \\ 0 & 0 & 0 & e_5 & f_5 \end{bmatrix}$$

5x5 matrix.

$$AX = B$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Thomas method

the aim is to find the LU-decomposition of the tridiagonal matrix.

$$e_k^{New} = \frac{e_k^{old}}{f_{k-1}}$$

$$f_k^{New} = -e_k g_{k-1} + f_k^{New}$$

$k = 2, 3, \dots, n$

e.g.: Find the LU-decomposition of

1) $A = \begin{bmatrix} 2 & 1 & 0 \\ 4 & -2 & -4 \\ 0 & -2 & 5 \end{bmatrix}$

$$\sim \begin{bmatrix} 2 & 1 & 0 \\ 2 & -4 & -4 \\ 0 & \frac{1}{2} & 7 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 0 \\ 2 & -4 & -4 \\ 0 & \frac{1}{2} & 7 \end{bmatrix}$$

e: $\frac{-2}{-4}$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & \frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ 0 & -4 & -4 \\ 0 & 0 & 7 \end{bmatrix}$$

L U

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2) $A = \begin{bmatrix} 2 & 2 & 0 & 0 \\ -2 & 3 & 2 & 0 \\ 0 & 6 & 5 & 4 \\ 0 & 0 & -2 & 2 \end{bmatrix}$ (3)

$\sim \begin{bmatrix} 2 & 2 & 0 & 0 \\ -1 & 5 & 2 & 0 \\ 0 & \frac{6}{5} & \frac{13}{5} & 4 \\ 0 & 0 & \frac{-10}{13} & \frac{66}{13} \end{bmatrix}$

$\frac{-6}{5} \times 2 + 5 = \frac{-12}{5} + \frac{25}{5} = \frac{13}{5}$

$-2 \div \frac{13}{5} = -2 * \frac{5}{13}$

$\frac{10}{13} \times 4 + 2 = \frac{40}{13} + \frac{26}{13} = \frac{66}{13}$

$\begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & \frac{6}{5} & 1 & 0 \\ 0 & 0 & \frac{-10}{13} & 1 \end{bmatrix} \begin{bmatrix} 2 & 2 & 0 & 0 \\ 0 & 5 & 2 & 0 \\ 0 & 0 & \frac{13}{5} & 4 \\ 0 & 0 & 0 & \frac{66}{13} \end{bmatrix}$

L U

KAMEL ELEIEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

[2] Symmetric Matrix (Cholesky Method)

Recall

A is symmetric if $A^T = A$

$$L = \begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix}, \quad U = L^T \quad \text{for } 2\times 2 \text{ matrix}$$

$$L = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix}, \quad U = L^T \quad \text{for } 3\times 3 \text{ matrix}$$

$$L = \begin{bmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ l_{31} & l_{32} & l_{33} & 0 \\ l_{41} & l_{42} & l_{43} & l_{44} \end{bmatrix}, \quad U = L^T, \quad \text{for } 4\times 4 \text{ matrix}$$

$$\vdots$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Use Cholesky method to find the LU-decomposition of

1) $A = \begin{bmatrix} 4 & 6 \\ 6 & 25 \end{bmatrix}$

$$\begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} \\ 0 & l_{22} \end{bmatrix} = \begin{bmatrix} 4 & 6 \\ 6 & 25 \end{bmatrix}$$

L $\quad$ U

$l_{11}^2 = 4 \rightarrow l_{11} = \sqrt{4} = 2 \rightarrow \boxed{l_{11} = 2}$

$l_{11} l_{21} = 6 \rightarrow 2 l_{21} = 6 \rightarrow \boxed{l_{21} = 3}$

$l_{21}^2 + l_{22}^2 = 25 \rightarrow 9 + l_{22}^2 = 25$

$\rightarrow l_{22}^2 = 16 \rightarrow \boxed{l_{22} = 4}$

$$\begin{bmatrix} 2 & 0 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 0 & 4 \end{bmatrix} = A$$

L $\quad$ U

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2) $A = \begin{bmatrix} 4 & 2 & 14 \\ 2 & 17 & -5 \\ 14 & -5 & 83 \end{bmatrix}$

$$\begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} & l_{31} \\ 0 & l_{22} & l_{32} \\ 0 & 0 & l_{33} \end{bmatrix} = \begin{bmatrix} 4 & 2 & 14 \\ 2 & 17 & -5 \\ 14 & -5 & 83 \end{bmatrix}$$

L $\quad$ U $\quad$ A

$l_{11}^2 = 4 \longrightarrow \boxed{l_{11} = 2}$

$l_{11} l_{21} = 2 \longrightarrow 2 l_{21} = 2 \rightarrow \boxed{l_{21} = 1}$

$l_{11} l_{31} = 14 \rightarrow 2 l_{31} = 14 \rightarrow \boxed{l_{31} = 7}$

$l_{21}^2 + l_{22}^2 = 17 \Rightarrow 1 + l_{22}^2 = 17$

$\Rightarrow l_{22}^2 = 16 \rightarrow \boxed{l_{22} = 4}$

$l_{21} l_{31} + l_{22} l_{32} = -5$

$(1)(7) + 4 l_{32} = -5 \Rightarrow 4 l_{32} = -12$

$\Rightarrow \boxed{l_{32} = -3}$

$l_{31}^2 + l_{32}^2 + l_{33}^2 = 83$

$49 + 9 + l_{33}^2 = 83 \rightarrow l_{33}^2 = 25 \rightarrow \boxed{l_{33} = 5}$

$$\begin{bmatrix} 2 & 0 & 0 \\ 1 & 4 & 0 \\ 7 & -3 & 5 \end{bmatrix} \begin{bmatrix} 2 & 1 & 7 \\ 0 & 4 & -3 \\ 0 & 0 & 5 \end{bmatrix} = A$$

L $\quad$ U

KAMEL ELGIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Gauss - Seidel method (iterative method)

Linear system $AX = B$

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & & & \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$$

This method will be convergent, if

$$|a_{ii}| > \sum_{\substack{j=1 \\ j \neq i}}^{n} |a_{ij}|$$

e.g.: By Gauss - Seidel method, solve

the system

$$6x - y - 2z = 12$$

$$3x + 5y + z = 3$$

$$2x - y + 7z = 36$$

initial guess

$x_0 = 0$, $y_0 = 0$, $z_0 = 0$

Do two iterations ?

Sol:

$$x = \frac{12 + y + 2z}{6}$$

$$y = \frac{3 - 3x - z}{5}$$

$$z = \frac{36 - 2x + y}{7}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

1st iteration $x_0=0, y_0=0, z_0=0$

$$x_1 = \frac{12+0+2(0)}{6} \rightarrow \boxed{x_1 = 2}$$

$$y_1 = \frac{3-3(2)-0}{5} \Rightarrow \boxed{y_1 = -0.6}$$

$$z_1 = \frac{36-2(2)+(-0.6)}{7} = \boxed{4.485714 = z_1}$$

2nd iteration $x_1=2, y_1=-0.6,$
$z_1 = 4.485714$

$$x_2 = \frac{12+(-0.6)+2(4.485714)}{6} \rightarrow \boxed{x_2 = 3.39524}$$

$$y_2 = \frac{3-3(3.39524)-(4.485714)}{5}$$

$$\boxed{y_2 = -2.334286}$$

$$z_2 = \frac{36-2(3.39524)+(-2.334286)}{7}$$

$$\boxed{z_2 = 3.89319}$$

Convergent

True solution $(\overset{x}{3}, \overset{y}{-2}, \overset{z}{4})$

·END·

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Least squares Regression (Best fitting curve)

Before starting

* Sigma - Notation $\sum$ (summation)

1) $\sum_{i=1}^{n} f(i) = f(1) + f(2) + \cdots + f(n)$

* $\sum_{i=1}^{4} i^2 = 1^2 + 2^2 + 3^2 + 4^2 = 1 + 4 + 9 + 16 = 30$

* $\sum_{i=1}^{5} x_i = x_1 + x_2 + x_3 + x_4 + x_5$

* $\sum_{i=1}^{3} (a + bx_i) = (a + bx_1) + (a + bx_2) + (a + bx_3)$

2) $\sum_{i=1}^{n} a = na$

* $\sum_{i=1}^{4} 5 = 5 + 5 + 5 + 5 = 20$

* $\sum_{i=1}^{100} 2 = \underbrace{2 + 2 + 2 + \cdots + 2}_{100 \text{ times}} = (100)(2) = 200$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

3) $\sum_{i=1}^{n}(f(i) \mp g(i)) = \sum_{i=1}^{n} f(i) \mp \sum_{i=1}^{n} g(i)$

* $\sum_{i=1}^{3}(i^2 + 2i - 3) = (1^2 + 2(1) - 3) + (2^2 + (2)(2) - 3) + (3^2 + 2(3) - 3)$

$= 0 + 5 + 12$

$= 17$ ✓

or $\sum_{i=1}^{3}(i^2 + 2i - 3) = \sum_{i=1}^{3} i^2 + \sum_{i=1}^{3} 2i - \sum_{i=1}^{3} 3$

$= (1^2 + 2^2 + 3^2) + (2(1) + 2(2) + 2(3)) - (3 + 3 + 3)$

$= 14 + 12 - 9$

$= 17$ ✓

4) $\sum_{i=1}^{n} c\, f(i) = c \sum_{i=1}^{n} f(i)$ c: Constant

* $\sum_{i=1}^{4}(5i^2 + 1) = (5(1^2) + 1) + (5(2^2) + 1) + (5(3^2) + 1) + (5(4^2) + 1)$

$= (5 + 1) + (20 + 1) + (45 + 1) + (80 + 1)$

$= 6 + 21 + 46 + 81$

$= 154$

or $\sum_{i=1}^{4}(5i^2 + 1) = 5\sum_{i=1}^{4} i^2 + \sum_{i=1}^{4} 1$

$= 5(1^2 + 2^2 + 3^2 + 4^2) + 4$

$= 5(30) + 4 = 154$ ✓

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Partial Derivatives

→ function in one variable.

$y = f(x) \longrightarrow \frac{dy}{dx} = f'(x)$ → ordinary derivative of f w.r.t x

→ Functions in two or more variables

* $Z = f(x, y) \longrightarrow \frac{\partial z}{\partial x} = f_x$ (Partial derivative of f w.r.t x)

$\frac{\partial z}{\partial y} = f_y$ (Partial derivative of f w.r.t y)

* $w = f(x, y, z) \longrightarrow \frac{\partial w}{\partial x} = f_x$ (Partial derivative of f w.r.t x)

$\frac{\partial w}{\partial y} = f_y$ (Partial derivative of f w.r.t y)

$\frac{\partial w}{\partial z} = f_z$ (Partial derivative of f w.r.t z)

⋮

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.9 Find all the partial derivatives of

1) $f(x,y) = x^3 + y^2 + xy + 1$

$f_x = 3x^2 + y$

$f_y = 2y + x$

2) $f(x,y,z) = xy + xz + yz$

$f_x = y + z$

$f_y = x + z$

$f_z = x + y$

3) $f(a,b,c) = (ax_i + by_i + cz_i)^2$ $i = 1, 2, 3, \ldots, n$

x_i, y_i, z_i are fixed

$f_a = 2(ax_i + by_i + cz_i)(x_i)$

$= 2x_i(ax_i + by_i + cz_i)$

$f_b = 2(ax_i + by_i + cz_i)(y_i)$

$= 2y_i(ax_i + by_i + cz_i)$

$f_c = 2(ax_i + by_i + cz_i)(z_i)$

$= 2z_i(ax_i + by_i + cz_i)$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Min. or Max. of any functions (in one or more variables) occur when all the first derivatives) are (is) zeros (zero).

e.g: Find the minimum of $f(x,y) = (x-2)^2 + (y-3)^2 + 1$

$f_x = 2(x-2)$

$f_y = 2(y-3)$

$f_x = 0 \rightarrow x = 2$

$f_y = 0 \rightarrow y = 3$

$\therefore$ the min. of f occurs at $(2,3)$

then min. $= f(2,3) = 1$.

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Now: Least-squares regression.

Assume we want to find a relationship (model) between two variables x, y after recording measurements of corresponding values of x and y. And assume we get n ordered pairs (data), namely

$(x_1, y_1), (x_2, y_2), \ldots, (x_n, y_n)$.

to see the shape of the better model, plot these points in the xy-plane.



scatter diagram





Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Linear Regression

Data : $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n) = \{(x_i, y_i)\}_{i=1}^{n}$

x : independent variable
y : dependent variable



y_i = the measured value of y when $x = x_i$

$\hat{y}_i = A + Bx_i$ the calculated value (from the model) when $x = x_i$

$\hat{y}_i$ is also called the predicted value of y when $x = x_i$

$$e_i = y_i - \hat{y}_i$$ ← e_i : error (residual)

Note
One of the important aims of the regression curve is to predict the value of dependent variable y at a given value (in the range) of the independent variable x

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

To obtain the best-fitting line (or curve) by least squares method, is, to minimize the sum of the squares of residuals

$$S_r = \sum_{i=1}^{n} e_i^2$$

Now, to find the best fitting line

$$y = A + Bx \quad \text{(Also called linear regression)}$$

$$S_r = \sum_{i=1}^{n} (y_i - \hat{y}_i)^2$$

$$S_r = \sum_{i=1}^{n} (y_i - A - Bx_i)^2$$

$$\frac{\partial S_r}{\partial A} = \sum_{i=1}^{n} 2(y_i - A - Bx_i)(-1)$$

$$= -2 \sum_{i=1}^{n} (y_i - A - Bx_i)$$

$$= -2 \left[\sum_{i=1}^{n} y_i - \sum_{i=1}^{n} A - \sum_{i=1}^{n} Bx_i \right]$$

$$= -2 \left[\sum_{i=1}^{n} y_i - nA - B \sum_{i=1}^{n} x_i \right]$$

$$\frac{\partial S_r}{\partial A} = 0 \rightarrow \boxed{nA + B \sum_{i=1}^{n} x_i = \sum_{i=1}^{n} y_i} \quad (1)$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\frac{\partial S_r}{\partial B} = \sum_{i=1}^{n} 2\,(y_i - A - Bx_i)(-x_i)$$

$$= -2 \sum_{i=1}^{n} (x_i y_i - A x_i - B x_i^2)$$

$$= -2 \left[\sum_{i=1}^{n} x_i y_i - A \sum_{i=1}^{n} x_i - B \sum_{i=1}^{n} x_i^2 \right]$$

$$\frac{\partial S_r}{\partial B} = 0 \rightarrow \boxed{A \sum_{i=1}^{n} x_i + B \sum_{i=1}^{n} x_i^2 = \sum_{i=1}^{n} x_i y_i} \quad (2)$$

so we get the following linear system

$$\begin{bmatrix} n & \Sigma x_i \\ \Sigma x_i & \Sigma x_i^2 \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \end{bmatrix}$$

we use Cramer's rule to find B.

$$B = \frac{\begin{vmatrix} n & \Sigma y_i \\ \Sigma x_i & \Sigma x_i y_i \end{vmatrix}}{\begin{vmatrix} n & \Sigma x_i \\ \Sigma x_i & \Sigma x_i^2 \end{vmatrix}} = \frac{n \Sigma x_i y_i - (\Sigma x_i)(\Sigma y_i)}{n \Sigma x_i^2 - (\Sigma x_i)^2}$$

$$\boxed{B = \frac{n \Sigma x_i y_i - (\Sigma x_i)(\Sigma y_i)}{n \Sigma x_i^2 - (\Sigma x_i)^2}}$$

- from statistics, $\bar{y} = A + B\bar{x}$

$$\rightarrow \boxed{A = \bar{y} - B\bar{x}}$$

where $\bar{y} = \frac{\Sigma y_i}{n}$, $\bar{x} = \frac{\Sigma x_i}{n}$.

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Fit a least-squares line to the following data. And hence, find the predicted value of y and the residual (error) when $x = 7$

x	1	3	5	7	9
y	4	2	6	8	10

Sol:

$y = A + Bx$ or $y_i = A + Bx_i + e_i$

or $y_i = \hat{y}_i + e_i$

$$\begin{bmatrix} n & \Sigma x_i \\ \Sigma x_i & \Sigma x_i^2 \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \end{bmatrix}$$

	x_i	y_i	x_i^2	$x_i y_i$
1	1	4	1	4
2	3	2	9	6
3	5	6	25	30
4	7	8	49	56
5	9	10	81	90
Σ	25	30	165	186

$$\Rightarrow \begin{bmatrix} 5 & 25 \\ 25 & 165 \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} = \begin{bmatrix} 30 \\ 186 \end{bmatrix}$$

$\Rightarrow A = 1.5$, $B = 0.9$

A: y-intercept, B: slope

$y = 1.5 + 0.9x$

$n = 5$

$x = 7 = x_4$

$\hat{y}_4 = 1.5 + 0.9(7) = 7.8$

$e_4 = y_4 - \hat{y}_4 = 8 - 7.8 = 0.2$

END

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Polynomial Regression

* Least squares Parabola (Quadratic)

Ex fitting the data $\{(x_i, y_i)\}_{i=1}^{n}$ by Least squares parabola

$\rightarrow$ $y = a + bx + cx^2 + e$ — residual (error)

$$y_i = a + bx_i + cx_i^2 + e_i$$

$$S_r = \sum_{i=1}^{n} e_i^2$$ sum of the squares of residuals

$$S_r = \sum_{i=1}^{n} (y_i - a - bx_i - cx_i^2)^2$$

solving the equations

$$\frac{\partial S_r}{\partial a} = 0 \quad , \quad \frac{\partial S_r}{\partial b} = 0 \quad \text{and} \quad \frac{\partial S_r}{\partial c} = 0$$

we get the following linear system

$$n a + b\Sigma x_i + c\Sigma x_i^2 = \Sigma y_i \quad -(1)$$

$$a\Sigma x_i + b\Sigma x_i^2 + c\Sigma x_i^3 = \Sigma x_i y_i \quad -(2)$$

$$a\Sigma x_i^2 + b\Sigma x_i^3 + c\Sigma x_i^4 = \Sigma x_i^2 y_i \quad -(3)$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Using matrices we can write the above system as

$$\begin{bmatrix} n & \Sigma x_i & \Sigma x_i^2 \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i^3 \\ \Sigma x_i^2 & \Sigma x_i^3 & \Sigma x_i^4 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \\ \Sigma x_i^2 y_i \end{bmatrix}$$

In the same manner, we can find the linear system, if we want a cubic regression

$$y_i = a + bx_i + cx_i^2 + dx_i^3 + e_i$$

$$S_r = \sum_{i=1}^{n} (y_i - a - bx_i - cx_i^2 - dx_i^3)^2$$

$$\begin{bmatrix} n & \Sigma x_i & \Sigma x_i^2 & \Sigma x_i^3 \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i^3 & \Sigma x_i^4 \\ \Sigma x_i^2 & \Sigma x_i^3 & \Sigma x_i^4 & \Sigma x_i^5 \\ \Sigma x_i^3 & \Sigma x_i^4 & \Sigma x_i^5 & \Sigma x_i^6 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \\ \Sigma x_i^2 y_i \\ \Sigma x_i^3 y_i \end{bmatrix}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Using least squares method, fit the following data by a quadratic regression

x_i	1.1	2	3.2	4	4.5	5.2
y_i	0.5	2.5	2	4	3.5	6.5

Sol: $y_i = a + bx_i + cx_i^2 + e_i$

$n = 6$,

$$\begin{bmatrix} n & \Sigma x_i & \Sigma x_i^2 \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i^3 \\ \Sigma x_i^2 & \Sigma x_i^3 & \Sigma x_i^4 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma y_i \\ \Sigma x_i y_i \\ \Sigma x_i^2 y_i \end{bmatrix}$$

	x_i	y_i	x_i^2	x_i^3	x_i^4	$x_i y_i$	$x_i^2 y_i$
1	1.1	0.5	1.21	1.331	1.4641	0.55	0.605
2	2	2.5	4	8	16	5	10
3	3.2	2	10.24	32.768	104.8576	6.4	20.48
4	4	4	16	64	256	16	64
5	4.5	3.5	20.25	91.125	410.0625	15.75	70.875
$n \to 6$	5.2	6.5	27.04	140.608	731.1616	33.8	175.76
	Σx_i	Σy_i	Σx_i^2	Σx_i^3	Σx_i^4	$\Sigma x_i y_i$	$\Sigma x_i^2 y_i$
Σ	20	19	78.74	337.832	1519.5458	77.5	341.72

$$\begin{bmatrix} 6 & 20 & 78.74 \\ 20 & 78.74 & 337.832 \\ 78.74 & 337.832 & 1519.5458 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 19 \\ 77.5 \\ 341.72 \end{bmatrix}$$

$a = 1.178264$, $b = -0.388651$, $c = 0.250234$

$\therefore y_i = 1.178264 - 0.388651 x_i + 0.250234 x_i^2 + e_i$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Multiple linear regression (Plane)

To fit the data

$$\{(x_i, y_i, z_i)\}_{i=1}^{n} = \{(x_1, y_1, z_1), (x_2, y_2, z_2), \dots (x_n, y_n, z_n)\}$$

by multiple linear regression, namely

$$z_i = a + bx_i + cy_i + e_i \qquad i = 1, 2, \dots, n$$

write $S_r = \sum_{i=1}^{n} e_i^2 = \sum_{i=1}^{n} (z_i - a - bx_i - cy_i)^2$

and then, put $\frac{\partial S_r}{\partial a} = 0$, $\frac{\partial S_r}{\partial b} = 0$ and $\frac{\partial S_r}{\partial c} = 0$

to get the following linear system

$$\begin{aligned} n a + b\Sigma x_i + c\Sigma y_i &= \Sigma z_i \quad (1) \\ a\Sigma x_i + b\Sigma x_i^2 + c\Sigma x_i y_i &= \Sigma x_i z_i \quad (2) \\ a\Sigma y_i + b\Sigma x_i y_i + c\Sigma y_i^2 &= \Sigma y_i z_i \quad (3) \end{aligned}$$

or by matrices

$$\begin{bmatrix} n & \Sigma x_i & \Sigma y_i \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i y_i \\ \Sigma y_i & \Sigma x_i y_i & \Sigma y_i^2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma z_i \\ \Sigma x_i z_i \\ \Sigma y_i z_i \end{bmatrix}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g : Use multiple Linear regression (plane), to fit the following data : (x, y, z)

x	1	3	5	7	9
y	2	6	4	8	10
z	7	10	16	20	30

and hence, find the predicted value of z when $x = 7, y = 8$ and the residual ?

Sol:

$$z_i = a + bx_i + cy_i + e_i$$

$$\begin{bmatrix} n & \Sigma x_i & \Sigma y_i \\ \Sigma x_i & \Sigma x_i^2 & \Sigma x_i y_i \\ \Sigma y_i & \Sigma x_i y_i & \Sigma y_i^2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \Sigma z_i \\ \Sigma x_i z_i \\ \Sigma y_i z_i \end{bmatrix}$$

$n = 5$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

	x_i	y_i	z_i	x_i^2	$x_i y_i$	$x_i z_i$	y_i^2	$y_i z_i$
1	1	2	7	1	2	7	4	14
2	3	6	10	9	18	30	36	60
3	5	4	16	25	20	80	16	64
4	7	8	20	49	56	140	64	160
$n=5$	9	10	30	81	90	270	100	300
	25	30	83	165	186	527	220	598

$$\begin{bmatrix} 5 & 25 & 30 \\ 25 & 165 & 186 \\ 30 & 186 & 220 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 83 \\ 527 \\ 598 \end{bmatrix}$$

$$a = \frac{262}{95} \quad , \quad b = \frac{55}{19} \quad , \quad c = -\frac{2}{19}$$

$$\therefore \boxed{Z_i = \frac{262}{95} + \frac{55}{19} x_i - \frac{2}{19} y_i + e_i}$$

Now

when $x = 7 = x_4$, $y = 8 = y_4$

the predicted value of $Z_4 = \hat{Z}_4 = \frac{262}{95} + \frac{55}{19}(7) - \frac{2}{19}(8)$

$$\therefore \hat{Z}_4 \simeq 22.17895 \quad (\text{to 5 d.p's})$$

$$\therefore \text{the residual } (e_4) = Z_4 - \hat{Z}_4$$

$$= 20 - 22.17895$$

$$= -0.17895$$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

e.g: By using least squares method, to fit the data $\{(x_i, y_i)\}_{i=1}^{n}$ by the model $y = Ax^4$, derive a formula to compute A?

$$y_i = Ax_i^4 + e_i$$

$$S_r = \sum_{i=1}^{n} e_i^2$$

$$= \sum_{i=1}^{n} (y_i - Ax_i^4)^2$$

$$\frac{dS_r}{dA} = \sum_{i=1}^{n} 2(y_i - Ax_i^4)^1(-x_i^4)$$

$$= -2 \sum_{i=1}^{n} (x_i^4 y_i - Ax_i^8)$$

$$\frac{dS_r}{dA} = -2\left[\sum_{i=1}^{n} x_i^4 y_i - A\sum x_i^8\right]$$

$$\frac{dS_r}{dA} = 0 \rightarrow -2\left[\sum_{i=1}^{n} x_i^4 y_i - A\sum_{i=1}^{n} x_i^8\right] = 0$$

$$\rightarrow \sum_{i=1}^{n} x_i^4 y_i - A\sum_{i=1}^{n} x_i^8 = 0$$

$$A = \frac{\sum_{i=1}^{n} x_i^4 y_i}{\sum_{i=1}^{n} x_i^8}$$

$$A = \frac{\sum_{i=1}^{n} x_i^4 y_i}{\sum_{i=1}^{n} x_i^8}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Linearization of Nonlinear relationships:

① $y = A e^{Bx}$ exponential model

$\ln y = \ln(A e^{Bx})$

$\ln y = \ln A + \ln e^{Bx}$

$\ln y = \ln A + Bx \ln e$

$\ln e = 1$

$\ln y = \ln A + Bx$

$y = \ln A + Bx$

$Y = \ln A + Bx$

B: slope

$\ln A$: intercept

② $y = A x^{B}$ simple power model

$\rightarrow \ln y = \ln(A x^{B})$

$\ln y = \ln A + \ln x^{B}$

$\ln y = \ln A + B \ln x$

$Y = \ln A + BX$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

3. $y = \frac{Ax}{B+x}$ saturation-growth-rate

$$\rightarrow \frac{1}{y} = \frac{B+x}{Ax}$$

$$\frac{1}{y} = \frac{B}{Ax} + \frac{x}{Ax}$$

$$\frac{1}{y} = \frac{B}{A}\left(\frac{1}{x}\right) + \left(\frac{1}{A}\right)$$

$$Y = \frac{B}{A}X + \frac{1}{A}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Fit the following data by the exponential model $y = Ae^{Bx}$ (by linearization)

x	2	4	6	8
y	2	3	3.5	4.5

Sol:

$$y = Ae^{Bx} \longrightarrow \ln y = \ln(Ae^{Bx})$$

$$\Rightarrow \boxed{\ln y = \ln A + Bx}$$

$$\begin{bmatrix} n & \Sigma x \\ \Sigma x & \Sigma x^2 \end{bmatrix} \begin{bmatrix} \ln A \\ B \end{bmatrix} = \begin{bmatrix} \Sigma \ln y \\ \Sigma x \ln y \end{bmatrix}$$

$n = 4$

	x	y	$\ln y$	x^2	$x \ln y$
1	2	2	0.693147	4	1.386294
2	4	3	1.098612	16	4.394448
3	6	3.5	1.252763	36	7.516578
$n=4$	8	4.5	1.504077	64	12.032616
Σ	20		4.548599	120	25.329936

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$\begin{bmatrix} 4 & 20 \\ 20 & 120 \end{bmatrix} \begin{bmatrix} \ln A \\ B \end{bmatrix} = \begin{bmatrix} 4.548599 \\ 25.329936 \end{bmatrix}$$

$\Rightarrow$ $\ln A = 0.4904145$ and $B = 0.129347$

$$A = e^{0.4904145}$$

$$A = 1.632993$$

$$\therefore y = 1.632993\, e^{0.129347x}$$

END

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Interpolation:

→ Polynomial interpolation:

If $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$ — (n+1) points — (precise data), then there exists a unique polynomial of the n^{th} degree (n^{th} order) that passing through these n+1 points.

The polynomial interpolation is to estimate the value of y corresponding to the intermediate value x between $x_0, x_1, \dots, x_n$, by substituting x in this function. i.e $y = f_n(x)$

Also, we can approximate a complicated function that passes through the points $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$ $\left(f(x_i) = y_i\right)_{i=0,1,2,\dots,n}$ by polynomial of degree n, namely $f_n(x)$ $x \in [x_0, x_n]$

Linear interpolation

$f_1(x) = a_0 + a_1 x$



$f_2(x) = a_0 + a_1 x + a_2 x^2$



Quadratic interpolation
(Prabola)

$f_3(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3$



Cubic interpolation

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

1) Newton's Divided Difference Interpolation Polynomials:

Def:

If we have $(x_0, f(x_0))$, $(x_1, f(x_1))$, $\cdots$, $(x_n, f(x_n))$

1) First finite divided difference

$$f[x_k, x_{k-1}] = \frac{f(x_k) - f(x_{k-1})}{x_k - x_{k-1}} \quad , k = 1, 2, \cdots, n.$$

2) Second finite divided difference

$$f[x_k, x_{k-1}, x_{k-2}] = \frac{f[x_k, x_{k-1}] - f[x_{k-1}, x_{k-2}]}{x_k - x_{k-2}}$$

$$k = 2, 3, \cdots, n.$$

3) third finite divided difference

$$f[x_k, x_{k-1}, x_{k-2}, x_{k-3}] = \frac{f[x_k, x_{k-1}, \cdots x_{k-2}] - f[x_{k-1}, \cdots x_{k-3}]}{x_k - x_{k-3}}$$

$$k = 3, 4, 5, \cdots, n.$$

$\vdots$

n) the n^{th} finite divided difference

$$f[x_n, x_{n-1}, x_{n-2}, \cdots x_0] = \frac{f[x_n, x_{n-1}, \cdots, x_1] - f[x_{n-1}, x_{n-2}, \cdots x_0]}{x_n - x_0}$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Examples

* $f[x_1, x_0] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$

* $f[x_2, x_1] = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$

* $f[x_5, x_4] = \frac{f(x_5) - f(x_4)}{x_5 - x_4}$

* $f[x_2, x_1, x_0] = \frac{f[x_2, x_1] - f[x_1, x_0]}{x_2 - x_0}$

* $f[x_3, x_2, x_1] = \frac{f[x_3, x_2] - f[x_2, x_1]}{x_3 - x_1}$

* $f[x_4, x_3, x_2] = \frac{f[x_4, x_3] - f[x_3, x_2]}{x_3 - x_2}$

* $f[x_3, x_2, x_1, x_0] = \frac{f[x_3, x_2, x_1] - f[x_2, x_1, x_0]}{x_3 - x_0}$

* $f[x_4, x_3, x_2, x_1] = \frac{f[x_4, x_3, x_2] - f[x_3, x_2, x_1]}{x_4 - x_1}$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Linear Interpolation

Data : $(x_0, f(x_0))$, $(x_1, f(x_1))$

$$f_1(x) = b_0 + b_1(x - x_0)$$ Linear interpolation formula



where $b_0 = f(x_0)$

$$b_1 = f[x_1, x_0] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$

---

* Quadratic Interpolation (Parabola)

Data : $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$



$$f_2(x) = b_0 + b_1(x - x_0) + b_2(x - x_0)(x - x_1)$$

↳ Quadratic formula

where $b_0 = f(x_0)$

$$b_1 = f[x_1, x_0] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$

$$b_2 = f[x_2, x_1, x_0] = \frac{f[x_2, x_1] - f[x_1, x_0]}{x_2 - x_0}$$

x_i	$f(x_i)$	First D.D	Second D.D
x_0	$f(x_0)$ b_0		
		$f[x_1, x_0]$ b_1	
x_1	$f(x_1)$		$f[x_2, x_1, x_0]$ b_2
		$f[x_2, x_1]$	
x_2	$f(x_2)$		

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Cubic Interpolation

Data : $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$, $(x_3, f(x_3))$

$$f_3(x) = b_0 + b_1(x-x_0) + b_2(x-x_0)(x-x_1) + b_3(x-x_0)(x-x_1)(x-x_2)$$

Cubic interpolation formula

when $b_0 = f(x_0)$

$b_1 = f[x_1, x_0]$

$b_2 = f[x_2, x_1, x_0]$

$b_3 = f[x_3, x_2, x_1, x_0]$



x_i	$f(x_i)$	First D.D	Second D.D	Third D.D
x_0	$f(x_0)$ ← b_0			
		$f[x_1, x_0]$ ← b_1		
x_1	$f(x_1)$		$f[x_2, x_1, x_0]$ ← b_2	
		$f[x_2, x_1]$		$f[x_3, x_2, x_1, x_0]$ ← b_3
x_2	$f(x_2)$		$f[x_3, x_2, x_1]$	
		$f[x_3, x_2]$		
x_3	$f(x_3)$			

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* n^{th} (order) degree polynomial Interpolation.

Data: $(x_0, f(x_0))$, $(x_1, f(x_1))$, ... , $(x_n, f(x_n))$

$$f_n(x) = b_0 + b_1(x-x_0) + b_2(x-x_0)(x-x_1) + \cdots + b_n(x-x_0)(x-x_1)\cdots(x-x_{n-1})$$

nth degree polynomial interpolation formula

where $b_0 = f(x_0)$, $b_1 = f[x_1, x_0]$, $b_2 = f[x_2, x_1, x_0]$, ...

$$b_n = f[x_n, x_{n-1}, \ldots, x_0]$$

---

e.g: Estimate $(\ln 3)$ using linear interpolation by taking a) $x_0 = 1$, $x_1 = 5$

b) $x_0 = 2$, $x_2 = 4$

and compute the percent absolute relative true error assume the true value of ln3 is 1.098612

Sol: $f(x) = \ln x$

a) $x_0 = 1 \rightarrow f(x_0) = 0$

$x_1 = 5 \longrightarrow f(x_1) = 1.609438$

$f_1(x) = b_0 + b_1(x-x_0)$

$b_0 = 0$, $b_1 = f[x_1, x_0] = \dfrac{f(x_1) - f(x_0)}{x_1 - x_0} = \dfrac{1.609438 - 0}{5 - 1}$

$\therefore b_1 = 0.402360$

$\therefore f_1(x) = 0 + 0.402360(x - 1)$

$f_1(x) = 0.402360(x-1)$ ✓

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$\therefore \ln 3 \cong f_1(3) = 0.402360\,(3-1) = 0.804719$ (7)

$\therefore \ln 3 = 0.804719$

$\therefore |\epsilon_t| * 100\% = \left| \frac{1.098612 - 0.804719}{1.098612} \right| * 100\%$

$= 26.75\%$



b) $x_0 = 2 \Rightarrow f(x_0) = 0.693147$

$x_1 = 4 \Rightarrow f(x_1) = 1.386294$

$f_1(x) = b_0 + b_1(x-2)$

$b_0 = f(x_0) = 0.693147$

$b_1 = f[x_1, x_0] = \frac{f(x_1) - f(x_0)}{x_1 - x_0} = \frac{1.386294 - 0.693147}{4-2}$

$= 0.3465735$

$\therefore f_1(x) = 0.693147 + 0.3465735(x-2)$

$f_1(x) = 0.693147 + 0.3465735(x-2)$

$f(3) = \ln 3 \cong f_1(3) = 1.0397205$

$|\epsilon_t| = 5.36\%$



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Use the given data, to find the Newton's 3rd-order interpolating polynomial, and hence estimate $f(3)$?

x	-2	2	4	8
$f(x)$	3	7	13	17

Sol:

$$f_3(x) = b_0 + b_1(x-x_0) + b_2(x-x_0)(x-x_1) + b_3(x-x_0)(x-x_1)(x-x_2)$$

x_i	$f(x_i)$	1st FDD	2nd FDD	3rd FDD
x_0 -2	3 (b_0)			
		1 (b_1)		
x_1 2	7		$\frac{1}{3}$ (b_2)	
		3		$-\frac{1}{15}$ (b_3)
x_2 4	13		$-\frac{1}{3}$	
		1		
x_3 8	17			

$$f_3(x) = 3 + 1(x+2) + \frac{1}{3}(x+2)(x-2) - \frac{1}{15}(x+2)(x-2)(x-4)$$

$$f(3) \cong f_3(3) = 3 + (5) + \frac{1}{3}(5)(1) - \frac{1}{15}(5)(1)(-1)$$

$$= 10$$

– END –

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

2) Lagrange Interpolating Polynomials:

Introduction: Product Notation (Π-notation)

* $\prod_{i=1}^{3} i^2 = (1^2)(2^2)(3^2) = (1)(4)(9) = 36$

* $\prod_{i=0}^{4} (x_i - 5) = (x_0 - 5)(x_1 - 5)(x_2 - 5)(x_3 - 5)(x_4 - 5)$

* $\prod_{\substack{i=0 \\ i \neq 3}}^{5} (x_i - x_3) = (x_0 - x_3)(x_1 - x_3)(x_2 - x_3)(x_4 - x_3)(x_5 - x_3)$

If $(x_0, f(x_0))$, $(x_1, f(x_1))$, ... $(x_n, f(x_n))$ $n+1$ - points (precise data)

then the Lagrange Interpolating polynomial of n-order is

$$f_n(x) = \sum_{i=0}^{n} L_i(x) f(x_i)$$

$$= L_0(x) f(x_0) + L_1(x) f(x_1) + L_2(x) f(x_2) + \cdots + L_n(x) f(x_n)$$

where $L_i(x) = \prod_{\substack{j=0 \\ j \neq i}}^{n} \frac{x - x_j}{x_i - x_j}$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Lagrange Interpolating Polynomial of first order

Data : $(x_0, f(x_0))$, $(x_1, f(x_1))$

$$f_1(x) = L_0(x) f(x_0) + L_1(x) f(x_1)$$

where $L_0(x) = \frac{x - x_1}{x_0 - x_1}$, $L_1(x) = \frac{x - x_0}{x_1 - x_0}$

* Lagrang Interpolating Polynomial of second order

Data : $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$

$$f_2(x) = L_0(x) f(x_0) + L_1(x) f(x_1) + L_2(x) f(x_2)$$

where

$$L_0(x) = \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)}$$

$$L_1(x) = \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)}$$

$$L_2(x) = \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

* Lagrange Interpolation Polynomial of third-order.

Data: $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$, $(x_3, f(x_3))$

$$f_3(x) = L_0(x) f(x_0) + L_1(x) f(x_1) + L_2(x) f(x_2) + L_3(x) f(x_3)$$

where

$$L_0(x) = \frac{(x-x_1)(x-x_2)(x-x_3)}{(x_0-x_1)(x_0-x_2)(x_0-x_3)}$$

$$L_1(x) = \frac{(x-x_0)(x-x_2)(x-x_3)}{(x_1-x_0)(x_1-x_2)(x_1-x_3)}$$

$$L_2(x) = \frac{(x-x_0)(x-x_1)(x-x_3)}{(x_2-x_0)(x_2-x_1)(x_2-x_3)}$$

$$L_3(x) = \frac{(x-x_0)(x-x_1)(x-x_2)}{(x_3-x_0)(x_3-x_1)(x_3-x_2)}$$

KAMEL FLEIFEL

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: If $f(x) = \frac{x^2+4}{x^2+1}$, $x_0 = -1$, $x_1 = 2$, $x_3 = 3$

use the data $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$

to find the Lagrange interpolating function (Polynomial) of 2nd order. Then use it to estimate $f(1)$ and compute the absolute percent relative true error ($|\epsilon_t| * 100\%$).

Sol:

x_0 $f(x_0)$, x_1 $f(x_1)$, x_2 $f(x_2)$

$(-1, 2.5)$, $(2, 1.6)$, $(3, 1.3)$

$$f_2(x) = L_0(x) f(x_0) + L_1(x) f(x_1) + L_2(x) f(x_2)$$

i.e $$f_2(x) = \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)} f(x_0) + \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_1-x_2)} f(x_1) + \frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)} f(x_2)$$

$$\therefore f_2(x) = \frac{(x-2)(x-3)}{(-1-2)(-1-3)}(2.5) + \frac{(x+1)(x-3)}{(2+1)(2-3)}(1.6) + \frac{(x+1)(x-2)}{(3+1)(3-2)}(1.3$$

$$f_2(x) = \frac{5}{24}(x-2)(x-3) - \frac{8}{15}(x+1)(x-3) + \frac{13}{40}(x+1)(x-2) \checkmark$$

$$f(1) \cong f_2(1) = 1.9 \checkmark$$

True value $= f(1) = \frac{1^2+4}{1^2+1} = \frac{5}{2} = 2.5 \checkmark$

$$\epsilon_t = \left|\frac{2.5 - 1.9}{2.5}\right| * 100\% = 24\%$$

- END -

KAMEL FLEIFEL

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

Numerical Integration:

Trapezoidal Rule

Introduction

If $[a,b]$ is a closed interval

$x_0 = a < x_1 < x_2 < \cdots < x_n = b$ then

$P = \{x_0, x_1, \ldots, x_n\}$ is called a uniform partition of the closed interval $[a,b]$ if

(step size) $\Delta x = h = x_1 - x_0 = x_2 - x_1 = \cdots = x_n - x_{n-1} = \frac{b-a}{n}$

and $[x_0, x_1], [x_1, x_2], \ldots, [x_{n-1}, x_n]$ are called subintervals or segments



Notes: ① The number of segments $= n$ and (subintervals)

the number of the points $x_0, x_1, \ldots, x_n$ equals $n+1$

i.e

the number of segments + 1 = number of points

② $h = \frac{b-a}{n}$

$x_i = a + ih \quad i = 0, 1, 2, \ldots, n \qquad x_0 = a$

$x_0 = a + 0h = a$

$x_1 = a + h$ ... $x_n = a + nh = a + n\frac{b-a}{n} = b$ $\quad x_2 = a + 2h$ $\quad x_n = a + nh$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Find the step size and the uniform partition for the interval $[-1,3]$, if number of subintervals (segments) is equal to 4?

Sol:

$a = -1$, $b = 3$, $n = 4$

$\Rightarrow \Delta x_i = h = \frac{3-(-1)}{4} = 1$ (step size)

$x_0 = a = -1$

$x_1 = a + h = -1 + 1 = 0$

$x_2 = x_1 + h = a + 2h = 0 + 1 = 1$

$x_3 = x_2 + h = 1 + 1 = 2$

$x_4 = x_3 + h = 2 + 1 = 3$

$P = \{-1, 0, 1, 2, 3\}$.

e.g: If P is a uniform partition of the interval $[1,2]$ and the step size $= 0.1$, find the number of elements of P.

Sol: $a = 1$, $b = 2$, $h = 0.1$, $n =$ number of segments.

$h = \frac{b-a}{n} = \frac{2-1}{n} \Rightarrow \frac{1}{n} = 0.1$

$\Rightarrow 0.1n = 1 \Rightarrow n = 10$ segments

$\therefore$ the number of the elements of P is 11.

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Trapezoidal Rule: (simple T.R)

$f(x) \geq 0 \quad \forall x \in [a,b]$

$\int_a^b f(x)\,dx$ = Area under curve and above x-axis.



$\Rightarrow \int_a^b f(x)dx \simeq$ Area of the trapezoid

$\simeq \frac{h}{2}[f(a)+f(b)]$

$\simeq \frac{b-a}{2}(f(a)+f(b))$ ✓

$h = \frac{x_1 - x_0}{1} = \frac{b-a}{1}$

$h = b-a$

$$\int_a^b f(x)dx = \frac{h}{2}(f(a)+f(b)) \qquad h = \frac{b-a}{1},\ n=1$$

$$= \frac{h}{2}(f(x_0)+f(x_1))$$

or

$$\int_a^b f(x)dx = \frac{b-a}{2}(f(a)+f(b))$$

Also

$$\int_a^b f(x)dx \simeq \int_a^b f_1(x)\,dx$$



$$\simeq \int_a^b \left[f(a) + \frac{f(b)-f(a)}{b-a}(x-a)\right]dx$$

$$\simeq \frac{b-a}{2}(f(a)+f(b)).$$ ✓

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Use simple trapezoidal rule, to approximate $\int_0^1 x^3 dx$ and find the absolute relative error ?

Sol:

$$n = 1 \Rightarrow h = \frac{b-a}{1} = \frac{1-0}{1} = 1$$

$$a = 0 \,, \quad b = 1 \,, \quad f(x) = x^3$$

$$\Rightarrow \int_0^1 x^3 dx = \frac{1}{2}[f(0) + f(1)]$$

$$= \frac{1}{2}[0 + 1]$$

$$= 0.5$$

True value $\int_0^1 x^3 dx = \frac{x^4}{4}\Big|_0^1 = 0.25$.

$$\epsilon_t = \frac{|0.25 - 0.5|}{0.25} * 100\% = 100\% .$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Multiple - Application Trapezoidal Rule

Composite integration formula :T, R

$$\int_a^b f(x)dx \cong \frac{h}{2}(f(x_0)+f(x_1)) + \frac{h}{2}(f(x_1)+f(x_2)) + \frac{h}{2}(f(x_2)+f(x_3)) + \cdots + \frac{h}{2}(f(x_{n-2})+f(x_{n-1})) + \frac{h}{2}(f(x_{n-1})+f(x_n))$$



$x_{i+1} - x_i = h$ for any i

$$\Rightarrow \int_a^b f(x)dx \cong \frac{h}{2}[f(x_0)+f(x_1) + f(x_1)+f(x_2) + f(x_2)+f(x_3) + \cdots + f(x_{n-2})+f(x_{n-1}) + f(x_{n-1})+f(x_n)]$$

$$\Rightarrow \int_a^b f(x)dx \cong \frac{h}{2}[f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)]$$

$$\Rightarrow \int_a^b f(x)dx \cong \frac{h}{2}\left[f(x_0) + 2\sum_{i=1}^{n-1} f(x_i) + f(x_n)\right]$$

$$h = \frac{b-a}{n}$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g.: Use Multiple-Application Trapezoidal Rule, take the number of segments to be 4, to approximate $\int_0^1 x^3 dx$ and compute $\epsilon_t\%$.

Sol:

$n = 4 \Rightarrow h = \frac{b-a}{4} = \frac{1-0}{4} = 0.25$, $f(x) = x^3$

$x_0 = a = 0$, $x_1 = 0.25$, $x_2 = 0.5$, $x_3 = 0.75$, $x_4 = 1 = b$.

$$\therefore \int_0^1 x^3 dx \simeq \frac{0.25}{2}\left[0^3 + 2\left(0.25^3 + 0.5^3 + 0.75^3\right) + 1^3\right]$$

$$\simeq 0.265625$$

True value: $\int_0^1 x^3 dx = 0.25$

$$\therefore \quad \epsilon_t\% = \left|\frac{0.25 - 0.265625}{0.25}\right| * 100\%$$

$$= 6.25\%$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g : Use Multiple-Application Trapezoidal formula (take 4 points) to approximate

$$\int_0^{\pi/2} \sin x^2 \, dx$$ and compute $\epsilon_t\%$?

Sol:

$$f(x) = \sin x^2$$

$$a = 0, \quad b = \frac{\pi}{2}, \quad n = 3$$

$$h = \frac{b-a}{3} = \frac{\frac{\pi}{2} - 0}{3} = \frac{\pi}{6}$$

$$\Rightarrow x_0 = 0, \quad x_1 = \frac{\pi}{6}, \quad x_2 = \frac{\pi}{3}, \quad x_3 = \frac{\pi}{2}$$

$$\int_0^{\pi/2} \sin x^2 dx \simeq \frac{h}{2}\left[f(x_0) + 2(f(x_1) + f(x_2)) + f(x_3)\right]$$

$$\simeq \frac{\pi}{12}\left[\sin 0^2 + 2\left(\sin\left(\frac{\pi}{6}\right)^2 + \sin\left(\left(\frac{\pi}{3}\right)^2\right)\right) + \sin\left(\left(\frac{\pi}{2}\right)^2\right)\right]$$

$\sin\left(\left(\frac{\pi}{2}\right)^2\right)$

$$\therefore \int_0^{\pi/2} \sin x^2 \, dx \simeq 1.542037743$$

True value = 0.828116328 8

$$\epsilon_t = \left|\frac{\text{True} - \text{app.}}{\text{True}}\right| * \%$$

END

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Simpson's Rules

Simple Simpson's $\frac{1}{3}$-rule

$n=2$ in order to replace the integrand by a quadratic function.

$$\int_a^b f(x)dx \cong \int_a^b f_2(x)\,dx$$



$$n=2 \Rightarrow h = \frac{b-a}{2}$$

$$\Rightarrow x_0 = a \quad , x_1 = a + \frac{b-a}{2} = \frac{a+b}{2} \quad , x_2 = b$$

$$f_2(x) = \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)} f(x_0) + \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_1-x_2)} f(x_1) +$$

$$\frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)} f(x_2)$$

After integrating $f_2(x)$ over the interval $[a,b]$

we get the following rule (Simpson $\frac{1}{3}$-rule)

$$\int_a^b f(x)dx = \frac{h}{3}\left[f(x_0) + 4f(x_1) + f(x_2) \right].$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Use simple Simpson's $\frac{1}{3}$-rule to approximate $\int_1^2 x^2 dx$.

Sol: $f(x) = x^2$, $n=2$, $a=1$, $b=2 \Rightarrow h = \frac{2-1}{2} = 0.5$

$x_0 = 1$, $x_1 = 1.5$, $x_2 = 2$.

$$\therefore \int_1^2 x^2 dx \cong \frac{h}{3}\left[f(x_0) + 4f(x_1) + f(x_2)\right]$$

$$\cong \frac{0.5}{3}\left[1^2 + 4(1.5^2) + 2^2\right]$$

$$\cong \frac{7}{3}.$$

e.g: Find the length of the curve of $y = e^{-x}$ from $x=0$ to $x=1$, by using simple simpson's $\frac{1}{3}$-rule to approximate the integral.

Hint: the length of the curve of y from $x=a$ to $x=b$ is given by

$$L = \int_a^b \sqrt{\left(\frac{dy}{dx}\right)^2 + 1}\, dx$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Sol:



$$L = \int_0^1 \sqrt{\left(\frac{dy}{dx}\right)^2 + 1}\, dx$$

$$\frac{dy}{dx} = -e^{-x}$$

$$\Rightarrow L = \int_0^1 \sqrt{(-e^{-x})^2 + 1}\, dx$$

$$L = \int_0^1 \sqrt{e^{-2x} + 1}\, dx$$

Now $a = 0$, $b = 1$, $n = 2$, $f(x) = \sqrt{e^{-2x}+1}$

$$\Rightarrow h = \frac{1-0}{2} = 0.5 \Rightarrow x_0 = 0,\ x_1 = 0.5,\ x_2 = 1$$

$$\therefore L \simeq \frac{0.5}{3}\left[f(x_0) + 4f(x_1) + f(x_2)\right]$$

$$\simeq \frac{0.5}{3}\left[\sqrt{1+1} + 4\sqrt{e^{-1}+1} + \sqrt{e^{-2}+1}\right]$$

$$L \simeq 1.1929983 04$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

The Multiple Applications of Simpson's $\frac{1}{3}$-Rule:

n should be even, i.e $n = 2m$, $m \in \mathbb{N}$



$$h = \frac{b-a}{n}$$

$$\int_a^b f(x)dx \simeq \frac{h}{3}\left[f(x_0) + 4f(x_1) + f(x_2)\right] + \frac{h}{3}\left(f(x_2) + 4f(x_3) + f(x_4)\right)$$

$$+ \frac{h}{3}\left(f(x_4) + 4f(x_5) + f(x_6)\right) + \cdots + \frac{h}{3}\left(f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)\right)$$

$$\Rightarrow \int_a^b f(x)dx = \frac{h}{3}\left[f(x_0) + 4f(x_1) + f(x_2) + f(x_2) + 4f(x_3) + f(x_4) + f(x_4) + 4f(x_5)\right.$$

$$\left. + f(x_6) + \cdots + f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)\right]$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$\Rightarrow \int_a^b f(x)\,dx = \frac{h}{3}\Big[f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) + 4f(x_5) + 2f(x_6) + \cdots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)\Big]$$

$$\Rightarrow \int_a^b f(x)\,dx = \frac{h}{3}\Big[f(x_0) + 4\big(f(x_1) + f(x_3) + \cdots + f(x_{n-1})\big) + 2\big(f(x_2) + f(x_4) + \cdots + f(x_{n-2})\big) + f(x_n)\Big]$$

$$\int_a^b f(x)\,dx \doteq \frac{h}{3}\Big[f(x_0) + 4\sum_{i=1}^{m} f(x_{2i-1}) + 2\sum_{i=1}^{m-1} f(x_{2i}) + f(x_n)\Big]$$

$n = 2m$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

e.g : Using Multiple - Application Simpson's $\frac{1}{3}$-rule

taking $n=6$, approximate $\int_0^3 x^2 dx$.

Sol:

$n=6$, $f(x)=x^2$, $a=0$, $b=3$

$$\Longrightarrow h = \frac{3-0}{6} = 0.5$$

$x_0=0$, $x_1=0.5$, $x_2=1$, $x_3=1.5$, $x_4=2$, $x_5=2.5$

$x_6=3$

$$\therefore \int_0^3 x^2 dx \cong \frac{h}{3}\left[f(x_0)+4f(x_1)+2f(x_2)+4f(x_3)+2f(x_4)+4f(x_5)+f(x_6)\right]$$

$$\cong \frac{0.5}{3}\left[0^2+4(0.5^2)+2(1^2)+4(1.5^2)+2(2^2)+4(2.5^2)+3^2\right]$$

$$\cong 9$$

True value $\int_0^3 x^2 dx = \frac{x^3}{3}\Big|_0^3 = 9$

END

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Simple Simpson's $\frac{3}{8}$ - Rule : $(n=3)$

To approximate the integral $\int_a^b f(x)\,dx$ using this method, we replace the integrand $f(x)$ by cubic function $f_3(x)$ (using Lagrange polynomial of order 3)

$$\Rightarrow \int_a^b f(x)\,dx \cong \int_a^b f_3(x)\,dx$$

Notice that

$n=3 \rightarrow h=\frac{b-a}{3}$

$x_0 = a,\ x_1 = a+h,\ x_2 = a+2h$

$x_3 = a+3h = b$

After Calculations

$$\Rightarrow \int_a^b f(x)\,dx = \frac{3h}{8}\left[f(x_0) + 3f(x_1) + 3f(x_2) + f(x_3)\right]$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Apply Simple Simpson's $\frac{3}{8}$-rule to approximate $\int_0^1 \sec\sqrt{x}\,dx$

Sol: $n=3$, $a=0$, $b=1 \longrightarrow h=\frac{b-a}{n}=\frac{1-0}{3}=\frac{1}{3}$

$\Rightarrow x_0=0$, $x_1=\frac{1}{3}$, $x_2=\frac{2}{3}$ and $x_3=1$.

$$f(x)=\sec\sqrt{x}=\frac{1}{\cos\sqrt{x}}$$

$$\Rightarrow \int_0^1 \sec\sqrt{x}\,dx \cong \frac{3h}{8}\left[f(x_0)+3f(x_1)+3f(x_2)+f(x_3)\right]$$

$$\cong \frac{3(\frac{1}{3})}{8}\left[\frac{1}{\cos\sqrt{0}}+\frac{3}{\cos\sqrt{\frac{1}{3}}}+\frac{3}{\cos\sqrt{\frac{2}{3}}}+\frac{1}{\cos\sqrt{1}}\right]$$

$$\cong 1.35154 \quad \checkmark$$

True value (by calculator) 1.351069947.

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Multiple - Application Simpson's $\frac{3}{8}$ - Rule

$n = 3m \quad , m \in \mathbb{N} \quad (n \equiv \text{multiple of } 3)$



$$\int_a^b f(x)dx = \frac{3h}{8}\left(f(x_0)+3f(x_1)+3f(x_2)+f(x_3)\right) + \frac{3h}{8}\left(f(x_3)+3f(x_4)+3f(x_5)+f(x_6)\right)$$

$$+ \quad \cdots \quad + \quad \frac{3h}{8}\left(f(x_{n-3})+3f(x_{n-2})+3f(x_{n-1})+f(x_n)\right)$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

$$\Rightarrow \int_a^b f(x)\,dx = \frac{3h}{8}\left[f(x_0) + 3f(x_1) + 3f(x_2) + 2f(x_3) + 3f(x_4) + 3f(x_5) + 2f(x_6) + \cdots + 2f(x_{n-3}) + 3f(x_{n-2}) + 3f(x_{n-1}) + f(x_n)\right]$$

$$n = 6 \rightarrow \int_a^b f(x)\,dx = \frac{3h}{8}\left[f(x_0) + 3f(x_1) + 3f(x_2) + 2f(x_3) + 3f(x_4) + 3f(x_5) + f(x_6)\right]$$

---

use Simpson's rules if

$n = 5$

$$\Rightarrow \int_a^b f(x)\,dx = \frac{3h}{8}\left[f(x_0) + 3f(x_1) + 3f(x_2) + f(x_3)\right] + \frac{h}{3}\left[f(x_3) + 4f(x_4) + f(x_5)\right]$$



- END -

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Numerical Differentiation.

Notations: If $y = f(x)$



Taylor series: about $x = x_i$

$$f(x) = f(x_i) + f'(x_i)(x - x_i) + \frac{f''(x_i)(x - x_i)^2}{2!} + \frac{f'''(x_i)(x - x_i)^3}{3!} + \cdots \quad \cdots (1)$$

* If $x = x_{i+1} \rightarrow x - x_i = x_{i+1} - x_i = h$

$$\Rightarrow f(x_{i+1}) = f(x_i) + f'(x_i)h + \frac{f''(x_i)h^2}{2!} + \frac{f'''(x_i)h^3}{3!} + \cdots \quad \cdots (2)$$

* If $x = x_{i-1} \rightarrow x - x_i = x_{i-1} - x_i = -h$

sub. in (1)

$$f(x_{i-1}) = f(x_i) - f'(x_i)h + \frac{f''(x_i)h^2}{2!} - \frac{f'''(x_i)h^3}{3!} + \cdots \quad (3)$$

* If $x = x_{i+2} \rightarrow x_{i+2} - x_i = x - x_i = 2h$

$$\Rightarrow f(x_{i+2}) = f(x_i) + f'(x_i)(2h) + \frac{f''(x_i)(4h^2)}{2!} + \cdots \quad (4)$$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

from ② $\Rightarrow$

$$f'(x_i)\,h = f(x_{i+1}) - f(x_i) - \frac{f''(x_i)h^2}{2!} - \frac{f'''(x_i)h^3}{3!} - \cdots$$

$\div h \Longrightarrow$

$$f'(x_i) = \frac{f(x_{i+1}) - f(x_i)}{h} \underbrace{- \frac{f''(x_i)h}{2} - \frac{f'''(x_i)h^2}{6} - \cdots}_{-O(h)}$$

$$\Rightarrow f'(x_i) = \frac{f(x_{i+1}) - f(x_i)}{h} - O(h)$$

or

$$\boxed{f'(x_i) \cong \frac{f(x_{i+1}) - f(x_i)}{h}}$$



from ③ $\Rightarrow$

$$f'(x_i)\,h = f(x_i) - f(x_{i-1}) + \frac{f''(x_i)h^2}{2!} - \frac{f'''(x_i)h^3}{3!} + \cdots$$

$\div h \Longrightarrow$

$$f'(x_i) = \frac{f(x_i) - f(x_{i-1})}{h} + \frac{f''(x_i)h}{2!} - \frac{f'''(x_i)h^2}{2!} + \cdots$$

$$\Rightarrow f'(x_i) = \frac{f(x_i) - f(x_{i-1})}{h} + O(h)$$

$$\boxed{f'(x_i) \cong \frac{f(x_i) - f(x_{i-1})}{h}}$$



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

(2) − (3) ⟹

$$f(x_{i+1}) - f(x_{i-1}) = 2f'(x_i)h + 2\frac{f'''(x_i)h^3}{3!} + 2\frac{f^{(5)}(x_i)h^5}{5!} + \cdots$$

$\div 2h$

$$\Rightarrow \frac{f(x_{i+1}) - f(x_{i-1})}{2h} = f'(x_i) + \frac{f'''(x_i)h^2}{3} + \frac{f^{(5)}(x_i)h^4}{5!} + \cdots$$

$$\Rightarrow f'(x_i) = \frac{f(x_{i+1}) - f(x_{i-1})}{2h} - \frac{f'''(x_i)h^2}{3} - \frac{f^{(5)}(x_i)h^4}{5!} - \cdots$$

$$\rightarrow f'(x_i) = \frac{f(x_{i+1}) - f(x_{i-1})}{2h} - O(h^2)$$

or

$$f'(x_i) \cong \frac{f(x_{i+1}) - f(x_{i-1})}{2h}$$



To approximate the first derivative we can use

1) $f'(x_i) \cong \frac{f(x_{i+1}) - f(x_i)}{h}$ — Forward divided Difference Formula



2) $f'(x_i) \cong \frac{f(x_i) - f(x_{i-1})}{h}$ — Backward divided Difference Formula



3) $f'(x_i) \cong \frac{f(x_{i+1}) - f(x_{i-1})}{2h}$ — Centered divided difference formula



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

(2) + (3) $\Rightarrow$

$$f(x_{i+1}) + f(x_{i-1}) = 2f(x_i) + 2\frac{f''(x_i)h^2}{2!} + 2\frac{f^{(4)}(x_i)h^4}{4!} + \cdots$$

$$\Rightarrow f''(x_i)h^2 = f(x_{i+1}) - 2f(x_i) + f(x_{i-1}) - \frac{f^{(4)}(x_i)h^4}{12} - \cdots$$

$$\div h^2 \Rightarrow f''(x_i) = \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1})}{h^2} - \frac{f^{(4)}(x_i)h^2}{12} - \cdots$$

$$\Rightarrow f''(x_i) = \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1})}{h^2} - O(h^2)$$

$$\Rightarrow \boxed{f''(x_i) \cong \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1})}{h^2}}$$

To approximate $f''(x_i)$ we can use

1) $f''(x_i) \cong \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1})}{h^2}$ centered finite divided difference formula



2) $f''(x_i) \cong \frac{f(x_{i+2}) - 2f(x_{i+1}) + f(x_i)}{h^2}$ Forward finite divided difference formula



3) $f''(x_i) \cong \frac{f(x_i) - 2f(x_{i-1}) + f(x_{i-2})}{h^2}$ Backward finite divided difference formula



Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Given that $f(x) = x^2$, approximate $f'(1)$ using Forward and centered divided difference formulas. take the step size $h = 0.1$

Sol:

$f(x) = x^2$



Forward: $f'(1) \cong \dfrac{f(x_{i+1}) - f(x_i)}{h}$

$\cong \dfrac{1.21 - 1}{0.1}$

$\cong \dfrac{0.21}{0.1} = \dfrac{21}{10} = 2.1$

True value
$f'(x) = 2x$.
$f'(1) = 2$

Centered: $f'(1) \cong \dfrac{f(x_{i+1}) - f(x_{i-1})}{2h}$

$\cong \dfrac{1.21 - 0.81}{2(0.1)}$

$\cong 2$ ..

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g: Given that $f(2.5) = 4$ and $f(3) = 5$, use a suitable divided difference formula to approximate $f'(3)$?

Sol:



$$f'(3) = \frac{f(x_i) - f(x_{i-1})}{h}$$

$h = 3 - 2.5 = 0.5$

$$= \frac{5 - 4}{0.5} = \frac{1}{0.5}$$

$$= 2$$

e.g: Given that $f(x) = x^3$, approximate $f''(2)$ use centered divided difference formula take the step size to be 0.5 ?

$h = 0.5$

$x_i = 2$

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

Sol: $f(x) = x^3$

(7)

$f(x_{i-1}) = 3.375$ $\quad$ $f(x_i) = 8$ $\quad$ $f(x_{i+1}) = 15.625$

$x_{i-1} = 1.5$ $\quad$ $x_i = 2$ $\quad$ $x_{i+1} = 2.5$

$$f''(2) \cong \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1})}{h^2}$$

$$\cong \frac{15.625 - 2(8) + 3.375}{0.5^2}$$

$$f''(2) \cong 12$$

True value $= f'(x) = 3x^2 \Rightarrow f'(2) = 12$

Also $f''(2) = 12$

Error $= 0$;

END.

Scanned with CamScanner

الممسوحة ضوئيا بـ CamScanner

Euler's Method

Introduction to Differential Equations:

Def:

A differential Equation (D.E) is an equation that contaiNts derivative(s) of unknown function, and its order is the order of the highest derivatives that it contains.

To solve a D.E is to find the unknown function that satisfies it.

Note: If the differential equation contains partial derivative(s), it is called partial differential equation (PDE)

but if it contains ordinary derivative(s) it is called ordinary differential equation (ODE)

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

Examples:

* $y'' + xy' + y = e^x$ ODE of 2nd order

* $\frac{\partial^2 u}{\partial y \partial x} + \frac{\partial^2 u}{\partial x^2} = 0$ PDE of 2nd order

* $x\,dy + y\,dx = 0$ 1st D.E

* $y^{(4)} + y^5 = 0$ 4th order D.E

* $(x^2 + y^2)\,dy + 2xy\,dx = 0$ 1st order D.E

* $y''' = \cos x$ D.E of 3rd order

* $f'(x) = 6x + 1$ D.E of 1st order

* $u_t = 9u_{xx}$ PDE of 2nd order. (Heat equation).

Scanned with CamScanner
الممسوحة ضوئيا بـ CamScanner

e.g.: Show that $y = \sin 2x + \cos 2x$ is a solution for the differential equation

$$y'' + 4y = 0$$

Soln:

$$y = \sin 2x + \cos 2x$$

$$y' = 2\cos 2x - 2\sin 2x$$

$$y'' = -4\sin 2x - 4\cos 2x$$

$$y'' + 4y = -4\sin 2x - 4\cos 2x + 4(\sin 2x + \cos 2x)$$

$$= -4\sin 2x - 4\cos 2x + 4\sin 2x + 4\cos 2x$$

$$= 0$$

$\therefore y = \sin 2x + \cos 2x$ satisfies the D.E.

Hence, it is a solution for the D.E.

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

e.g: Solve the differential equation $\frac{dy}{dx} = 2x$, $y(0) = 4$

Sol:

$\Rightarrow dy = 2x\,dx$

$\Rightarrow \int dy = \int 2x\,dx$

$y = x^2 + C$ general solution

, $y(0) = 4$ ← condition

$y(0) = 4 \rightarrow 0 + C = 4$

$\rightarrow C = 4$

$\therefore y = x^2 + 4$



Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

D.E + condition(s) at the same value of the independent variable ⟶ Initial Value Problem (IVP)

$$y'' + y = 0 \quad , \quad y(0) = 5 \; , \quad y'(0) = 4$$

$$y' = 2x + 1 \quad , \quad y(1) = 7$$

D.E + conditions at different values of the independent variable(s) ⟶ Boundary Value Problem (BVP)

$$y'' + y = 0 \quad , \quad y(0) = 2 \; , \quad y(\pi) = 7$$

$$u_{xx} + u_{yy} = 0$$

$$u(x,0) = 0 \; , \quad u(x,L) = 0$$

$$u(0,y) = 0 \; , \quad u(L,y) = f(y)$$



Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Euler's Method:

To solve (approximation) the IVP $\frac{dy}{dx} = f(x,y)$, $y(x_0) = y_0$ by Euler's method

$$\frac{y_{i+1} - y_i}{h} = f(x_i, y_i)$$

$$\Longrightarrow \boxed{y_{i+1} = y_i + h\, f(x_i, y_i)}$$

$x_0 \longrightarrow y_0$

$+h \downarrow \quad x_1 \quad y_1$

$+h \downarrow \quad x_2 \quad y_2$

$+h \downarrow \quad x_3 \quad y_3$

$\vdots$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Using Euler's method, Solve the D.E (IVP)

$y' = 2x$, $y(0) = 1$ over the interval $[0, 4]$, take $h = 1$ (step size)

then find the analytic solution of this IVP.

Sol:

$y' = 2x$, $[0, 4]$

$y_{i+1} = y_i + h(2x_i)$, $y(0) = 1$

$x_0 = 0$	$y_0 = 1$
$x_1 = 1$	$y_1 = 1$
$x_2 = 2$	$y_2 = 3$
$x_3 = 3$	$y_3 = 7$
$x_4 = 4$	$y_4 = 13$

$h = 1 \rightarrow$

$y_{i+1} = y_i + 2x_i$

$y_1 = y_0 + 2x_0$

$= 1 + 2(0) = 1$

$y_2 = y_1 + 2x_1$

$y_2 = 1 + 2(1) = 3$

$y_3 = y_2 + 2x_2$

$y_3 = 3 + 2(2) = 7$

$y_4 = y_3 + 2x_3$

$= 7 + 2(3)$

$= 13$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$\Rightarrow$ (0,1) , (1,1) , (2,3) , (3, 7) , (4, 13)



Analytic solution:

$y' = 2x$, $y(0) = 1$

$\Rightarrow dy = 2x\,dx$

$\Rightarrow \int dy = \int 2x\,dx$

$y = x^2 + C$

$y(0) = 1 \rightarrow C = 1$

$\Rightarrow y = x^2 + 1$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Recall ; Parachutist Problem:

$$\frac{dv}{dt} = g - \frac{c}{m}v \qquad v(0)=0$$



$$mg - cv = m\frac{dv}{dt}$$

$$\Rightarrow \frac{dv}{dt} = g - \frac{c}{m}v$$

Analytic solution

$$v = \frac{gm}{c}\left(1 - e^{-\frac{c}{m}t}\right)$$

Ex: A parachutist of mass 68.1 kg, jumps out of a stationary hot air baloon. if the drag coefficient $c = 12.5$ kg/s $g = 9.81\ m/s^2$

then $\frac{dv}{dt} = 9.81 - \frac{12.5}{68.1}v$

$$v = \frac{(9.81)(68.1)}{12.5}\left(1 - e^{-\frac{12.5}{68.1}t}\right)$$

Complete the following table

Sol:

Notice that

$$v_{i+1} = v_i + h\left(9.81 - \frac{12.5}{68.1}v_i\right)$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$V = \frac{(9.81)(68.1)}{12.5}\left(1 - e^{-\frac{12.5}{68.1}(t)}\right)$$

$$V_{i+1} = V_i + 2\left(9.81 - \frac{12.5}{68.1} V_i\right)$$

t (sec)	V_i (Euler)	V (analytic)
0	0	0
2		
4		
6		
8		
10		
12		
14		
16		
18		
20		

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Use Euler's method to approximate $y(1.6)$ by solving the IVP

$$\frac{dy}{dx} = x^2 + y^2 \quad , y(1) = 2$$

take the step size to be 0.2

Solution:

$$\frac{dy}{dx} = x^2 + y^2 \quad , y(1) = 2$$

$$\Rightarrow f(x,y) = x^2 + y^2 \quad \text{(slope)} \quad , h = 0.2$$

$$\Rightarrow \boxed{y_{i+1} = y_i + 0.2(x_i^2 + y_i^2)}$$

$y(1.6) \cong ??$

i

x_i	y_i
$x_0 = 1$	$y_0 = 2$
$x_1 = 1.2$	$y_1 = 3$
$x_2 = 1.4$	$y_2 = 5.088$
$x_3 = 1.6$	$y(1.6) = 10.657$ 5488 $= y_3$

END

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

Runge-Kutta 2nd order Methods:

To solve (numerically) the IVP

$$\frac{dy}{dx} = f(x,y) \quad , y(x_0) = y_0$$

Find

1) $K_1 = f(x_i, y_i)$

2) $K_2 = f(x_i + ph \ , \ y_i + ph K_1)$ h: step size

where $a_1 + a_2 = 1$

$a_2 p = \frac{1}{2}$

then apply recurrence formula

$$y_{i+1} = y_i + h(a_1 K_1 + a_2 K_2) \quad i = 0, 1, 2, \ldots$$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

① Heun's Method:

$a_2 = \frac{1}{2}$

(Predictor - Corector)

$a_2 = \frac{1}{2} \longrightarrow a_1 + \frac{1}{2} = 1 \longrightarrow a_1 = \frac{1}{2}$

$a_2 p = \frac{1}{2} \longrightarrow \frac{1}{2} p = \frac{1}{2} \longrightarrow p = 1$

$\Rightarrow k_1 = f(x_i, y_i)$

$k_2 = f(x_i + h, y_i + k_1 h)$ ← Predictor

$$y_{i+1} = y_i + h\left(\frac{1}{2}k_1 + \frac{1}{2}k_2\right)$$

or $$y_{i+1} = y_i + \frac{h}{2}(k_1 + k_2)$$

Corrector

$i = 0, 1, 2, 3, \ldots$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

(2) Midpoint Method :

$a_2 = 1$

$a_2 = 1 \longrightarrow a_1 + 1 = 1 \longrightarrow a_1 = 0$

$a_2 p = \frac{1}{2} \longrightarrow 1p = \frac{1}{2} \longrightarrow p = \frac{1}{2}$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \tfrac{1}{2}h, y_i + \tfrac{1}{2}h k_1)$$

$$y_{i+1} = y_i + h k_2$$

$i = 0, 1, 2, 3, \ldots$

(3) Ralston Method :

$a_2 = \frac{2}{3}$

$a_1 + a_2 = 1 \longrightarrow a_1 + \frac{2}{3} = 1 \longrightarrow a_1 = \frac{1}{3}$

$a_2 p = \frac{1}{2} \longrightarrow \frac{2}{3} p = \frac{1}{2} \longrightarrow p = \frac{3}{4}$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + \tfrac{3}{4}h, y_i + \tfrac{3}{4}h k_1)$$

$$y_{i+1} = y_i + h\left(\tfrac{1}{3}k_1 + \tfrac{2}{3}k_2\right)$$

$i = 0, 1, 2, \ldots$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: Use Heun's method to solve the (IVP)

$$\frac{dy}{dx} = y\sin x \quad , \quad y(0) = 1$$

over the interval $[0, 1]$, assume the step size $h = 0.5$.

Sol: $f(x, y) = y\sin x$

$x_0 = 0 \; , \; y_0 = 1 \; , \; h = 0.5$

$x_0 = 0 \quad y_0 = 1$

$x_1 = 0.5 \quad y_1$

$x_2 = 1 \quad y_2 = ?$

1st iteration:

$x_0 = 0 \; , \; y_0 = 1$

$K_1 = f(x_i, y_i) \longrightarrow K_1 = y_0 \sin x_0$

$K_1 = 1 \sin 0 \rightarrow \boxed{K_1 = 0}$

$K_2 = f(x_0 + \overset{h}{0.5} \; , \; y_0 + K_1 (\overset{h}{0.5}))$

$K_2 = f(0.5, \; 1 + 0(0.5))$

$K_2 = f(0.5, 1.0) = 1.0 \sin(0.5) \rightarrow \boxed{K_2 = 0.4794255}$

$y_1 = y_0 + h(\frac{1}{2}K_1 + \frac{1}{2}K_2)$

$y_1 = 1 + \frac{0.5}{2}(0 + 0.4794255)$

$\rightarrow \boxed{y_1 = 1.1198564}$

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

$x_1 = 0.5$, $y_1 = 1.1198564$

The analytic solution: $y = e^{1-\cos x}$

$y(0.5) = e^{1-\cos(0.5)} = 1.1302258$

2nd iteration

$x_1 = 0.5$, $y_1 = 1.1198564$

$K_1 = f(0.5, 1.1198564)$

$= 1.1198564 \sin 0.5$

$\Rightarrow \boxed{K_1 = 0.5368878}$

$K_2 = f(0.5 + 0.5, 1.1198564 + 0.5(0.5368878))$

$K_2 = f(1, 1.3883003)$

$= 1.3883003 \sin 1$

$\boxed{K_2 = 1.1682442}$

$\therefore y_2 = 1.1198564 + \frac{0.5}{2}[0.5368878 + 1.1682442]$

$y_2 = 1.5461394$

True value

$y(1) = e^{1-\cos 1}$

$= 1.583595183$

(0, 1), (0.5, 1.1198564) and (1, 1.5461394)

Ex: Solve the above example by Midpoint and Ralston methods

- END -

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

Numerical Solution for BVP

by

Finite Difference Method:

Idea: replace

y'' by $\frac{y_{i+1} - 2y_i + y_{i-1}}{h^2}$	centered F.D formula for 2nd derivative
y' by $\frac{y_{i+1} - y_{i-1}}{2h}$	center F.D formula for 1st derivative
y by y_i	

$h = \Delta x$ = step size

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

e.g: By using finite difference method, solve the BVP:

$$y'' - y = 0 \quad , \quad y(0) = 1 \quad , \quad y(1) = 2$$

Use 4-segments (5 point (values)(nodes))

Sol:

$$n = 4 \quad , \quad a = 0 \quad , \quad b = 1$$

$$h = \frac{b-a}{n} \rightarrow h = \frac{1-0}{4} \longrightarrow h = 0.25$$

$$\Rightarrow a = x_0 = 0 \quad , x_1 = 0.25 \quad , x_2 = 0.5 \quad , x_3 = 0.75 \quad , x_4 = 1 = b$$



Now replace y'' by $\frac{y_{i+1} - 2y_i + y_{i-1}}{h^2}$ and y by y_i in the D.E $y'' - y = 0$, to get

$$\frac{y_{i+1} - 2y_i + y_{i-1}}{h^2} - y_i = 0$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$* h^2 \Rightarrow \quad y_{i+1} - 2y_i + y_{i-1} - h^2 y_i = 0$

$$y_{i-1} - (2+h^2)y_i + y_{i+1} = 0$$

but $h^2 = 0.125^2 = 0.0625$

$$\Rightarrow \boxed{y_{i-1} - 2.0625 y_i + y_{i+1} = 0 \qquad i = 1, 2, 3}$$

$i=1 \rightarrow y_0 - 2.0625 y_1 + y_2 = 0$

$\Rightarrow 1 - 2.0625 y_1 + y_2 = 0$

$$\Rightarrow \boxed{-2.0625 y_1 + y_2 = -1} \qquad (1)$$

$$i=2 \rightarrow \boxed{y_1 - 2.0625 y_2 + y_3 = 0} \qquad (2)$$

$i=3 \longrightarrow y_2 - 2.0625 y_3 + y_4 = 0$

$\rightarrow y_2 - 2.0625 y_3 + 2 = 0$

$$\Rightarrow \boxed{y_2 - 2.0625 y_3 = -2} \qquad (3)$$

$$\begin{bmatrix} -2.0625 & 1 & 0 \\ 1 & -2.0625 & 1 \\ 0 & 1 & -2.0625 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ -2 \end{bmatrix}$$

by solving this system we get

$$y_1 = 1.130193 \;, \quad y_2 = 1.331023 \;, \quad y_3 = 1.615041$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

The analytic solution of the BVP

$$y = 0.69440\, e^{x} + 0.30560\, e^{-x}$$

$$y(0) = y_0 = 1$$

$1.12962876 = y(0.25) \cong y_1 = 1.130193$

$1.33022782 = y(0.5) \cong y_2 = 1.331023$

$1.61440003 = y(0.75) \cong y_3 = 1.615041$

$1.99999859 = y(1) \cong y_4 = 2$

2

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner

Application:

A noninsulated uniform rod positioned between two bodies of constant but different temperature.

If L is non-insulated along its length and the system in the steady state the equation



$$\frac{d^2T}{dx^2} + h'(T_a - T) = 0$$

$$T(0) = A \ , \ T(L) = B$$

h': heat transfer coefficient

T_a: the temperature of the surroundling.

$T = T(x)$ = the temperature at x.

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

e.g: If the system in the steady state. and the differential equation of this model:



$$\frac{d^2T}{dx^2} + h'(T_a - T) = 0$$

$T(0) = 40^\circ C$, $T(10) = 200^\circ C$

and the ~~trans~~ heat transfer coefficient $= 0.01\ m^{-2}$

and the surrounding temperature $= 20^\circ C$

Find the temperature when $x = 2, 4, 6, 8$ m

using finite difference method.

Sol $h = 2$ (we have 5-segments)

The diff. equation

$$\frac{d^2T}{dx^2} + 0.01(20 - T) = 0 \quad , T(0) = 40,$$

$$T(10) = 200$$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$\frac{d^2T}{dx^2} + 0.2 - 0.01T = 0$$

$$\Rightarrow \frac{d^2T}{dx^2} - 0.01T = -0.2$$

$$\Rightarrow \frac{T_{i+1} - 2T_i + T_{i-1}}{h^2} - 0.01\,T_i = -0.2$$

but $h = 2 \rightarrow h^2 = 4$

$$\Rightarrow \frac{T_{i+1} - 2T_i + T_{i-1}}{4} - 0.01\,T_i = -0.2$$

$$*4 \Rightarrow T_{i+1} - 2T_i + T_{i-1} - 0.04\,T_i = -0.8$$

$$\boxed{T_{i-1} - 2.04\,T_i + T_{i+1} = -0.8}$$

$T(0) = 40, \quad T(10) = 200$



4 - unknowns

$\Rightarrow i = 1, 2, 3, 4$

Scanned with CamScan
الممسوحة ضوئيا بـ CamScanner

$$T_{i-1} - 2.04\,T_i + T_{i+1} = -0.8$$

$i=1 \rightarrow T_0 - 2.04T_1 + T_2 = -0.8$

$40 - 2.04T_1 + T_2 = -0.8$

$$-2.04T_1 + T_2 = -40.8 \quad -(1)$$

$i=2 \rightarrow$ $$T_1 - 2.04T_2 + T_3 = -0.8 \quad -(2)$$

$i=3 \rightarrow$ $$T_2 - 2.04T_3 + T_4 = -0.8 \quad -(3)$$

$i=4 \rightarrow T_3 - 2.04T_4 + T_5 = -0.8$

$\rightarrow T_3 - 2.04T_4 + 200 = -0.8$

$\Rightarrow$ $$T_3 - 2.04T_4 = -200.8 \quad (4)$$

$$\begin{bmatrix} -2.04 & 1 & 0 & 0 \\ 1 & -2.04 & 1 & 0 \\ 0 & 1 & -2.04 & 1 \\ 0 & 0 & 1 & -2.04 \end{bmatrix} \begin{bmatrix} T_1 \\ T_2 \\ T_3 \\ T_4 \end{bmatrix} = \begin{bmatrix} -40.8 \\ -0.8 \\ -0.8 \\ -200.8 \end{bmatrix}$$

Solving this system we get

$T_1 = 65.9698°C$; $T_2 = 93.7785°C$, $T_3 = 124.5382°C$

$T_4 = 159.4795°C$

END

Scanned with CamScan

الممسوحة ضوئيا بـ CamScanner