

From: Scientific Perspectives™
<education@scientificperspectives.com>
To: [\[HCP email\]](#)
Subject 1: What percentage of patients do not respond to oral antidepressants alone?
Subject 2: Did you know this about medication response in patients with MDD?
Alternate Subject Line 1: Are your patients struggling to find relief from MDD?
Alternate Subject Line 2: Do you treat patients with MDD cycling through oral antidepressants?
Pre-header 1: Test your knowledge now
Pre-header 2: Uncover information from the STAR*D trial

Information from Industry
Developed under the direction and sponsorship of Johnson & Johnson

Are your patients struggling to find relief from major depressive disorder (MDD)?



What Do You Think?

Based on the STAR*D trial, what percentage of patients with MDD do not respond to oral antidepressants alone?¹

- ☒ A. 33%
- ☐ B. 15%
- ☐ C. 17%
- ☐ D. 25%

The STAR*D trial enrolled adult patients aged 18 to 75 diagnosed with a current major depressive episode who were treated in 41 participating psychiatric and primary care sites. Patients were evaluated across 4 successive steps of therapy, either switching or augmenting treatment if an adequate response was not achieved.¹

STAR*D=Sequenced Treatment Alternatives to Relieve Depression.

Reference:

1. Rush AJ, Trivedi MH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one of several treatment steps: a STAR*D report. *Am J Psychiatry*. 2006;163(11):1905-1917. doi:10.1176/ajp.2006.163.11.1905

© Johnson & Johnson, Inc. 2024 09/24 cp-470066v1

This is a service from Relevate Health.
To ensure you continue to receive these emails, please add education@scientificperspectives.com to your address book.
Your participation in this activity is governed by the Relevate Health [Privacy Policy](#) and [Terms of Use](#).

[Unsubscribe](#)