

Simplified INLA, automatic differentiation and adaptive Gauss-Hermite quadrature for fast and accurate approximate inference

Integrated nested Laplace approximations for extended latent Gaussian models, with application to the Naomi HIV model

Adam Howes^{1, 2}
@adamhowes
ath19@ic.ac.uk

Alex Stringer³ Seth R. Flaxman⁴ Jeffrey W. Eaton²

¹ Department of Mathematics, Imperial College London
² MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London
³ Department of Statistics and Actuarial Science, University of Waterloo
⁴ Department of Computer Science, University of Oxford

The Naomi model

Inference procedure
Laplace approximation
Adaptive Gauss-Hermite Quadrature

Approximate integrals by

$$\int_{\Theta} p(\theta) d\theta \approx |L| \sum_{z \in \mathcal{Q}(m,k)} p(\hat{\theta} + Lz) \omega(z)$$

with Gauss-Hermite quadrature rule $z \in \mathcal{Q}(m, k)$ adapted based upon the mode $\hat{\theta} = \operatorname{argmax}_{\theta \in \Theta} p(\theta)$ and lower Cholesky $LL^\top = -\partial_\theta^2 \log p(\theta)|_{\theta=\hat{\theta}}$ of the target.

Our algorithm

Given C++ user template for $-\log p(y, x, \theta)$:

Comparison
Conclusions

Funding AH was supported by the EPSRC and Bill & Melinda Gates Foundation. This research was supported by the MRC Centre for Global Infectious Disease Analysis.

References