


Time stepping review of open-source solvers

Guided research

Marc Amorós Trepát

Supervisor: Prof. Dr. Hans-Joachim Bungartz

Advisor: M.Sc. Benjamin Rodenberg



Outline

- 1 Introduction
 - Motivation
 - Open-Source solvers
 - Time stepping schemes
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions

- Simulations of physical phenomena are useful in many fields.

- Simulations of physical phenomena are useful in many fields.
- Can we reduce the error of solutions?

- Simulations of physical phenomena are useful in many fields.
- Can we reduce the error of solutions?
- Higher order of a numerical method \implies higher precision?

- Simulations of physical phenomena are useful in many fields.
- Can we reduce the error of solutions?
- Higher order of a numerical method \implies higher precision?
- How does this error behave when we couple simulations?




Figure 1 Diagram showing setup for coupling simulations with preCICE library.

Source: <https://precice.org>

Time stepping schemes

We focus on temporal discretization error $\varepsilon_{\Delta t}$.

Second-order time stepping schemes $\rightarrow \varepsilon_{\Delta t}$ decreases $\mathcal{O}(\Delta t^2)$.

We focus on temporal discretization error $\varepsilon_{\Delta t}$.

Second-order time stepping schemes $\rightarrow \varepsilon_{\Delta t}$ decreases $\mathcal{O}(\Delta t^2)$.

When solving a PDE of the form:

$$\frac{\partial u}{\partial t} = F(u, t) \quad (1)$$

we need to discretize the time-derivative. Easiest way is the Euler explicit method:

$$\frac{u^{n+1} - u^n}{\Delta t} = F(u^n, t^n) \quad (2)$$

OpenFOAM

- Implicit Euler (1st order):

$$\frac{u^{n+1} - u^n}{\Delta t} = \frac{1}{2} [F(u^n, t^n) + F(u^{n+1}, t^{n+1})] \quad (3)$$

- Crank-Nicolson (2nd order):

$$\frac{u^{n+1} - u^n}{\Delta t} = \frac{\theta}{2} F(u^n, t^n) + \left(1 - \frac{\theta}{2}\right) F(u^{n+1}, t^{n+1}) \quad (4)$$

OpenFOAM

- Implicit Euler (1st order):

$$\frac{u^{n+1} - u^n}{\Delta t} = \frac{1}{2} [F(u^n, t^n) + F(u^{n+1}, t^{n+1})] \quad (3)$$

- Crank-Nicolson (2nd order):

$$\frac{u^{n+1} - u^n}{\Delta t} = \frac{\theta}{2} F(u^n, t^n) + \left(1 - \frac{\theta}{2}\right) F(u^{n+1}, t^{n+1}) \quad (4)$$

CalculiX

- Implicit α -method (2nd order):

$$\mathbf{v}^{n+1} = \mathbf{v}^n + \Delta t [(1 - \gamma)\mathbf{a}^n + \gamma\mathbf{a}^{n+1}] \quad (5)$$

$$\mathbf{u}^{n+1} = \mathbf{u}^n + \Delta t \mathbf{v}^n + \frac{1}{2} (\Delta t)^2 [(1 - 2\beta)\mathbf{a}^n + 2\beta\mathbf{a}^{n+1}] \quad (6)$$

- 1 Introduction
- 2 OpenFOAM**
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions

The error has the form $\varepsilon_u = \varepsilon_{\Delta t} + \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$.






Figure 2 Visualization of the Taylor-Green vortex scenario.



(a) Error compared to reference ($\Delta t = 10^{-5}$).

$$|u - \tilde{u}| = \varepsilon_{\Delta t} + \varepsilon_{\text{num}}$$

plateau when $\varepsilon_{\Delta t} < \varepsilon_{\text{num}}$




(b) Error compared to analytical solution.

$$|u - u^*| = \varepsilon_{\Delta t} + \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$$


plateau when $\varepsilon_{\Delta t} < \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX**
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions

We use the α -method as time stepping scheme.



(a) Convergence study, showing higher-order convergence.



(b) Perpendicular elastic flap scenario.

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation**
- 5 Sources of error
- 6 Conclusions




Figure 5 Example solution of the FSI simulation, coupled with preCICE.




Figure 6 Convergence study of the coupled perpendicular flap scenario. Results with the Crank-Nicolson (CN) and the implicit Euler time stepping schemes, and using v2 and v3 of preCICE.

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error**
- 6 Conclusions

Verification of CalculiX adapter

We coupled the CalculiX adapter with a fake-fluid participant that applied a force $f^n = f_{max} \sin(t^n + \phi)$ on the tip.





Figure 7 Convergence study of the CalculiX adapter coupled with a fake fluid participant. Reference solution is $\Delta t = 10^{-4}$.

Verification of fluid participant

- Single-solver simulations with fluid participant.
- Hard to convergence due to high CFL numbers.
- Precomputed initial conditions increased stability.
- Time variable inflow conditions.
- Sublinear error convergence.



- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions**

- Does it make sense to use second order methods? Yes, but not always.

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.
- Linear convergence error on FSI simulation achievable.

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.
- Linear convergence error on FSI simulation achievable.
- CalculiX adapter maintains error convergence. OpenFOAM adapter?

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.
- Linear convergence error on FSI simulation achievable.
- CalculiX adapter maintains error convergence. OpenFOAM adapter?
- Try different setups and different numerical methods.

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.
- Linear convergence error on FSI simulation achievable.
- CalculiX adapter maintains error convergence. OpenFOAM adapter?
- Try different setups and different numerical methods.
- This work offers some insights in the error behaviour.

- Does it make sense to use second order methods? Yes, but not always.
- CalculiX and OpenFOAM can reach higher-order convergence.
- Linear convergence error on FSI simulation achievable.
- CalculiX adapter maintains error convergence. OpenFOAM adapter?
- Try different setups and different numerical methods.
- This work offers some insights in the error behaviour.
- Useful for researchers surrounding preCICE.