


Time stepping review of open-source solvers

Guided research

Marc Amorós Trepát

Supervisor: Prof. Dr. Hans-Joachim Bungartz

Advisor: M.Sc. Benjamin Rodenberg



Outline

- 1 Introduction
 - Motivation
 - Open-Source solvers
 - Time stepping schemes
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions




Figure 1 Diagram showing setup for coupling simulations with preCICE library.

Time stepping schemes

When solving a PDE of the form:

$$\frac{\partial u}{\partial t} = F(u, t) \quad (1)$$

we need to discretize the time-derivative. Easiest way is the Euler explicit method:

$$\frac{u^{n+1} - u^n}{\Delta t} = F(u^n, t^n) \quad (2)$$

We focus on second-order time stepping schemes \rightarrow temporal discretization error $\varepsilon_{\Delta t}$ decreases $\mathcal{O}(\Delta t^2)$.

- 1 Introduction
- 2 OpenFOAM**
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions

The error has the form $\varepsilon_u = \varepsilon_{\Delta t} + \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$.

We use Crank-Nicolson time stepping method.





Figure 2 Visualization of the Taylor-Green vortex scenario.


OpenFOAM - convergence study



(a) Error compared to reference ($\Delta t = 10^{-5}$).

$$|u - \tilde{u}| = \varepsilon_{\Delta t} + \varepsilon_{\text{num}}$$

plateau when $\varepsilon_{\Delta t} < \varepsilon_{\text{num}}$




(b) Error compared to analytical solution.

$$|u - u^*| = \varepsilon_{\Delta t} + \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$$


plateau when $\varepsilon_{\Delta t} < \varepsilon_{\Delta x} + \varepsilon_{\text{num}}$

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX**
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions

We use the α -method as time stepping scheme.



(a) Convergence study, showing higher-order convergence.



(b) Perpendicular elastic flap scenario.

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation**
- 5 Sources of error
- 6 Conclusions




Figure 5 Example solution of the FSI simulation, coupled with preCICE.




Figure 6 Convergence study of the coupled perpendicular flap scenario. Results with the Crank-Nicolson (CN) and the implicit Euler time stepping schemes, and using v2 and v3 of preCICE.

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error**
- 6 Conclusions

Verification of CalculiX adapter

We coupled the CalculiX adapter with a fake-fluid participant that applied a force $f^n = f_{max} \sin(t^n + \phi)$ on the tip.




Figure 7 Convergence study of the CalculiX adapter coupled with a fake fluid participant. Reference solution is $\Delta t = 10^{-4}$.

Verification of fluid participant

We ran the fluid-participant as a single-solver scenario and found:

- Difficult convergence due to high CFL numbers.
-

- 1 Introduction
- 2 OpenFOAM
- 3 CalculiX
- 4 FSI simulation
- 5 Sources of error
- 6 Conclusions**

Conclusions and future work

- Does it make sense to use second order methods? Yes, but not always.
- Single-solver simulations showed how CalculiX and OpenFOAM can reach higher-order convergence.
-