

Advanced radiation and remote sensing

Lukas Kluft, Manfred Brath, Stefan Bühler

November 15, 2019

Exercise No. 3 – Line shape

You can reuse the python script from the first exercise as a starting point for this exercise. First, copy the python script from the first exercise into the directory of exercise no. 3:

```
$ cd ~/arts-lectures/exercises/03-line_shape/  
$ cp ../01-rotational_spectra/absorption.py line_shape.py
```

The “calculate_absxsec” function within your python script calculates absorption cross sections and uses several keyword arguments as inputs, among other lineshape and forefactor. Please use “Voigt_Kuntz6” as line shape and “VVH” as fore factor:

1. Choose an individual line of for example water vapor and perform calculations over a restricted frequency range for a number of different pressures. Keep the temperature constant.

- How does the shape of the spectral lines change?

By now we investigated absorption in terms of the absorption cross-section σ . Another widely used unit is the absorption coefficient α . It takes the number concentration n of the absorber into account:

$$\alpha = n \cdot \sigma \tag{1}$$

- How does the absorption coefficient in the line centre change, if pressure is changed?
2. The full-width at half maximum (FWHM) is a measure of the line width. Typhon provides the function `typhon.spectroscopy.linewidth()` to calculate the FWHM for a given absorption spectrum.
 - Make a plot of this as a function of altitude (pressure) for a microwave line and an infrared absorption line.