

Doctoral Dissertation

博士論文

Effects of Replicated N-body Simulation Boxes in Simulating Weak Lensing Observables

(弱重力レンズ統計量の模擬データ生成における
有限体積のN体シミュレーションの影響の研究)

A Dissertation Submitted for the Degree of Doctor of Philosophy

December 2024

令和6年12月博士（理学）申請

Department of Physics, Graduate School of Science,

The University of Tokyo

東京大学大学院理学系研究科
物理学専攻

Akira Tokiwa

常盤 晟

論文の内容の要旨

Effects of Replicated N-body Simulation Boxes in Simulating Weak Lensing Observables

(弱重力レンズ統計量の模擬データ生成における
有限体積のN体シミュレーションの影響の研究)

氏名

常盤 晟

This thesis examines how the super-sample effect and the box replication effect influence the accuracy and interpretation of higher-order statistics from weak gravitational lensing data. These effects are critical in understanding the limitations and potential biases in simulations used for cosmological studies. The study begins with a review of modern cosmology. It highlights how weak lensing measures small distortions in the shapes of distant galaxies caused by intervening matter. This can help us understand dark matter, dark energy, and the overall structure of the Universe. It then focuses on beyond-two-point statistics, a method reveals non-Gaussian features in the matter distribution and thereby enhance constraints on cosmological parameters.

To achieve this, we use a simulation-based framework with two N -body simulation strategies: the BIGBOX simulation set from the HalfDome project, which has 6144^3 particles in a $3750 \text{ Mpc}/h$ box, and the TILED simulation set, composed of multiple 1024^3 -particle boxes, each $625 \text{ Mpc}/h$ in size. Both simulations share the same cosmological parameters but differ in volume coverage and box replication schemes, allowing us to isolate super-sample modes and other effects. We project three-dimensional matter distributions into full-sky convergence maps at various redshifts under the Born approximation, then incorporate shape noise and Gaussian smoothing to approximate realistic observing conditions. From these maps, we employ a Fibonacci grid to extract numerous $10^\circ \times 10^\circ$ patches and evaluate a comprehensive set of lensing observables—namely, the angular power spectrum, bispectrum, probability distribution function (PDF), peak and minima counts, and Minkowski functionals. Comparisons of the mean values, variances, and covariances of these observables between BIGBOX and TILED are further contextualized by theoretical models such as `Halofit` and `hmpdf`.

Our findings show that the mean values of all investigated statistics from BIGBOX and TILED agree at the 1% level, confirming consistency under the same cosmological framework. However, variance and covariance estimates differ by up to 10%-30% at higher multipoles and redshifts, primarily due to super-sample modes present in the larger-volume BIGBOX simulations but absent in

TILED. Although shape noise and Gaussian smoothing affect certain observables, most notably the angular power spectrum and the counts of peaks and minima, the super-sample effect remains the dominant contributor to these discrepancies. Furthermore, we detect 2%-5% systematic biases introduced by box replication artifacts, which can obscure super-sample signals by shifting the means of selected observables and altering the distribution of prominent peaks and deep minima. These results highlight several key considerations for future research and the design of upcoming weak lensing surveys.

First, large-scale modes must be incorporated into simulations to ensure realistic uncertainty estimates. We adopt the approximate dimensional analysis $R_{\sigma^2} \propto (L^2/\Omega_W)^\alpha$, where L is the simulation box size, Ω_W is the survey area, and α is an undetermined exponent. This implies that super-sample variance becomes increasingly significant for larger surveys employing relatively small simulation boxes. For instance, matching the variance levels of TILED in an 18,000 deg² survey (e.g., LSST) would require a simulation box of approximately 8 Gpc/ h .

Secondly, shape noise and Gaussian smoothing must be carefully managed to maintain the fidelity of higher-order statistics. Although these effects can shift certain observables, the super-sample effect remains robust under various observational conditions.

Finally, mitigating box replication artifacts is essential for high-precision analyses. These artifacts introduce 2%-5% biases in mean observables and can modify covariance structures by up to 10%. Nonetheless, improved simulation and data analysis strategies can address these systematic issues.

This work highlights the importance of quantifying super-sample effects, particularly in higher-order statistics, an area where no prior work existed. Additionally, it provides one of the first comprehensive investigations into the impact of box replication effect on summary statistics and their covariance. As forthcoming surveys, including the Rubin Observatory's LSST, Euclid, and the Roman Space Telescope, aim for increasingly stringent constraints, properly accounting for super-sample effects and replication artifacts will be indispensable. Future efforts will focus on reducing replication biases, exploring additional cosmological models and baryonic physics, and integrating these advances into survey planning and analysis pipelines. By refining both simulation methodologies and interpretive frameworks, this thesis helps establish weak lensing as an even more robust and informative probe of the Universe's fundamental properties.

Abstract

We investigate the influence of the super-sample effect on the mean and covariance of the power spectrum and higher-order statistics using two distinct simulation approaches: BIGBOX and TILED. Our analysis draws on 11 realizations from the state-of-the-art HalfDome project, each containing 6144^3 particles within a $3750 \text{ Mpc}/h$ volume, as well as 20 TILED realizations, each originating from smaller 1024^3 -particle boxes spanning $625 \text{ Mpc}/h$. Convergence fields are constructed under the Born approximation. To achieve uniform sky coverage, we employ a Fibonacci grid scheme to extract $10^\circ \times 10^\circ$ patches and subsequently evaluate a comprehensive set of higher-order statistics, including the bispectrum, probability distribution function (PDF), peak and minima counts, and Minkowski functionals, together with the angular power spectrum. These metrics enable a robust assessment of non-Gaussian features in convergence maps and reveal the impact of the super-sample effect.

Our comparative analysis shows that mean summary statistics from BIGBOX and TILED simulations agree to within 1% across all statistics, redshifts, and multipoles, thereby confirming both simulations' fidelity to the underlying cosmological model. In contrast, variance ratios exhibit notable discrepancies of 10%–30%, particularly at higher multipoles and redshifts, driven by large-scale modes present in BIGBOX but missing in TILED. Although shape noise affects several observables, the super-sample effect remains the dominant factor. Gaussian smoothing modifies small-scale structures but does not significantly alter covariance or correlation properties. Box replication artifacts, however, introduce additional systematic biases and increase variance, underscoring the need for careful simulation design and execution.

These findings underscore the importance of accurately modeling the super-sample effect for high-precision cosmological analyses, especially in light of upcoming weak-lensing surveys such as LSST, Euclid, and Roman. A reliable variance estimate mandates the inclusion of large-scale modes while ensuring robustness against shape noise and smoothing procedures. Mitigating systematic challenges like box replication artifacts is essential to avoid biased interpretations of higher-order statistics. Future efforts will focus on resolving these artifacts, incorporating diverse cosmological models and baryonic physics, and integrating refined approaches into survey design and data analysis pipelines to bolster cosmological parameter inference. Our study marks a critical step toward a deeper understanding of the super-sample effect and its implications for generating and analyzing next-generation weak-lensing mock data.

Acknowledgements

This research was conducted during my Ph.D. studies in the Department of Physics, Graduate School of Science, at The University of Tokyo, under the supervision of Masahiro Takada. I would like to express my deepest gratitude to him for his continuous support from my master's degree to the present. I am also grateful to Jia Liu, Adrian Bayer, Joaquin Armijo, Leander Thiele, and Ryo Terasawa for their valuable comments and discussions. Special thanks to the members of the Takada group and the researchers at the Kavli Institute for their support and encouragement. I would like to thank the staff of Kavli IPMU and The University of Tokyo for their assistance. I extend my heartfelt thanks to my wife, Yuuka, my family, and friends for their unwavering support.

I acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin and the RIKEN Center for Computational Science for providing grid resources that have significantly contributed to this work.

The HalfDome project is supported by JSPS KAKENHI Grants JP23KJ0392 (YZ), 23K13095 and 23H00107 (JL). MM and AL acknowledge support from NASA grant 21-ATP21-0145. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility located at Lawrence Berkeley National Laboratory, operated under Contract No. DE-AC02-05CH11231 using NERSC award HEP-ERCAP0023125. We acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing grid resources that have contributed to our work. This work was supported by MEXT as Program for Promoting Researches on the Supercomputer “Fugaku” (Multi-wavelength Cosmological Simulations for Next-generation Surveys, JPMXP1020230407) and used computational resources of supercomputer Fugaku provided by the RIKEN Center for Computational Science (Project ID: hp230202). We thank Julian Borrill and Reijo Kesitalo for their support in computation and data storage. The HalfDome logo is designed by James Sunseri.

Contents

1	Introduction	1
1.1	Historical Overview	1
1.2	Astronomical Surveys and Observations	2
1.3	Constraint from Weak Lensing	5
1.4	Higher-Order Statistics in Weak Lensing	6
1.5	Aim of this Thesis	9
1.5.1	Motivation: Why Not a Single Huge Simulation?	10
1.5.2	Super-Sample Effect	10
1.5.3	Box Replication Effects	11
1.5.4	Simulation Strategies	12
1.6	Structure of the Thesis	12
2	Foundational Concepts in Modern Cosmology	14
2.1	Einstein Field Equations	14
2.1.1	The Stress-Energy Tensor for a Perfect Fluid	15
2.1.2	Equations of State for Different Cosmic Components	15
2.2	FLRW Metric and the Friedmann Equations	17
2.2.1	The FLRW Metric	17
2.2.2	Derivation of the Friedmann Equations	17
2.2.3	Critical Density, Density Parameters, and H_0	18
2.2.4	Evolution of Density Parameters with Scale Factor	18
2.3	Redshift and Cosmological Distances	19
2.3.1	Luminosity Distance	20
2.3.2	Angular Diameter Distance	21
2.4	Initial Conditions and the Primordial Universe	21
2.4.1	Primordial Fluctuations and the Power Spectrum	21
2.4.2	Baryon Acoustic Oscillations (BAOs)	24
2.5	Growth of Matter Density Fluctuations	24
2.6	The Spherical Collapse Model	26

2.7	Dark Matter Halos	28
2.7.1	The Halo Mass Function: Press-Schechter	28
2.7.2	Halo Bias	29
2.7.3	Halo Density Profiles: NFW and Einasto Models	30
3	Theoretical Foundations of Weak Gravitational Lensing	31
3.1	Introduction to Weak Gravitational Lensing	31
3.1.1	Derivation of the Lens Equation	31
3.1.2	The Jacobian (Lensing) Matrix $\mathcal{A}$	33
3.2	Convergence Field	34
3.2.1	Definition of Convergence in Real and Fourier Space	34
3.2.2	Convergence from Density Contrast	35
3.3	Shear Field	36
3.3.1	Definition of Shear	36
3.3.2	E-mode and B-mode Decomposition	36
3.4	Lensing Estimation	37
3.4.1	Galaxy Ellipticity and Reduced Shear	37
3.4.2	From Shear to Convergence	38
4	Summary Statistics in Weak Lensing	40
4.1	Two-point Correlation Function	40
4.2	Convergence Power Spectrum	41
4.2.1	Derivation of the Convergence Power Spectrum	41
4.2.2	Halofit Model	43
4.3	Convergence Bispectrum	44
4.4	Probability Density Functions	44
4.4.1	Smoothing and Normalization Procedures	45
4.4.2	Practical Estimation of the Convergence PDF	45
4.4.3	hmpdf Model	46
4.5	Peak and Minimum Counts	46
4.6	Minkowski Functionals	47
4.6.1	Definition of Minkowski Functionals	47
4.6.2	Analytical Expressions in Gaussian Random Fields	49
5	Estimation of Covariance Matrix for Weak Lensing Statistics	50
5.1	The Role of the Covariance Matrix in Parameter Inference	51
5.1.1	Likelihood Functions and the Gaussian Approximation	51

5.1.2	Fisher Information and Parameter Constraints	51
5.2	Covariance of the Matter Power Spectrum	52
5.2.1	Shot Noise and Super-Sample Covariance	53
5.3	Covariance of the Angular Convergence Power Spectrum	54
5.4	Covariance of Higher-Order Weak Lensing Statistics	55
6	Numerical Simulations for Large Scale Structure and Weak Lensing	57
6.1	Introduction to Numerical Simulations in Cosmology	57
6.1.1	Dark Matter-Only vs. Hydrodynamical Simulations	57
6.1.2	Historical Evolution and Growth in Particle Number	58
6.2	Initial Condition Generation	58
6.2.1	Initial Density Field	58
6.2.2	Initial Displacement Field	59
6.2.3	Initial Velocities	60
6.3	N-Body Simulation Methods	60
6.3.1	Direct Summation	60
6.3.2	Particle-Mesh (PM) Method	61
6.3.3	Particle-Particle Particle-Mesh (P3M) Method	62
6.3.4	Tree-Particle-Mesh (Tree-PM) Method	62
6.4	Computational Tools and Optimization Techniques	63
6.4.1	Fast Fourier Transforms (FFT)	63
6.4.2	Mass Assignment Schemes	63
6.4.3	Parallelization Techniques	64
6.4.4	Adaptive Mesh Refinement (AMR)	65
6.4.5	Tree Construction	65
6.5	Advanced Codes and Methods: FASTPM	66
6.5.1	Modified Kick and Drift Factors	67
6.5.2	Algorithm Steps	68
6.6	Generating Weak Lensing Maps from Simulations	68
6.6.1	Conventional Ray-Tracing Algorithm	69
6.6.2	Born-approximated Ray-tracing	70
7	Simulation-Based Methodology for Assessing Higher-Order Weak Lensing Covariance	73
7.1	Constructing the BIGBOX and TILED Datasets	73
7.2	Producing Weak Lensing Convergence Maps at Multiple Redshifts	75
7.3	Incorporating Galaxy Shape Noise into Convergence Maps	76
7.4	Patch Selection and Projection	77

7.5	Applying Gaussian Kernels to Convergence Fields	80
7.6	Extracting Weak Lensing Statistics from Convergence Maps	81
7.6.1	Angular Power Spectrum	81
7.6.2	Bispectrum	83
7.6.3	PDF	83
7.6.4	Peak/Minima Counts	83
7.6.5	Minkowski Functionals	83
7.7	Comparing Covariances from BIGBOX and TILED	84
8	Comparative Analysis of Statistical Measures and Covariance Structures in BIGBOX and TILED	86
8.1	Comparison of Mean and Variance for Summary Statistics	86
8.2	Analysis of Covariance and Correlation Matrices	90
8.3	Effects of Noise on Summary Statistics	96
8.4	Influence of Smoothing Scale	99
8.5	Systematic Effects: Box Replication Artifact	102
9	Conclusions	108
A	Filtering Techniques in Cosmological Lensing	110
A.1	Gaussian Filter	110
A.2	Wiener Filter	110
A.3	Starlet Filter	111

List of Figures

1.1	Comparison of S_8 values measured from different surveys.	3
1.2	Forecasted constraints on Ω_m and σ_8 from the Euclid mission using ten different higher-order statistics	7
1.3	Examples of simulation strategies for generating lightcones	8
1.4	Visualization of a partial sky convergence map covering $5 \times 5 \text{ deg}^2$, showing the effects of box replication along the line of sight	9
1.5	Contributions of Gaussian, SSC, and cNG terms to the diagonal elements of the total covariance matrix for the lensing convergence power spectrum	11

2.1	Angular Power Spectrum (TT) of CMB Anisotropy	22
2.2	Linear Matter Power Spectrum by CLASS	23
3.1	Schematic representation of the lensing geometry	33
3.2	Illustration of Gravitational Lensing	35
4.1	Concept of Peaks/Minima Identification	47
4.2	Excursion sets of different thresholds	48
6.1	Evolution of the number of particles used in N -body simulations	59
6.2	Illustration of three mass assignment schemes	64
6.3	Illustration of adaptive mesh refinement	65
6.4	Illustration of an Octree decomposition	66
6.5	Parameter bias induced by the Born approximation	71
7.1	Spatial and redshift setup for the BIGBOX and TILED simulations	75
7.2	Illustration of a 5×5 patch at $z = 1.5$	76
7.3	Normalized lensing efficiency for multiple source redshifts	77
7.4	Convergence maps from the BIGBOX and TILED simulations	78
7.5	Visualization of the Fibonacci grid	78
7.6	Extraction of a patch from the full-sky convergence map using a Fibonacci grid	80
7.7	Gaussian smoothing on a noisy convergence map for multiple smoothing scales	81
7.8	Average standard deviation of noiseless convergence maps	82
7.9	Average $\sigma_1 = \sqrt{\kappa_x^2 + \kappa_y^2}$ of the noiseless convergence maps	84
8.1	Comparison of Mean and Variance for $C_\ell^{\kappa\kappa}$ and Bispectrum	88
8.2	Comparison of Mean and Variance for PDF, Peak/Minima Counts, and Minkowski Functionals	89
8.3	Average BIGBOX/TILED Ratios of Covariance and Correlation Matrices	90
8.4	Covariance Matrices of Summary Statistics in BIGBOX Simulations	91
8.5	Covariance Matrices of Summary Statistics in TILED Simulations	92
8.6	Correlation Matrices of Summary Statistics in BIGBOX and TILED Simulations	93
8.7	Ratios of Covariance Matrices between BIGBOX and TILED Simulations	94
8.8	Ratios of Correlation Matrices between BIGBOX and TILED Simulations	95
8.9	Average BIGBOX/TILED Ratio of Covariance for Multiple Noise Levels	97
8.10	BIGBOX/TILED Ratio of Covariance for Multiple Noise Levels	97
8.11	Average BIGBOX/TILED Ratio of Correlation for Multiple Noise Levels	98
8.12	BIGBOX/TILED Ratio of Correlation for Multiple Noise Levels	98

8.13	Average BIGBOX/TILED Ratio of Covariance for Multiple Smoothing Scales	100
8.14	BIGBOX/TILED Ratio of Covariance for Multiple Smoothing Scales	100
8.15	Average BIGBOX/TILED Ratio of Correlation for Multiple Smoothing Scales	101
8.16	BIGBOX/TILED Ratio of Correlation for Multiple Smoothing Scales	101
8.17	Angular Power Spectrum for Heavily Influenced RIPs	102
8.18	Average BIGBOX / TILED ratios of covariance matrices for the RIPs and the RMPs	103
8.19	Average BIGBOX / TILED ratios of correlation matrices for the RIPs and the RMPs	103
8.20	BIGBOX/TILED Ratios of the mean and variance of summary statistics for the RIPs and the RMPs	105
8.21	Covariance Ratios for RIP	106
8.22	Correlation Ratios for RIP	107

List of Tables

1.1	Comparison of Key Galaxy Surveys for Weak Lensing	4
1.2	Comparison of Large-Volume N-body Simulations	10
6.1	Comparison of N -body simulation methods	61
7.1	Cosmological parameters used in the N-body simulations.	74
7.2	Summary of the summary statistics employed in this study, including their respective value ranges and the computational subroutines used for analysis.	82

Introduction

Contents

1.1	Historical Overview	1
1.2	Astronomical Surveys and Observations	2
1.3	Constraint from Weak Lensing	5
1.4	Higher-Order Statistics in Weak Lensing	6
1.5	Aim of this Thesis	9
1.6	Structure of the Thesis	12

1.1 Historical Overview

Over the past century, cosmological observations have profoundly reshaped our understanding of the universe. From observations of [Hubble \(1925\)](#) measuring the distances to spiral nebulae, including the Andromeda Galaxy, it showed that these ‘nebulae’ were actually galaxies outside the Milky Way. Later in [Hubble \(1929\)](#), it was shown that galaxies are receding from us at velocities proportional to their distances, leading to the formulation of Hubble’s Law, and the revolutionary concept of an expanding universe. Years later, the discovery of the cosmic microwave background (CMB) radiation by [Penzias & Wilson \(1965\)](#) provided compelling evidence for the Big Bang theory, suggesting that the universe originated from a hot and dense state approximately 13.8 billion years ago. Subsequent satellite missions like the Cosmic Background Explorer (COBE; [Smoot et al. 1992](#)) and the Wilkinson Microwave Anisotropy Probe (WMAP; [Bennett et al. 2003](#)) measured the CMB with unprecedented precision finding anisotropies of $\sim 10^{-6}$ K. In the 1970s, Rubin and Ford analyzed galaxy rotation curves and found that galaxies rotate at speeds that cannot be accounted for by the visible matter alone ([Rubin & Ford, 1970](#); [Rubin et al., 1980](#)). This discrepancy provided strong evidence for the existence of dark matter, a mysterious form of matter that does not emit, absorb, or reflect light. Dark matter is now understood to constitute about 26% of the universe’s mass-energy content, while Baryon, making

up our visible world, constitutes only 5 %. The late 1990s witnessed the surprising discovery that the expansion of the universe is accelerating, based on observations of distant Type Ia supernovae by the Supernova Cosmology Project and the High-Z Supernova Search Team (Riess et al., 1998; Perlmutter et al., 1999). This acceleration implies the existence of dark energy, an enigmatic force that permeates all of space and makes up approximately 68% of the universe’s total mass-energy budget. In 2015, the Laser Interferometer Gravitational-Wave Observatory (LIGO) made the first direct detection of gravitational waves providing direct evidence for the existence of gravitational waves, confirming a key prediction of General Relativity, opened a new window for observing cosmic events (Abbott et al., 2016). As observational technologies continue to advance, ongoing and future discoveries promise to further refine our understanding of the universe, addressing the remaining mysteries of dark matter, dark energy, and the fundamental forces shaping the universe.

From a theoretical standpoint, the cornerstone of modern cosmology is Einstein’s General Theory of Relativity, formulated in 1915 (Einstein, 1915). General Relativity provides the fundamental framework for explaining gravitational phenomena on cosmic scales, including the dynamics of the universe’s expansion, black holes, and gravitational lensing effects. Building upon Einstein’s equations, Friedmann (1922) and Lemaître (1931) independently derived solutions that describe a homogeneous and isotropic universe. These solutions led to the concept of an expanding or contracting universe and forms the mathematical foundation for the standard cosmological model. In the late 1940s, Gamow (1948), along with his collaborators Alpher & Herman (1948), proposed the Big Bang nucleosynthesis theory, which explains the formation of light elements like hydrogen and helium in the early universe. Peebles (1968), Zeldovich & Sunyaev (1969) further developed the theory of the recombination era, when the universe cooled enough for electrons and protons to combine into neutral hydrogen, allowing photons to travel freely, creating the CMB radiation. Subsequently, the theory of Baryonic Acoustic Oscillations (BAO) was introduced by Sunyaev & Zeldovich (1970), and independently by Peebles & Yu (1970), which describes the imprint of primordial sound waves in the distribution of galaxies and the CMB. To address the Big-Bang’s challenges, the concept of cosmic inflation was introduced by Guth (1981), Linde (1982), and others. Together, these theories contribute to a comprehensive picture of the universe’s origin, composition, and evolution.

1.2 Astronomical Surveys and Observations

Astronomical surveys are comprehensive observational programs designed to map extensive regions of the sky with high depth and precision, producing critical datasets that address fundamental questions in astrophysics and cosmology. These surveys aim to test the standard cosmological model (Λ CDM) by providing precise measurements that can either confirm or challenge its validity. They tackle key issues such as the Hubble tension, a discrepancy in the measured expansion rate of the




Figure 1.1: Comparison of S_8 values measured from different surveys. The figure highlights the discrepancy between S_8 values inferred from Planck CMB measurements and those derived from late-time cosmic distance ladder observations. Figure adapted from [Madhavacheril et al. \(2024\)](#)

universe derived from early ([Planck Collaboration et al., 2016](#)) and late ([Riess et al., 2019](#)) observations, and inconsistencies in cosmological parameters like S_8 . Figure 1.1 illustrates the S_8 values obtained from various surveys, highlighting the tension between measurements inferred from Planck CMB data and those from late-time cosmic distance ladder observations. Additionally, astronomical surveys investigate the formation and evolution of cosmic structures by mapping millions of galaxies and dark matter distributions using advanced techniques such as cosmic shear and galaxy-galaxy lensing ([Mandelbaum et al., 2013](#); [Abbott et al., 2022](#)).

These surveys employ different methodologies:

Imaging surveys capture wide-field images across multiple wavelengths to map cosmic structures and analyze galaxy populations (e.g., HSC ([Aihara et al., 2018](#)), SDSS ([Kollmeier et al., 2019](#)), DES ([Abbott et al., 2018](#)), LSST ([Ivezić et al., 2019](#))), while spectroscopic surveys collect spectral data revealing redshifts, compositions, and kinematics essential for studying galaxy dynamics and the universe's expansion (e.g., PFS ([Tamura et al., 2016](#)), BOSS ([Dawson et al., 2013](#)), DESI ([DESI Collaboration et al., 2016](#)), KiDS with spectroscopic extensions ([de Jong et al., 2013](#))).

They can be ground-based, utilizing Earth-based telescopes but limited by atmospheric effects (e.g., HSC, DES, KiDS), or space-based, operating above Earth's atmosphere for higher clarity and sensitivity, especially in inaccessible wavelengths (e.g., HST ([Freedman et al., 2001](#)), the upcoming Nancy Grace Roman Space Telescope ([Spergel et al., 2015](#)), and the Euclid mission ([Refregier et al.,](#)

2010)).

Surveys are also classified into Stage-III and Stage-IV based on technological sophistication and scale (Albrecht et al., 2006). Stage-III surveys (e.g., DES, KiDS, HSC) represent the current generation aiming to refine cosmological parameters and deepen understanding of dark energy and dark matter. Stage-IV surveys (e.g., Rubin Observatory (Ivezić et al., 2019), DESI, the upcoming Roman Space Telescope) are the next generation characterized by scale and precision, aiming for high-precision cosmological measurements and deeper exploration of dark energy and dark matter.

Several significant galaxy surveys have been designed to measure weak lensing signals with high precision. Table 1.1 provides a comprehensive overview of four pivotal surveys focusing on their observational capabilities.

Table 1.1: Comparison of Key Galaxy Surveys for Weak Lensing

Survey	Area (deg ²)	Approx. Galaxy Density (arcmin ⁻²)	Median Redshift
DES/KiDS	~5,000	7	0.4
HSC Wide	~1,400	15	0.7
LSST	~18,000	30	1.0
<i>Roman/Euclid</i>	~2,000	50	1.5

The Dark Energy Survey (DES; The Dark Energy Survey Collaboration 2005; Abbott et al. 2018, 2021) utilized the 570-megapixel Dark Energy Camera (DECam; Flaugher et al. 2015) mounted on the 4-m Blanco Telescope at the Cerro Tololo Inter-American Observatory (CTIO) in Chile. Over the course of its operation, DES observed more than 300 million galaxies across approximately 5,000 deg² of the southern sky in five optical bands (g, r, i, z, Y). It achieved an effective galaxy density of about ~ 6 arcmin⁻² and provided photometric redshift estimates up to $z \sim 1.2$. The data collected by DES has made significant contributions to cosmology and astrophysics, including precise measurements of cosmic shear (Amon et al., 2022) and galaxy clustering (Abbott et al., 2022).

The Hyper Suprime-Cam Subaru Strategic Program (HSC-SSP; Aihara et al. 2018) comprises three layers: Wide, Deep, and UltraDeep, conducted with the 8.2-m Subaru Telescope equipped with the 870-megapixel Subaru Hyper Suprime-Cam (HSC; Miyazaki et al. 2018). The Wide layer covers approximately 1,400 deg², yielding galaxy densities of around ~ 15 arcmin⁻². Photometric redshifts extend up to $z \sim 2$. The superior imaging quality of HSC enhances the accuracy of weak lensing measurements and contributes to tighter cosmological constraints (Hikage et al., 2019). Currently, HSC is preparing its final data release (Y6) ending STAGE-III surveys.

The future Legacy Survey of Space and Time (LSST; LSST Science Collaboration et al. 2009; Ivezić et al. 2019) is conducted at the Vera C. Rubin Observatory. Over a 10-year period, LSST will survey approximately 18,000 deg² of the sky. It is expected to detect around 20 billion galaxies, corresponding to galaxy densities exceeding ~ 30 arcmin⁻², with redshift measurements up to $z \sim 3$. LSST's vast dataset will substantially improve the statistical precision of weak lensing analyses and

further refine cosmological models (LSST Dark Energy Science Collaboration, 2012).

Finally, the *Nancy Grace Roman Space Telescope* (Roman; Spergel et al. 2015) will conduct wide-field near-infrared imaging and spectroscopy from space scheduled for launch in the mid-2020s. Covering approximately $2,000 \text{ deg}^2$. The expected galaxy densities exceed $\sim 50 \text{ arcmin}^{-2}$, facilitated by its space-based observations. The mission aims to provide spectroscopic redshifts higher than $z \sim 3$, significantly enhancing the precision of weak lensing measurements.

1.3 Constraint from Weak Lensing

While Λ CDM has been successful in explaining a wide range of cosmological observations, several tensions have emerged between different datasets. statistically significant discrepancy of about 4 to 5σ between the value of the Hubble constant (H_0) inferred from the Planck CMB measurements (Di Valentino et al., 2021) and the late-time measurements of local universe cosmic distance ladder measurements (Riess et al., 2022). In addition to the Hubble tension, discrepancies have been observed in the measurements of the parameter $S_8 \equiv \sigma_8 \sqrt{\Omega_m/0.3}$, where σ_8 represents the root-mean-square amplitude fluctuation of matter density measured in spheres of $8 h^{-1} \text{ Mpc}$, and Ω_m is the present-day matter density parameter. Several large-scale structure (LSS) experiments have reported 2 to 3σ lower values of S_8 compared to those inferred from Planck CMB data (Hikage et al., 2019; Asgari et al., 2021; García-García et al., 2021).

Some of the strongest constraints on S_8 from LSS observations come from the study of cosmic shear, which is the weak gravitational lensing of distant galaxies by the intervening LSS along the line of sight. These small, coherent distortions in the shapes of background galaxies are sensitive to both the amplitude of matter density fluctuations (σ_8) and the growth of these fluctuations over cosmic time (Bartelmann & Schneider, 2001; Bartelmann, 2010; Kilbinger, 2015). While there is a degeneracy between Ω_m and σ_8 in cosmic shear analyses, the product S_8 is tightly constrained (Kilbinger, 2015; Mandelbaum, 2018). Assuming a flat Λ CDM cosmological model and utilizing cosmic shear catalogs, recent analyses from major surveys have measured the cosmological parameter S_8 : DES year 3 analysis reports $S_8 = 0.759^{+0.025}_{-0.023}$ (Amon et al., 2022; Secco et al., 2022) using two-point correlation functions (2PCFs) and $S_8 = 0.793^{+0.038}_{-0.025}$ (Doux et al., 2022) using power spectra (C_ℓ s). The KiDS-1000 analysis reports $S_8 = 0.759^{+0.024}_{-0.021}$ (Asgari et al., 2021) using 2PCFs and $S_8 = 0.754^{+0.027}_{-0.029}$ (Loureiro et al., 2022) using C_ℓ s. The HSC year 3 analysis measure $S_8 = 0.769^{+0.031}_{-0.034}$ (Li et al., 2023) using 2PCFs and $S_8 = 0.776^{+0.032}_{-0.033}$ (Dalal et al., 2023) using C_ℓ s.

As demonstrated in recent analyses, fully and robustly quantifying the cosmological constraints from cosmic shear necessitates accurate estimation and mitigation of various systematic sources and astrophysical effects, which are natural phenomena influencing observations (Dalal et al., 2023). Imperfect modeling of the Point Spread Function (PSF), which characterizes the telescope’s response to

point sources, can introduce systematic errors in galaxy shape measurements (Zhang et al., 2023b). Additionally, uncertainties in the redshift distribution, referring to errors in determining the distances to galaxies, can bias estimates of cosmological parameters (Zhang et al., 2023a). Shear calibration biases, which are systematic errors in measuring the distortion of galaxy shapes, encompass several specific issues: galaxy model bias resulting from incorrect models of galaxy shapes (Bernstein, 2010), noise bias introduced by measurement noise (Refregier et al., 2012), selection bias arising from the preferential selection of certain types of galaxies (Kaiser, 2000), and detection bias caused by errors in the detection process (Sheldon et al., 2020). These biases significantly impact the measured cosmic shear signal and are typically calibrated using image simulations (Mandelbaum et al., 2018b). Furthermore, baryonic feedback processes, which involve interactions from supernovae and active galactic nuclei (Mead et al., 2016; Chisari et al., 2018), and intrinsic alignments, referring to the non-random orientations of galaxies due to local gravitational fields (Joachimi et al., 2015; Kirk et al., 2015; Kiessling et al., 2015), must be carefully controlled to avoid biased cosmological inferences.

1.4 Higher-Order Statistics in Weak Lensing

Traditionally, the power spectrum has been the primary summary statistic used to quantify the distribution of matter density fluctuations in the universe (Hikage et al., 2019; Dalal et al., 2023). However, the weak lensing field is inherently non-Gaussian due to the nonlinear evolution of structures under gravity, leading to features resulting from gravitational collapse, mergers, and other astrophysical processes. The power spectrum, being a two-point statistic, captures only the Gaussian aspects of the field and thus cannot fully characterize these non-Gaussian features.

To fully leverage the information contained in the weak lensing field, it is essential to employ higher-order statistics that are sensitive to the non-Gaussianities. Various higher-order statistics have been studied, such as: higher-order moments (Petri et al., 2015; Gatti et al., 2020), Minkowski functionals (Marques et al., 2019; Grewal et al., 2022; Armijo et al., 2024), probability distribution function (Boyle et al., 2021; Thiele et al., 2023; Uhlemann et al., 2023), Peak and Minima Counts (Martinet et al., 2018; Marques et al., 2024), three-point statistics (Takada & Jain, 2004; Fu et al., 2014), and deep learning (Fluri et al., 2018, 2022). Employing these higher-order statistics enhances the cosmological constraining power of weak lensing analyses, potentially alleviating tensions in parameter estimates and providing deeper insights into the underlying physics.

Figure 1.2 (Euclid Collaboration et al., 2023) shows the forecasted constraints on the cosmological parameters Ω_m and σ_8 from the Euclid mission using ten different higher-order statistics. It is demonstrated that by combining any of these statistics with the standard two-point statistics, The constraints on Ω_m and σ_8 improve by a factor of 2 to 3 for individual parameters and by a factor of 4.5 for their combination. This highlights the potential of higher-order statistics to probe non-Gaussian

features in the weak lensing field and enhance the precision of cosmological parameter estimates.






Figure 1.2: Forecasted constraints on Ω_m and σ_8 from the Euclid mission using ten different higher-order statistics. The figure demonstrates the potential improvements in parameter constraints achievable by incorporating higher-order statistics along with two-point statistics in weak lensing analyses. Figure adapted from [Euclid Collaboration et al. \(2023\)](#)

However, accurately estimating cosmological constraints from these higher-order statistics necessitates precise knowledge of the model uncertainties measured from the covariance matrices. The covariance matrix quantifies the uncertainties and correlations between different summary statistics and is crucial for techniques like Fisher forecasting and likelihood analyses that predict parameter constraints ([Tegmark, 1997](#)). To estimate these covariance matrices, we apply the same statistical measurements to a large ensemble of mock datasets that mimic real observations, accounting for the survey’s specific characteristics, such as cosmic variance ([Seljak, 2009](#)), shot noise ([Wu et al., 2019](#)) and super-sample covariance (SSC; [Takada & Hu 2013](#)). SSC arises from the correlation between observed modes and modes whose wavelengths are larger than the survey size. For instance, if the observed region is embedded in a large-scale super-survey overdensity, the structures within the survey have evolved faster compared to a region embedded in the cosmic mean density. For weak lensing surveys, these mock datasets are generated through ray-tracing simulations of light propagation through the universe, using light cones constructed from cosmological N-body simulations ([Shirasaki et al., 2019](#); [Euclid Collaboration et al., 2024](#)).

One common approach to generating non-repeating lightcones that span a wide range of redshifts involves stacking multiple multi-resolution simulation boxes (Sato et al., 2009; Fosalba et al., 2015b; Crocce et al., 2015; Takahashi et al., 2017; DeRose et al., 2019), as illustrated in the left panel of Figure 1.3. This method effectively captures the evolution of cosmic structures over time; however, it often faces challenges in achieving high redshift resolution and demands substantial computational resources. Alternatively, constructing lightcones by repetitively tiling a single simulation box along the line of sight preserves high redshift resolution and is computationally efficient (Sehgal et al., 2010; Liu et al., 2018; Stein et al., 2020; Omori, 2024). Nonetheless, this repetitive box method introduces artefacts such as box replication effects (Chen & Yu, 2024) (see Figure 1.4). These artefacts can result in the underestimation of variance on large scales and introduce biases in the mean values of summary statistics, potentially impacting the accuracy of cosmological parameter estimations (Zürcher et al., 2021). To mitigate these issues, various simulation strategies have been developed. For instance, Omori (2024) performed random rotations of the projected HEALPix shells every 1 Gpc/ h to avoid the repeating structures along the line of sight (shown in the right panel of Figure 1.3). However, this strategy is found to introduce decorrelation on large scales, potentially affecting the summary statistics and their covariance estimation (Upadhye et al., 2024).



Left: Stacking multiple multi-resolution simulation boxes to construct non-repeating lightcones.



Right: Replicating a single simulation box along the line of sight with random rotations to avoid repeating structures.

Figure 1.3: Examples of simulation strategies for generating lightcones. Figures adapted from Sato et al. (2009) and Omori (2024).




Figure 1.4: Visualization of a partial sky convergence map covering $5 \times 5 \text{ deg}^2$. Different box repetition times, denoted by N , indicate the number of times the simulation box is replicated along the line of sight. The upper panels correspond to the most affected direction $(1, 0, 0)$, while the middle and lower panels represent the directions $(1, 1, 0)$ and $(1, 1, 1)$, respectively. The replication effects become more pronounced with increasing N . Figure adapted from [Chen & Yu \(2024\)](#).

1.5 Aim of this Thesis

The primary goal of this thesis is to improve the precision and robustness of cosmological parameter constraints derived from higher-order weak lensing statistics. In pursuing this objective, the thesis addresses two interrelated challenges:

1. quantifying the influence of super-sample covariance (SSC) on higher-order observables
2. evaluating how replicated box structures bias summary statistics and their covariance.

Accurate covariance estimation is critical for reliable parameter inference; hence, mitigating these effects is essential for maximizing the scientific yield of current and future weak lensing surveys. The findings presented here provide practical guidelines on incorporating large-scale modes and managing periodic tiling, thereby enhancing the reliability and interpretability of weak lensing data.

1.5.1 Motivation: Why Not a Single Huge Simulation?

A simple way to avoid both the super-sample effect and the errors caused by repeating smaller simulation boxes is to run one extremely large simulation. Although this approach seems ideal, current computer power is not yet sufficient for such massive simulations. Table 1.2 shows several large-volume N-body simulations (Angulo et al., 2012; Skillman et al., 2014; Fosalba et al., 2015a; Heitmann et al., 2019; Cheng et al., 2020; Maksimova et al., 2021; Euclid Collaboration et al., 2024; Bayer et al., 2024), each of which has helped us understand the universe’s large-scale structure, cosmic evolution, and galaxy formation. However, these simulations require huge computing resources, so researchers can run only a limited number of them. This small number is not enough for studies such as covariance estimation or other methods that rely on many simulation runs. Therefore, despite ongoing improvements in computing technology, scientists often use smaller simulations and “tile” or repeat them to reach the necessary volume and statistical accuracy.

Table 1.2: Comparison of Large-Volume N-body Simulations. The table lists the box size, particle count, and mass resolution of several prominent N-body simulations. The simulations are ordered by box size, with the largest simulations at the top. The last two rows correspond to the simulations used in this work.

Simulation	Box Size [h^{-1} Gpc]	Particle Count	Mass Resolution [$h^{-1} M_{\odot}$]
Euclid Flagship	3.6	16000^3	1×10^9
OuterRim	4.225	10240^3	1.85×10^9
Millennium XXL	3	6720^3	6.17×10^9
DarkSky	8	10240^3	3.9×10^{10}
Uchuu	2	12800^3	3.27×10^8
AbacusSummit	2	6912^3	2×10^9
Cosmo-π	3.2	16384^3	$\sim 10^8$
MICE-GC	3.072	4096^3	2.93×10^{10}
HalfDome (BigBox)	3.75	6144^3	1.95×10^{10}
Tiled (This Work)	0.625	1024^3	1.95×10^{10}

1.5.2 Super-Sample Effect

The super-sample covariance (SSC) arises from correlations with modes whose wavelengths exceed the size of a given survey. In a single, modestly sized simulation box, these large-scale modes are not fully captured, leading to an underestimation of super-survey fluctuations and, consequently, biases in cosmological parameter constraints. Although SSC has been investigated thoroughly for the power spectrum (Takada & Hu, 2013; Barreira & Schmidt, 2017; Barreira et al., 2018), its role in higher-order weak lensing statistics remains comparatively underexplored.

Figure 1.5 illustrates the contributions of the Gaussian, SSC, and connected non-Gaussian (cNG) terms to the diagonal elements of the total covariance for the lensing convergence power spectrum. As

shown, SSC becomes increasingly significant at high multipole moments (ℓ), corresponding to small angular scales, and can account for up to 50% of the total covariance. Although theoretical predictions incorporating SSC (Linke et al., 2023; Uhlemann et al., 2023) have been proposed, they have yet to be validated comprehensively for higher-order weak lensing observables (Euclid Collaboration et al., 2023). Furthermore, while the effects of SSC have been explored in three-dimensional (3D) box simulations (Bayer et al., 2023), their impact on two-dimensional (2D) weak lensing simulations remains insufficiently characterized.




Figure 1.5: Contributions of Gaussian, super-sample covariance (SSC), and connected non-Gaussian (cNG) terms to the diagonal elements of the total covariance matrix for the lensing convergence power spectrum. The figure highlights the increasing significance of SSC at high multipole moments (ℓ), corresponding to small scales. Figure adapted from Barreira et al. (2018).

1.5.3 Box Replication Effects

A commonly used strategy to simulate large cosmological volumes involves periodically replicating smaller simulation boxes along different axes. Although computationally efficient and effective at providing high redshift resolution, this method introduces artificial periodicity, which can lead to the underestimation of variance on large scales and to biases in the mean of key summary statistics (Zürcher et al., 2021). Past studies have mostly focused on the power spectrum and lower-order moments (Chen & Yu, 2024), leaving higher-order statistics relatively unexplored. Although Fluri et al. (2019) investigated the replication scheme’s influence on Convolutional Neural Network (CNN) predictions and the power spectrum by increasing box size and particle count, the corresponding effects on the covariance of higher-order weak lensing observables remain unclear.

1.5.4 Simulation Strategies

To address these gaps, we employ two simulation approaches:

BigBox Simulations Large-volume simulations that include super-survey modes, naturally capturing the SSC effects inherent in the Universe’s large-scale structure.

Tiled Simulations Simulations that replicate smaller boxes to cover the desired light cone; by construction, these do not contain super-survey modes and thereby underestimate SSC.

By comparing these two approaches, we quantify the influence of the super-sample effect on higher-order statistics and their associated covariance matrices, thus determining its impact on cosmological parameter estimates. We also investigate the role of varying smoothing scales and shape noise levels—both key factors in weak lensing analyses—to identify the conditions under which SSC substantially alters the covariance of higher-order statistics.

Subsequently, we examine how replication artifacts bias higher-order statistics and their covariance by focusing on heavily tiled regions. As higher-order observables play an increasingly important role in weak lensing studies, understanding and mitigating these systematic effects is critical for deriving robust cosmological inferences. Our findings yield practical guidelines for reducing replication-induced biases and enhance the reliability of future weak lensing surveys.

1.6 Structure of the Thesis

This thesis is organized into nine chapters that systematically develop the theoretical framework, methodologies, and empirical analyses relevant to the research objectives outlined in the previous sections.

Chapter 2 introduces the foundations of modern cosmology, tracing the field’s historical development and emphasizing the key theoretical advances that inform our current understanding of the Universe. In Chapter 3, the fundamental concepts and observational aspects of weak gravitational lensing are examined. Chapter 4 then focuses on the summary statistics most commonly used to extract cosmological information from weak lensing data, with particular attention to the power spectrum and higher-order statistics.

Chapter 5 presents a detailed discussion of covariance matrices, including super-sample covariance (SSC), and explores their role in shaping cosmological parameter constraints. Chapter 6 provides an overview of numerical methods in astrophysics and cosmology, with a focus on N-body and weak lensing simulations. The methodological framework employed in this thesis—encompassing simulation strategies, data generation, statistical measurements, and covariance matrix estimation—is described in Chapter 7.

In Chapter 8, the outcomes of our analyses are reported, comparing mean values and covariance matrices of higher-order statistics obtained from the BIGBOX and TILED simulations. The influence of SSC on variances and correlations is also assessed. Finally, Chapter 9 concludes the thesis by summarizing the primary findings and contributions, emphasizing how they extend our understanding of super-sample covariance and box replication effects in higher-order weak lensing statistics.

Chapter 2

Foundational Concepts in Modern Cosmology

Contents

2.1	Einstein Field Equations	14
2.2	FLRW Metric and the Friedmann Equations	17
2.3	Redshift and Cosmological Distances	19
2.4	Initial Conditions and the Primordial Universe	21
2.5	Growth of Matter Density Fluctuations	24
2.6	The Spherical Collapse Model	26
2.7	Dark Matter Halos	28

In this chapter, we provide an overview of the fundamental concepts and equations that underpin the field of cosmology following [Dodelson \(2003\)](#) and [Weinberg \(2008\)](#) textbooks.

2.1 From Einstein to Cosmology: The Einstein Field Equations

The Einstein Field Equations (EFE) are the fundamental equations of General Relativity, describing how matter and energy influence the curvature of spacetime. Introduced by [Einstein \(1915\)](#), these equations extend Newton’s law of universal gravitation to a relativistic context, generalizing the definition of inertial observer in a spacetime which can be curved under the presence of a strong gravitational potential.

The EFE establish a relationship between the geometry of spacetime and the distribution of matter within it. They are expressed with the Minkowski Metric $g_{\mu\nu}$, which describes the curvature of spacetime, and the stress-energy tensor $T_{\mu\nu}$, which characterizes the distribution of matter and energy. The EFE are given by:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}, \quad (2.1)$$

where Λ is the cosmological constant, G is the gravitational constant, and c is the speed of light. The

Einstein tensor $G_{\mu\nu}$ is defined by:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}, \quad (2.2)$$

the $R_{\mu\nu}$ is the Ricci tensor, which can be written using the Christoffel symbols $\Gamma_{\mu\nu}^\lambda$ of metric connection as:

$$R_{\mu\nu} = \partial_\lambda \Gamma_{\mu\nu}^\lambda - \partial_\nu \Gamma_{\mu\lambda}^\lambda + \Gamma_{\lambda\sigma}^\lambda \Gamma_{\mu\nu}^\sigma - \Gamma_{\mu\sigma}^\lambda \Gamma_{\lambda\nu}^\sigma, \quad (2.3)$$

the Ricci scalar R is given by:

$$R = g^{\mu\nu} R_{\mu\nu}, \quad (2.4)$$

2.1.1 The Stress-Energy Tensor for a Perfect Fluid

Assuming a perfect fluid as the source of the gravitational field, the stress-energy tensor is given by

$$T_{\mu\nu} = (\rho + p) u_\mu u_\nu + pg_{\mu\nu}, \quad (2.5)$$

where ρ is the energy density, p is the pressure, and u_μ is the four-velocity of the fluid. In a homogeneous and isotropic universe, u_μ is given by

$$u_\mu = (-c, 0, 0, 0), \quad (2.6)$$

Therefore, each component of the stress-energy tensor can be expressed as

$$T_{00} = \rho c^2, \quad T_{ij} = pg_{ij}, \quad (2.7)$$

2.1.2 Equations of State for Different Cosmic Components

Different speices of matter and energy contribute to the energy density ρ and pressure P in the universe. The equation of state parameter w is defined as the ratio of pressure P to energy density ρ :

$$w = \frac{p}{\rho}. \quad (2.8)$$

For radiation, the equation of state parameter can be derived by considering the trace of the stress-energy tensor:

$$0 = T = g^{\mu\nu} T_{\mu\nu} = (\rho + p)(-1) + 4p = -\rho + 3p, \quad (2.9)$$

For non-relativistic matter where $p = 0$, the equation of state parameter is $w = 0$. For cosmological constant, the equation of state parameter can be derived by comparing the effective stress-energy of

the cosmological constant $T_{\mu\nu}^{(\Lambda)} = \rho_\Lambda g_{\mu\nu}$ to the stress-energy tensor of a perfect fluid:

$$\rho_\Lambda = \frac{\Lambda c^4}{8\pi G} \quad (2.10)$$

$$(\rho_\Lambda + p_\Lambda) u_\mu u_\nu + p_\Lambda g_{\mu\nu} = T_{\mu\nu}^{(\Lambda)} = \rho_\Lambda g_{\mu\nu} \quad (2.11)$$

$$\rho_\Lambda + p_\Lambda = 0 \quad (\text{valid for all } \mu, \nu)$$

$$p_\Lambda = -\rho_\Lambda \quad (2.12)$$

Therefore, the equation of state parameter for cosmological constant is $w = -1$.

To summarize, the Equation of States for each component of the universe are:

$$w = \begin{cases} 0 & \text{matter,} \\ \frac{1}{3} & \text{Radiation,} \\ -1 & \text{Cosmological Constant.} \end{cases} \quad (2.13)$$

In realistic universe, the energy density and pressure are contributed by multiple components. Assuming the interaction between different components is negligible, the total stress-energy tensor is the sum of the individual stress-energy tensors:

$$T_{\mu\nu} = \sum_i (T_i)_{\mu\nu}, \quad (2.14)$$

where i denotes the different components of the universe. Therefore, the total energy density and pressure are given by:

$$\rho = \sum_i \rho_i, \quad p = \sum_i p_i. \quad (2.15)$$

Unlike the conditions in the early Universe, when photons and baryons were tightly coupled, we now consider an epoch in which each component evolves independently, without any exchange of heat or other forms of energy. Under these assumptions, the conservation of energy-momentum for each component may be derived from the first law of thermodynamics. Specifically:

$$\begin{aligned} d(\rho_i a^3) &= -p_i d(a^3) \\ a^3 d\rho_i &= -\rho_i(1 + w_i) d(a^3) \quad (p_i = w_i \rho_i) \\ \ln \rho_i &= -3(1 + w_i) \ln a + \text{const.} \\ \rho_i &\propto a^{-3(1+w_i)}. \end{aligned} \quad (2.16)$$

where w_i is the equation of state parameter for the i -th component.

2.2 FLRW Metric and the Friedmann Equations

2.2.1 The FLRW Metric

The Friedmann-Lemaître-Robertson-Walker (FLRW) metric describes a homogeneous and isotropic universe. As obtained by [Weinberg \(1972\)](#):

$$ds^2 = -c^2 dt^2 + a^2(t) [d\chi^2 + f_K^2(\chi) (d\theta^2 + \sin^2 \theta d\phi^2)], \quad (2.17)$$

where $a(t)$ is the time evolving scale factor, χ is the comoving radial coordinate, and $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$ represents the metric on the unit two-sphere. The function $f_K(\chi)$ encodes the spatial curvature of the universe and is defined as:

$$f_K(\chi) = \begin{cases} \frac{1}{\sqrt{-K}} \sinh(\sqrt{-K}\chi) & \text{for } K < 0, \\ \chi & \text{for } K = 0, \\ \frac{1}{\sqrt{K}} \sin(\sqrt{K}\chi) & \text{for } K > 0, \end{cases} \quad (2.18)$$

where K is the spatial curvature constant, with $K < 0$ corresponding to an open universe, $K = 0$ to a flat universe, and $K > 0$ to a closed universe.

2.2.2 Derivation of the Friedmann Equations

For the FLRW metric, the non-zero components of the Einstein tensor are:

$$G_{00} = 3 \left(\frac{\dot{a}}{a} \right)^2 + 3 \frac{Kc^2}{a^2}, \quad (2.19)$$

$$G_{ij} = - \left(2 \frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + \frac{Kc^2}{a^2} \right) a^2 g_{ij}, \quad (2.20)$$

where the dot denotes differentiation with respect to cosmic time t .

Substituting the components of $G_{\mu\nu}$ and $T_{\mu\nu}$ into the Einstein field equations (2.1), we obtain the Friedmann equations ([Friedmann, 1922](#)):

- First Friedmann equation (00 component):

$$3 \left(\frac{\dot{a}}{a} \right)^2 + 3 \frac{Kc^2}{a^2} + \Lambda c^2 = \frac{8\pi G}{c^4} \rho c^2 \quad (2.21)$$

$$\left(\frac{\dot{a}}{a} \right)^2 = \frac{8\pi G}{3} \rho - \frac{Kc^2}{a^2} + \frac{\Lambda c^2}{3} \quad (2.22)$$

- Second Friedmann equation (ii component):

$$-\left(2\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{Kc^2}{a^2}\right)g_{ii} + \Lambda c^2 g_{ii} = \frac{8\pi G}{c^4} P g_{ii} \quad (2.23)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{c^4} P - \frac{1}{2} \left(\left(\frac{\dot{a}}{a}\right)^2 + \frac{Kc^2}{a^2} - \Lambda c^2 \right) \quad (2.24)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3P}{c^2} \right) + \frac{\Lambda c^2}{3} \quad (2.25)$$

2.2.3 Critical Density, Density Parameters, and H_0

Introducing the Hubble parameter H and the critical density ρ_c , we can simplify the Friedmann equations. The Hubble parameter is defined as:

$$H = \frac{\dot{a}}{a}, \quad (2.26)$$

and the critical density is defined as:

$$\rho_c = \frac{3H^2}{8\pi G}. \quad (2.27)$$

Substituting H and ρ_c into the first Friedmann equation (2.22), we obtain:

$$H^2 = H^2 \frac{\rho}{\rho_c} - \frac{Kc^2}{a^2} + \frac{\Lambda c^2}{3}. \quad (2.28)$$

Rearranging terms, we get:

$$1 = \frac{\rho}{\rho_c} - \frac{Kc^2}{a^2 H^2} + \frac{\Lambda c^2}{3H^2}. \quad (2.29)$$

Defining the density parameters:

$$\Omega_m = \frac{\rho_m}{\rho_c}, \quad \Omega_r = \frac{\rho_r}{\rho_c}, \quad \Omega_\Lambda = \frac{\rho_\Lambda}{\rho_c} = \frac{\Lambda c^2}{3H^2}, \quad \Omega_K = -\frac{Kc^2}{a^2 H^2}, \quad (2.30)$$

where ρ_m and ρ_r are the energy densities of matter and radiation, respectively, and ρ_Λ is the effective energy density associated with the cosmological constant, we can write the first Friedmann equation as:

$$1 = \Omega_r + \Omega_m + \Omega_K + \Omega_\Lambda. \quad (2.31)$$

2.2.4 Evolution of Density Parameters with Scale Factor

The evolution of the density parameters with the scale factor a can be derived from the conservation of energy-momentum and the equations of state. For matter-dominated and radiation-dominated

universes, the energy densities scale as:

$$\rho_m \propto a^{-3}, \quad \rho_r \propto a^{-4}. \quad (2.32)$$

Therefore, the corresponding density parameters vary with a as:

$$\Omega_m(a) = \Omega_{m0} a^{-3} \left(\frac{H_0}{H(a)} \right)^2, \quad \Omega_r(a) = \Omega_{r0} a^{-4} \left(\frac{H_0}{H(a)} \right)^2, \quad (2.33)$$

where the subscript 0 denotes present-day values, and H_0 is the current Hubble parameter. Conventionally, the Hubble parameter is parametrized as:

$$H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (2.34)$$

where h is a dimensionless parameter that accounts for the uncertainty in the exact value of H_0 and the cosmology assumed. It allows cosmological quantities to be expressed in a way that separates the dependence on the H_0 . Combining these expressions, the Friedmann equation (2.31) can be rewritten in terms of the present-day density parameters:

$$\left(\frac{H(a)}{H_0} \right)^2 = \Omega_{r0} a^{-4} + \Omega_{m0} a^{-3} + \Omega_{K0} a^{-2} + \Omega_{\Lambda0}, \quad (2.35)$$

which describes the evolution of the Hubble parameter with scale factor a in terms of the contributions from radiation, matter, curvature, and the cosmological constant.

2.3 Redshift and Cosmological Distances

The expansion of the universe causes the wavelength of light to stretch as it propagates through space. This effect, known as cosmological redshift, is a direct consequence of the expansion of the universe and is described by the scale factor $a(t)$.

Before going into the details of the redshift, we first introduce the concept of the null geodesic $ds^2 = 0$, which describes the path of light in a curved spacetime. Then, the radial coordinate distance for a photon traveling from a source to the observer is obtained from the null condition:

$$ds^2 = 0 \quad \Rightarrow \quad d\chi = \frac{c dt}{a(t)}. \quad (2.36)$$

Integrating this expression, we obtain the comoving radial distance χ as:

$$\chi = \int_{t_e}^{t_0} \frac{c dt'}{a(t')} \quad (2.37)$$

where t_0 is the present time, and t_e is the time at which the light is emitted. Suppose a light is emitted at time $t = t_e$ with wavelength λ_e and received at time $t = t_0$ with wavelength λ_0 . Then, the same phase emitted at time $t = t_e + \Delta t_e$ is received at time $t = t_0 + \Delta t_0$. The comoving distance between the source and the observer is given by:

$$\chi = \int_{t_e}^{t_0} \frac{c dt'}{a(t')} = \int_{t_e + \Delta t_e}^{t_0 + \Delta t_0} \frac{c dt'}{a(t')}. \quad (2.38)$$

Assuming the scale factor $a(t)$ change is negligible over the time interval Δt , we can expand the integrand in Eq. (2.38) to first order:

$$\int_{t_e + \Delta t_e}^{t_0 + \Delta t_0} \frac{c dt'}{a(t')} - \int_{t_e}^{t_0} \frac{c dt'}{a(t')} = \frac{c \Delta t_e}{a(t_e)} - \frac{c \Delta t_0}{a(t_0)} = 0. \quad (2.39)$$

Therefore, the fractional change in the wavelength of light is given by:

$$\frac{\lambda_0 - \lambda_e}{\lambda_e} = \frac{a(t_0)}{a(t_e)} - 1, \quad (2.40)$$

This shift in the wavelength of light is known as the cosmological redshift z , defined as:

$$z := \frac{\lambda_0 - \lambda_e}{\lambda_e} = \frac{a(t_0)}{a(t_e)} - 1, \quad (2.41)$$

Therefore, the redshift z serves us a measure of position in the expanding universe, with $z = 0$ corresponding to the present time and $z \rightarrow \infty$ corresponding to the time of the Big Bang. However, the space-time is not static nor flat, there are several distance measures that are used to describe the separation between two objects in the universe. A characteristic length scale is the Hubble distance, defined as the inverse of the Hubble parameter:

$$d_H = \frac{c}{H_0} = 9.26 \times 10^{25} h^{-1} \text{m} \quad (2.42)$$

The most common distance measures are the comoving distance, the luminosity distance, and the angular diameter distance.

2.3.1 Luminosity Distance

The luminosity distance $d_L(z)$ is a measure of observed distance broadly used in Astronomy and Cosmology, relating the intrinsic luminosity L of a source to the observed flux F via the inverse-square law (Carroll et al., 1992):

$$F = \frac{L}{4\pi d_L^2}. \quad (2.43)$$

In an expanding universe, the luminosity distance accounts for the effects of redshift on both the energy of photons and the rate at which they are received. It is defined as (Hogg, 1999):

$$d_L(z) = (1 + z) f_K(\chi(z)). \quad (2.44)$$

The luminosity distance is crucial for determining cosmological parameters using standard candles, such as Type Ia supernovae, whose intrinsic luminosities are known (Riess et al., 1998). By measuring the observed flux F and applying Eq. (2.43), we can infer $d_L(z)$ and constrain cosmological models.

2.3.2 Angular Diameter Distance

The angular diameter distance $d_A(z)$ relates the physical size D of an object to its observed angular size:

$$\theta = \frac{D}{d_A}. \quad (2.45)$$

In an expanding universe, the angular diameter distance is given by (Hogg, 1999):

$$d_A(z) = \frac{f_K(\chi(z))}{1 + z}. \quad (2.46)$$

The angular diameter distance is essential for studying standard rulers, such as the scale of baryon acoustic oscillations (BAO) in the cosmic microwave background and large-scale structure (Eisenstein et al., 2005). By measuring the angular size θ of these features and knowing their physical size D , we can determine $d_A(z)$ and thus constrain cosmological parameters.

2.4 Initial Conditions and the Primordial Universe

The initial conditions of the universe are believed to originate from quantum fluctuations during the inflationary epoch. These fluctuations are imprinted on the cosmic microwave background (CMB) radiation, which provides a snapshot of the universe at the time of recombination ($z \sim 1100$), and these primordial perturbations served as the seeds for the formation of the large-scale structures (LSS) observed today

Figure 2.1 illustrates the CMB temperature power spectrum (TT) measured by the Planck satellite (Planck Collaboration et al., 2014), showing the primordial fluctuations imprinted at large angular scales ($\ell \lesssim 30$).

2.4.1 Primordial Fluctuations and the Power Spectrum

The standard single-field slow-roll inflation model predicts that the primordial fluctuations are nearly scale-invariant and Gaussian (Dodelson, 2003). The matter power spectrum of these primordial cur-




Figure 2.1: Angular Power Spectrum of Cosmic Microwave Background Temperature Fluctuations as a function of the multipole moment (ℓ) and the corresponding angular scale. The observational data are depicted by red markers with associated error bars, whereas the theoretical prediction derived from the Λ CDM cosmological model is represented by the green curve (Planck Collaboration et al., 2014). The primordial fluctuations imprinted at large angular scales ($\ell \lesssim 30$).

vature perturbations is described by a nearly scale-invariant power-law form:

$$P_p(k) = A_s \left(\frac{k}{k_*} \right)^{n_s - 1}, \quad (2.47)$$

where A_s is the amplitude of the scalar fluctuations, k_* is the pivot scale, and n_s is the spectral index. The observational constraints on these parameters are provided by the Planck satellite (Planck Collaboration et al., 2020):

$$A_s = (2.101^{+0.031}_{-0.034}) \times 10^{-9}, \quad n_s = 0.965 \pm 0.004. \quad (2.48)$$

for the pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$.

As the universe evolves, various physical processes, such as radiation pressure, baryon-photon interactions, and dark matter dynamics, influence the growth of these initial perturbations. These effects are encapsulated in the transfer function $T(k)$, which modifies the primordial power spectrum to give the linear matter power spectrum at redshift z (Dodelson, 2003):

$$P(k, z) = P_p(k) T^2(k) D^2(z), \quad (2.49)$$

where $D(z)$ is the linear growth factor that describes the growth of perturbations in the linear regime, where each k -mode evolves independently of the others. The growth factor is given by:

$$D(a) = \frac{5\Omega_m a}{2} \int_0^1 \frac{dx}{(\Omega_m/x + \Omega_\Lambda x^2 + 1 - \Omega_m - \Omega_\Lambda)^{3/2}}, \quad (2.50)$$

where at the limit $a \rightarrow 0$, $D(a) \rightarrow a$.

The shape of $T(k)$ is determined by the Einstein-Boltzmann equations of a mixture of various energy components. Thus, there is no exact analytical form for $T(k)$; instead, it is typically computed using numerical codes such as CAMB (Lewis et al., 2000) and CLASS (Blas et al., 2011).

Qualitatively, the transfer function behaves differently on scales relative to the equality scale k_{eq} . Since in the radiation-dominated era, the growth of perturbations on super-horizon scales is suppressed compared to those on sub-horizon scales. In the matter-dominated era, the growth of perturbations is the same on super-horizon and sub-horizon scales. Due to this, the transfer function behaves as:

$$T(k) \propto \begin{cases} 1 & \text{for } k \ll k_{\text{eq}} \text{ Top-hat,} \\ k^{-2} & \text{for } k \gg k_{\text{eq}} \text{ Power-law.} \end{cases} \quad (2.51)$$

Figure 2.2 shows the linear matter power spectrum computed using the CLASS code. It is clear that the scaling of the linear power spectrum changes around k_{eq} . The wiggles around $k \sim 0.1 h/\text{Mpc}$ are due to Baryon Acoustic Oscillations (BAOs) imprinted.




Figure 2.2: The linear matter power spectrum as a function of the wavenumber k computed using the CLASS code. The transition from the radiation-dominated to the matter-dominated era is evident around $k_{\text{eq}} = 0.0104 h/\text{Mpc}$ (Planck Collaboration et al., 2020).

2.4.2 Baryon Acoustic Oscillations (BAOs)

BAOs serve as a standard ruler for cosmological distance measurements and are crucial for constraining cosmological parameters. These features originate from the oscillatory behavior of the photon-baryon plasma in the primordial universe prior to the epoch of recombination. During this period, photons and baryons are tightly coupled, effectively forming a coherent photon-baryon fluid. The coupling between photons and baryons is quantitatively characterized by the baryon-to-photon momentum density ratio, R , defined as:

$$R = \frac{\Pi_b}{\Pi_\gamma} = \frac{\rho_b \mathbf{v}_b}{(1 + \frac{1}{3}) \rho_\gamma \mathbf{v}_\gamma} = \frac{3\rho_b}{4\rho_\gamma}, \quad (2.52)$$

where ρ_b and ρ_γ denote the baryon and photon energy densities, respectively, while $\mathbf{v}_b$ and $\mathbf{v}_\gamma$ represent their respective velocities.

The propagation of acoustic waves within the photon-baryon plasma is governed by the effective sound speed, c_s , which arises from the interplay between radiation pressure and gravitational infall. The effective sound speed is derived from the effective pressure and energy density of the photon-baryon fluid:

$$\begin{aligned} c_s^2 &= \frac{\partial p_{\text{eff}}}{\partial \rho_{\text{eff}}} = \frac{\partial p_\gamma}{\partial (\rho_b + \rho_\gamma)} \\ &= \frac{1}{1 + R} \frac{\partial p_\gamma}{\partial \rho_\gamma} = \frac{1}{1 + R} \cdot \frac{1}{3} \\ c_s &= \frac{1}{\sqrt{3(1 + R)}}, \end{aligned} \quad (2.53)$$

where p_γ and ρ_γ are the photon pressure and energy density, respectively.

The acoustic oscillations in the photon-baryon plasma can be described by solutions to the linearized perturbation equations (Eq (2.66); we will discuss in the next section). Therefore, the evolution of the photon-baryon plasma scales as:

$$\delta_\gamma(k, t) \propto \cos(kc_s t + \phi), \quad (2.54)$$

where k is the wavenumber and ϕ is the phase constant.

2.5 Growth of Matter Density Fluctuations

Density fluctuations arise from quantum fluctuations during inflation and grow under the influence of gravity. Starting from the continuity and Euler equations, which govern the conservation of mass

and momentum, respectively:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (2.55)$$

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla P}{\rho} - \nabla \Phi, \quad (2.56)$$

where ρ is the density, $\mathbf{v}$ is the peculiar velocity field, P is the pressure, and Φ is the gravitational potential.

To analyze perturbations in an expanding universe, we move to comoving coordinates and express the density as a perturbation around the mean density, $\rho = \bar{\rho}(1 + \delta)$, where δ is the density contrast. The continuity and Euler equations then become:

$$\frac{\partial \delta}{\partial t} + \frac{1}{a} \nabla \cdot ((1 + \delta) \mathbf{v}) = 0, \quad (2.57)$$

$$\frac{\partial \mathbf{v}}{\partial t} + H \mathbf{v} + \frac{1}{a} (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla \delta P}{a \bar{\rho} (1 + \delta)} - \frac{1}{a} \nabla \Phi, \quad (2.58)$$

where $a(t)$ is the scale factor, and $H = \dot{a}/a$ is the Hubble parameter.

To derive the equation of motion for the density contrast, we linearize the above equations under the assumption that $\delta \ll 1$ and $\mathbf{v}$ is small. Neglecting higher-order terms in δ and $\mathbf{v}$, we obtain the linearized equations:

$$\frac{\partial \delta}{\partial t} + \frac{1}{a} \nabla \cdot \mathbf{v} = 0, \quad (2.59)$$

$$\frac{\partial \mathbf{v}}{\partial t} + H \mathbf{v} = -\frac{\nabla \delta P}{a \bar{\rho}} - \frac{1}{a} \nabla \Phi. \quad (2.60)$$

The gravitational potential Φ is related to the density contrast via Poisson's equation in comoving coordinates:

$$\nabla^2 \Phi = 4\pi G \bar{\rho} a^2 \delta, \quad (2.61)$$

where G is the gravitational constant. Assuming adiabatic perturbations, the pressure perturbation is related to the density perturbation by $\delta P = c_s^2 \delta \rho = c_s^2 \bar{\rho} \delta$, where c_s is the sound speed.

Taking the time derivative of the linearized continuity equation (2.59) and substituting the divergence of $\mathbf{v}$ from the linearized Euler equation (2.60), we obtain:

$$\frac{\partial^2 \delta}{\partial t^2} + 2H \frac{\partial \delta}{\partial t} - \frac{c_s^2}{a^2} \nabla^2 \delta = 4\pi G \bar{\rho} \delta. \quad (2.62)$$

Transforming to Fourier space, where $\nabla^2 \delta \rightarrow -k^2 \tilde{\delta}(k, t)$, the equation becomes:

$$\ddot{\tilde{\delta}}(k, t) + 2H\dot{\tilde{\delta}}(k, t) + \left(\frac{c_s^2 k^2}{a^2} - 4\pi G \bar{\rho} \right) \tilde{\delta}(k, t) = 0, \quad (2.63)$$

where $\tilde{\delta}(k, t)$ is the Fourier transform of the density contrast.

Defining the effective frequency squared $\omega^2(k, t) = 4\pi G \bar{\rho} - \frac{c_s^2 k^2}{a^2}$, the equation simplifies to:

$$\ddot{\tilde{\delta}}(k, t) + 2H\dot{\tilde{\delta}}(k, t) - \omega^2(k, t)\tilde{\delta}(k, t) = 0. \quad (2.64)$$

The solutions to this differential equation depend on the sign of $\omega^2(k, t)$:

- **Gravity-Dominated Regime** ($\omega^2(k, t) > 0$): For large-scale perturbations where gravity overcomes pressure forces (i.e., small k), the solutions are exponential:

$$\tilde{\delta}(k, t) = C_1 e^{\lambda t} + C_2 e^{-\lambda t}, \quad (2.65)$$

where $\lambda = \sqrt{\omega^2(k, t)}$. The growing mode ($e^{\lambda t}$) leads to the amplification of perturbations and structure formation, while the decaying mode ($e^{-\lambda t}$) becomes negligible over time.

- **Pressure-Dominated Regime** ($\omega^2(k, t) < 0$): For small-scale perturbations where pressure resists gravitational collapse (i.e., large k), the solutions are oscillatory:

$$\tilde{\delta}(k, t) = e^{-Ht} (C_1 \cos(|\omega(k, t)|t) + C_2 \sin(|\omega(k, t)|t)). \quad (2.66)$$

The perturbations oscillate with frequency $|\omega(k, t)|$ and are damped by the cosmic expansion, preventing collapse on small scales.

These results illustrate the Jeans instability criterion, which states that perturbations grow only if their wavelength exceeds the Jeans length $\lambda_J = c_s \sqrt{\frac{\pi}{G \bar{\rho}}}$ (Jeans, 1902).

2.6 The Spherical Collapse Model

As gravity is attractive force, ambient matter falls into such high density regions, which results in formation of halos. The spherical collapse model (Gunn & Gott, 1972) provides a simplified description of the formation of cosmic structures by considering the evolution of a spherically symmetric overdensity in an expanding universe. Suppose that there is spherical matter around a certain point in the Universe and initial density contrast is denoted as $\delta_i \ll 1$. The equation of motion of the shell which the radius R is given by:

$$\ddot{R} = -\frac{GM(< R)}{R^2}, \quad (2.67)$$

where $M(< R)$ is the mass enclosed within the radius R . Multiplying both sides by $\dot{R}$ and integrating over time, we get:

$$\dot{R}^2 = \frac{2GM(< R)}{R} + E. \quad (2.68)$$

The constant E corresponds to the energy, $E < 0$ for bound systems. From this expression, we can obtain a parametric solution for the radius $R(t)$ in term of θ :

$$R(t) = (GM)^{1/3} A^2 (1 - \cos \theta) \quad (2.69)$$

$$t = A^3 (\theta - \sin \theta) \quad (2.70)$$

where A is a constant. For a matter-dominated universe, the mean density $\bar{\rho} = (6\pi G t^2)^{-1}$. The density contrast within the shell is given by:

$$1 + \delta = \frac{\rho}{\bar{\rho}} = \frac{3M}{4\pi R^3} \frac{6\pi G t^2}{M} = \frac{9}{2} \frac{(\theta - \sin \theta)^2}{(1 - \cos \theta)^3} \quad (2.71)$$

$\theta = \pi$ corresponds to the turnaround point, where the shell reaches its maximum radius and starts collapsing. The density contrast at the turnaround point is given by:

$$\delta_{ta} = \frac{9\pi^2}{16} - 1 \approx 4.55 \quad (2.72)$$

where the corresponding radius R_{ta} and time t_{ta} are:

$$R_{ta} = 2(GM)^{1/3} A^2, \quad t_{ta} = \pi A^3 \quad (2.73)$$

At $\theta = 2\pi$, the shell reaches a singularity where the radius R goes to zero and the density contrast δ diverges. However, in reality, this does not occur because the shell undergoes virialization and forms a halo. We assume the shell virializes at $R = R_{ta}/2$ at time $t = t_{coll}$. The density contrast at virialization is given by:

$$\delta_{coll} = \delta_{ta} \times 4 \times 2^3 = 18\pi^2. \quad (2.74)$$

In the early epoch ($\theta \ll 1$), the density contrast follows the linear theory. If we expand the density contrast and time around $\theta = 0$, we get:

$$\delta = \frac{3}{20} \theta^2 + \mathcal{O}(\theta^4), \quad t = \frac{A^3}{6} \theta^3 + \mathcal{O}(\theta^5) \quad (2.75)$$

This yields $\delta \propto t^{2/3}$, which is consistent with linear theory. Denoting this linear fluctuation as $\delta_L(t)$:

$$\delta_L(t) = \frac{3(6t)^{2/3}}{20A^2} \quad (2.76)$$

Substituting $t = t_{\text{ta}}$ and $t = t_{\text{coll}}$, we obtain:

$$\delta_L(t_{\text{ta}}) = \frac{3(6\pi)^{2/3}}{20} \approx 1.06, \quad \delta_L(t_{\text{coll}}) = \frac{3(12\pi)^{2/3}}{20} \approx 1.69. \quad (2.77)$$

Therefore, when the linear density contrast exceeds $\delta_L \approx 1.69$, the shell virializes and forms a halo.

2.7 Dark Matter Halos

Dark matter halos are the fundamental building blocks of cosmic structures. They form through the gravitational collapse of overdense regions in the early universe and provide the potential wells in which baryonic matter accumulates to form galaxies and galaxy clusters.

2.7.1 The Halo Mass Function: Press-Schechter

The halo mass function (HMF) describes the number density of dark matter halos as a function of their mass and redshift. The Press-Schechter (PS) formalism ([Press & Schechter, 1974](#)) provides an analytical approach to calculate the HMF based on the initial Gaussian density field and the theory of gravitational collapse.

Let us consider the density field which follows Gaussian distribution at each position. The probability distribution function is:

$$P(\delta)d\delta = \frac{1}{\sqrt{2\pi}\sigma^2} \exp\left(-\frac{\delta^2}{2\sigma^2}\right) d\delta \quad (2.78)$$

where σ^2 is the variance of the density field. Supposed that the sphere of radius R contains mass $M = \frac{4}{3}\pi R^3 \bar{\rho}$, where $\bar{\rho}$ is the mean density. The density contrast within this sphere is:

$$\delta_M(\mathbf{q}, t) = \frac{3}{4\pi} \int_{|\mathbf{q}' - \mathbf{q}| < R} \delta(\mathbf{q}', t) d^3 \mathbf{q}' \quad (2.79)$$

This density contrast follows Gaussian distribution. The probability distribution function of the density contrast is:

$$P(\delta_M) = \frac{1}{\sqrt{2\pi}\sigma^2(M)} \exp\left(-\frac{\delta_M^2}{2\sigma^2(M)}\right) \quad (2.80)$$

The halo formation happens when the density contrast exceeds a critical value δ_c . The fraction of Lagrangian volume which collapses to form halos is:

$$P_{>\delta_c} = \int_{\delta_c}^{\infty} P(\delta_M) d\delta_M = \frac{1}{\sqrt{2\pi}} \int_{\delta_c/\sigma(M)}^{\infty} e^{-x^2/2} dx \quad (2.81)$$

Thus, mass which forms dark halo with more than mass M can be calculated as $\bar{\rho}_0 P_{>\delta_c}$. In Press-

Schechter formalism, the mass function is given by:

$$n(M)M dM = 2\bar{\rho}_0 \left| \frac{P_{\delta_c}}{d\sigma(M)} \right| \left| \frac{d\sigma(M)}{dM} \right| dM \quad (2.82)$$

Substituting Eq. (2.81) into the above equation, we obtain:

$$n(M) = \sqrt{\frac{2}{\pi}} \frac{\bar{\rho}_0}{M} \frac{\delta_c}{\sigma^2(M)} \left| \frac{d\sigma}{dM} \right| e^{-\delta_c^2/2\sigma^2(M)} \quad (2.83)$$

The mass function can be further simplified by using the relation between the variance of the density field and the linear power spectrum $P(k)$:

$$\sigma^2(M) = \int \frac{d^3k}{(2\pi)^3} P(k) W^2(kR) \quad (2.84)$$

where $W(kR)$ is the Fourier transform of the top-hat window function:

$$\begin{aligned} W(kR) &= \int d^3x e^{i\mathbf{k}\cdot\mathbf{x}} W_R(\mathbf{x}) \\ &= 4\pi \int_0^R x^2 dx \frac{\sin(kx)}{kx} W_R(x) \\ &= \frac{3}{(kR)^3} [\sin(kR) - kR \cos(kR)] \end{aligned} \quad (2.85)$$

Finally, we can construct the Press-Schechter mass function only from the linear power spectrum $P(k)$. In reality, matter collapses into halos non-spherically. One of the most popular extensions of the Press-Schechter formalism is the Sheth-Tormen mass function (Sheth & Tormen, 1999), which provides a better fit to numerical simulations by incorporating ellipsoidal collapse.

2.7.2 Halo Bias

Halo bias quantifies how dark matter halos are biased tracers of the underlying matter distribution. Massive halos tend to form in regions of higher density contrast, leading to a mass-dependent bias factor $b(M)$ which is independent for a convenient scale. Mo & White (1996) proposed a simple model for halo bias based on the Press-Schechter formalism and the spherical collapse model. They derive the analytical expression for the linear halo bias, which defined as:

$$b_h(M, z) := \frac{\delta_h(M, z)}{\delta_m} \quad (2.86)$$

where $\delta_h(M, z)$ is the overdensity of halos of mass M at redshift z , and δ_m is the overdensity of the matter field. In the lowest-order approximation, the linear halo bias is given by:

$$b_h(M, z) = 1 + \frac{\nu(M, z)^2 - 1}{\delta_c} \quad (2.87)$$

where $\nu(M, z) = \delta_c/\sigma(M, z)$ is the peak height, and δ_c is the critical density contrast for collapse. It is then extended by [Sheth et al. \(2001\)](#) to include ellipsoidal collapse.

2.7.3 Halo Density Profiles: NFW and Einasto Models

Since halos undergo nonlinear gravitational evolution, their density profiles need numerical simulations to be determined. The most widely used model is the Navarro-Frenk-White (NFW) profile ([Navarro et al., 1996, 1997](#)), which is found universally in numerical simulations. The NFW profile is given by:

$$\rho_{\text{NFW}}(r) = \frac{\rho_s}{(r/r_s)(1 + r/r_s)^2} \quad (2.88)$$

where ρ_s and r_s are the characteristic density and scale radius, respectively. At large radii, the NFW profile follows a power-law behavior $\rho_{\text{NFW}} \propto r^{-3}$, while at small radii, it behaves as $\rho_{\text{NFW}} \propto r^{-1}$. The enclosed mass is not well defined for the NFW profile, so the virial radius r_{vir} and r_{200c} , which encloses a density contrast of 200 times the critical density, are used instead.

After specifying the boundary, one can compute the halo mass as ([Oguri & Hamana, 2011](#)):

$$M_{\text{vir}} = 4\pi r_s^3 \rho_s m(c_{\text{vir}}) \quad (2.89)$$

with

$$m(c) = \int_0^c \frac{x dx}{(1+x)^2} = \ln(1+c) - \frac{c}{1+c} \quad (2.90)$$

where $c_{\text{vir}} = r_{\text{vir}}/r_s$ is the concentration parameter.

Another widely used profile is the Einasto profile ([Einasto, 1965](#)), which provides an alternative fitting function with more accuracy at relatively small halo radius. The Einasto profile is given by:

$$\rho_{\text{Ein}}(r) = \rho_s \exp\left(-\frac{2}{\alpha} \left[\left(\frac{r}{r_s}\right)^\alpha - 1\right]\right) \quad (2.91)$$

where ρ_s and r_s are the characteristic density and scale radius, respectively, and α is the shape parameter.

Theoretical Foundations of Weak Gravitational Lensing

Contents

3.1	Introduction to Weak Gravitational Lensing	31
3.2	Convergence Field	34
3.3	Shear Field	36
3.4	Lensing Estimation	37

Weak gravitational lensing (WL), also known as cosmic shear, refers to the subtle distortions in the images of distant source caused by the gravitational fields of intervening mass distributions. Unlike strong lensing, which produces noticeable effects such as multiple images or arcs, weak lensing induces small, coherent distortions that require statistical analysis to detect and interpret. For the standard approach to lensing, we refer to [Schneider et al. \(1992\)](#), [Bartelmann & Schneider \(2001\)](#) and [Bartelmann \(2010\)](#).

3.1 Introduction to Weak Gravitational Lensing

3.1.1 Derivation of the Lens Equation

To derive the lens equation, we consider a perturbed FLRW metric, which incorporates gravitational potential perturbations. The metric is expressed as

$$ds^2 = - \left(1 + \frac{2\Phi}{c^2} \right) c^2 dt^2 + a^2(t) \left(1 - \frac{2\Psi}{c^2} \right) [d\chi^2 + f_K^2(\chi) \omega_{ab} dx^a dx^b] \quad (a, b = 2, 3), \quad (3.1)$$

where the angular part of the metric is defined by

$$\omega_{ab} dx^a dx^b := d\theta^2 + \sin^2 \theta d\phi^2. \quad (3.2)$$

In this context, Φ and Ψ represent the scalar gravitational potentials, and $f_K(\chi)$ encodes the spatial curvature as previously defined in Eq. (2.18).

The trajectory of light within this spacetime is governed by the geodesic equation, which is given by

$$\frac{d^2 x^\mu}{d\lambda^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = 0, \quad (3.3)$$

where λ is an affine parameter and $\Gamma_{\alpha\beta}^\mu$ are the Christoffel symbols corresponding to the metric in Equation (3.1).

To facilitate the derivation, we reparametrize the geodesic equation by substituting the affine parameter λ with the comoving radial distance χ . Applying the chain rule, the geodesic equation transforms to

$$\frac{d^2 x^\mu}{d\chi^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\chi} \frac{dx^\beta}{d\chi} - \frac{d^2 \lambda}{d\chi^2} \left(\frac{d\lambda}{d\chi} \right)^{-1} \frac{dx^\mu}{d\chi} = 0. \quad (3.4)$$

Setting $\mu = 1$ (where $x^1 = \chi$) in Equation (3.4) and simplifying, we obtain

$$\frac{d^2 x^\mu}{d\chi^2} + \left(\Gamma_{\alpha\beta}^\mu - \Gamma_{\alpha\beta}^1 \right) \frac{dx^\alpha}{d\chi} \frac{dx^\beta}{d\chi} = 0. \quad (3.5)$$

The evaluation of Equation (3.5) requires the computation of the Christoffel symbols. Additionally, the derivative $c dt/d\chi$ is derived from the null condition

$$g_{\mu\nu} \frac{dx^\mu}{d\chi} \frac{dx^\nu}{d\chi} = 0, \quad (3.6)$$

yielding

$$\frac{c dt}{d\chi} = -a(t) \left[1 - \frac{\Phi}{c^2} - \frac{\Psi}{c^2} + \frac{f_K^2(\chi)}{2} \omega_{ab} \frac{dx^a}{d\chi} \frac{dx^b}{d\chi} \right], \quad (3.7)$$

valid to first order in Φ and second order in $dx^a/d\chi$. Notably, for the evaluation of Equation (3.5), only the zeroth-order term $c dt/d\chi = -a(t)$ is required. However, the inclusion of perturbative terms in Equation (3.7) is essential for subsequent derivations of the lens equation.

Focusing on the angular components ($\mu = a$) of Equation (3.5), we derive the following differential equation:

$$\frac{d^2 x^a}{d\chi^2} + 2 \frac{f'_K(\chi)}{f_K(\chi)} \frac{dx^a}{d\chi} + \omega^{ab} \frac{\Phi_b + \Psi_b}{c^2 f_K^2(\chi)} = 0, \quad (3.8)$$

where Φ_b and Ψ_b denote the derivatives of the gravitational potentials with respect to the angular

coordinates x^a . Integrating Equation (3.8) twice with respect to χ , we obtain

$$\begin{aligned}
 x^a(\chi_s) - x^a(0) &= -\frac{1}{c^2} \int_0^{\chi_s} d\chi' \int_0^{\chi'} d\chi \omega_{ab} [\Phi_b(\chi, \theta(\chi)) + \Psi_b(\chi, \theta(\chi))] \\
 &= -\frac{1}{c^2} \int_0^{\chi_s} d\chi \omega_{ab} [\Phi_b(\chi, \theta(\chi)) + \Psi_b(\chi, \theta(\chi))] \int_{\chi}^{\chi_s} d\chi' \frac{1}{f_K^2(\chi')} \\
 &= -\frac{1}{c^2} \int_0^{\chi_s} d\chi \omega_{ab} [\Phi_b(\chi, \theta(\chi)) + \Psi_b(\chi, \theta(\chi))] \frac{f_K(\chi_s - \chi)}{f_K(\chi_s) f_K(\chi)}, \quad (3.9)
 \end{aligned}$$

where χ_s denotes the comoving radial distance to the source, and $x^a(0)$ is the angular position at the observer's location. Defining $(\nabla_\theta)^a = \omega^{ab} \partial_b$, the angular position of the source galaxy β is related to the observed angular position θ by

$$\theta(\chi_s) = \theta(0) - \frac{1}{c^2} \int_0^{\chi_s} d\chi \nabla_\theta [\Phi(\chi, \theta(\chi)) + \Psi(\chi, \theta(\chi))] \frac{f_K(\chi_s - \chi)}{f_K(\chi_s) f_K(\chi)}. \quad (3.10)$$

Applying the Born approximation (Born, 1926), and assume that $\Phi = \Psi$, we can simplify the lens equation to:

$$\theta(\chi_s) = \theta(0) - \frac{2}{c^2} \int_0^{\chi_s} d\chi \nabla_\theta [\Phi(\chi, \theta(0))] \frac{f_K(\chi_s - \chi)}{f_K(\chi_s) f_K(\chi)}. \quad (3.11)$$

This final expression constitutes the lens equation, encapsulating the deflection of light due to the gravitational potentials Φ and Ψ along the line of sight.

3.1.2 The Jacobian (Lensing) Matrix $\mathcal{A}$




Figure 3.1: Schematic representation of the lensing geometry. The source is located at β , while the observed image is at θ . The deflection angle α is the difference between the observed and true angular positions.

Redefining the notation in Eq. (3.11) and considering the angular position of the source $\beta = \theta(\chi_s)$ and the observed angular position $\theta = \theta(0)$ (see Fig. 3.1), we can express the lens equation as (Bartelmann & Schneider, 2001; Hilbert et al., 2009; Kilbinger, 2015):

$$\beta = \theta - \alpha(\theta), \quad (3.12)$$

where the deflection angle $\alpha(\theta)$ is defined by:

$$\alpha(\theta) = \nabla_{\theta} \psi(\theta), \quad \psi(\theta) = \frac{2}{c^2} \int_0^{\chi_s} d\chi \frac{f_K(\chi_s - \chi)}{f_K(\chi_s) f_K(\chi)} \Phi(f_K(\chi) \theta, \chi). \quad (3.13)$$

The mapping between the source plane and the image plane can be described by the Jacobian matrix $\mathcal{A}$, which relates infinitesimal displacements in the source position to displacements in the image position:

$$\mathcal{A} := \frac{\partial \beta}{\partial \theta} = \begin{pmatrix} 1 - \kappa - \gamma_1 & -\gamma_2 \\ -\gamma_2 & 1 - \kappa + \gamma_1 \end{pmatrix} = (1 - \kappa) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - |\gamma| \begin{pmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \quad (3.14)$$

where κ is the convergence and $\gamma = \gamma_1 + i\gamma_2 = |\gamma|e^{2i\phi}$ is the shear. The quantities κ and γ will be discussed in detail in the subsequent sections. Figure 3.2 illustrates the effects of gravitational lensing on the shapes of background sources through the lensing matrix $\mathcal{A}$. The panels demonstrate how the combined effects of convergence and shear components in the Jacobian matrix $\mathcal{A}$ lead to complex distortions of background sources.

3.2 Convergence: Mass Distribution to Lensing

3.2.1 Definition of Convergence in Real and Fourier Space

From the lensing matrix in Eq. (3.14), the convergence κ is defined as:

$$\kappa(\theta) := \frac{1}{2} \left(\frac{\partial^2 \psi}{\partial \theta_1^2} + \frac{\partial^2 \psi}{\partial \theta_2^2} \right) = \frac{1}{2} \nabla_{\theta}^2 \psi(\theta) \quad (3.15)$$

with θ_1 and θ_2 representing the angular coordinates on the sky. In Fourier space, the convergence field could be expressed as:

$$\tilde{\kappa}(\ell) = \int d^2\theta e^{-i\ell \cdot \theta} \kappa(\theta) = \frac{1}{2} \ell^2 \tilde{\psi}(\ell), \quad (3.16)$$

where $(\tilde{})$ denotes the Fourier transform of the corresponding quantity and $\ell = |\ell|$ is the Fourier counterpart to the angular position θ .




Figure 3.2: Illustration of the distortion of background sources due to gravitational lensing. The left panel depicts the effect of the convergence κ and the middle and right panels show the components of the shear $\gamma = \gamma_1 + i\gamma_2$ on circular background sources. Positive and negative values of κ cause isotropic magnification or demagnification, while γ_1 and γ_2 introduce anisotropic distortions, stretching the sources along or at an angle to the principal axes.

3.2.2 Convergence from Density Contrast and the Lensing Efficiency Function

In a flat universe, the Poisson equation in comoving coordinates is expressed as

$$\begin{aligned}\nabla_{\mathbf{x}}^2 \Phi(\mathbf{x}, \chi) &= 4\pi G a^2(\chi) \bar{\rho}_m(\chi) \delta(\mathbf{x}, \chi) \\ &= 4\pi G a^2(\chi) \left[\frac{3H_0^2 \Omega_m}{8\pi G} a^{-3}(\chi) \right] \delta(\mathbf{x}, \chi) \\ &= \frac{3}{2} \Omega_m H_0^2 a^{-1}(\chi) \delta(\mathbf{x}, \chi),\end{aligned}\tag{3.17}$$

where we utilized Eq. (2.35) and Eq. (2.27). Substituting the expression for Φ from Eq. (3.17) into the lensing potential (Eq. (3.13)) and subsequently into the convergence (Eq. (3.15)), we derive:

$$\begin{aligned}\kappa(\boldsymbol{\theta}) &= \frac{1}{c^2} \int_0^{\chi_s} d\chi \frac{f_K(\chi_s - \chi)}{f_K(\chi_s) f_K(\chi)} \left[\frac{1}{f_K^2(\chi)} \nabla_{\mathbf{x}}^2 \Phi(\mathbf{x}, \chi) \right] \\ &= \int_0^{\chi_s} d\chi \frac{3\Omega_m H_0^2}{2c^2} \frac{f_K(\chi_s - \chi) f_K(\chi)}{f_K(\chi_s)} \frac{a^{-1}(\chi)}{f_K^3(\chi)} \delta(\mathbf{x}, \chi) \\ &= \int_0^{\chi_s} d\chi W(\chi, \chi_s) \delta(\mathbf{x}, \chi),\end{aligned}\tag{3.18}$$

where the lensing efficiency function $W(\chi, \chi_s)$ is defined by:

$$W(\chi, \chi_s) := \frac{3\Omega_m H_0^2}{2c^2} \frac{f_K(\chi) f_K(\chi_s - \chi)}{f_K(\chi_s)} \frac{a^{-1}(\chi)}{f_K^2(\chi)}.\tag{3.19}$$

In a flat universe ($f_K(\chi) = \chi$), this simplifies to:

$$W(\chi, \chi_s) = \frac{3\Omega_m H_0^2}{2c^2} a^{-1}(\chi) \frac{\chi(\chi_s - \chi)}{\chi_s}. \quad (3.20)$$

3.3 Shear: Anisotropic Distortions of Background Sources

The shear γ encapsulates the anisotropic stretching of galaxy images induced by gravitational lensing. Unlike convergence, which affects the size and brightness of images isotropically, shear induces distortions that alter the shapes of background galaxies coherently.

3.3.1 Definition of Shear

[Shear Components and Complex Representation] The shear components γ_1 and γ_2 describe distortions along different axes and are related to the lensing potential ψ by:

$$\gamma_1 := \frac{1}{2} \left(\frac{\partial^2 \psi}{\partial \theta_1^2} - \frac{\partial^2 \psi}{\partial \theta_2^2} \right), \quad \gamma_2 := \frac{\partial^2 \psi}{\partial \theta_1 \partial \theta_2}. \quad (3.21)$$

In Fourier space, the shear field can be expressed as:

$$\tilde{\gamma}_1(\ell) = \frac{1}{2} (\ell_1^2 - \ell_2^2) \tilde{\psi}(\ell) \quad \tilde{\gamma}_2(\ell) = \ell_1 \ell_2 \tilde{\psi}(\ell), \quad (3.22)$$

where $\tilde{\gamma}_1(\ell)$ and $\tilde{\gamma}_2(\ell)$ are the Fourier transforms of $\gamma_1(\theta)$ and $\gamma_2(\theta)$, respectively. Similar as Eq. (3.14), the shear field can be expressed in complex form as:

$$\tilde{\gamma}(\ell) := \tilde{\gamma}_1 + i\tilde{\gamma}_2 = |\tilde{\gamma}(\ell)| e^{2i\phi_\ell}, \quad \tan(2\phi_\ell) = \tilde{\gamma}_2/\tilde{\gamma}_1. \quad (3.23)$$

Therefore, the shear field in Fourier space is directly related to the convergence field combining Eq.(3.16) and Eq.(3.22) (Kaiser & Squires, 1993):

$$\tilde{\kappa}(\ell) = \frac{\ell_1^2 - \ell_2^2 + 2i\ell_1 \ell_2}{\ell^2} \tilde{\gamma}(\ell). \quad (3.24)$$

3.3.2 E-mode and B-mode Decomposition

The shear field can be decomposed into two distinct modes: the **E-mode** (gradient component) and the **B-mode** (curl component). By rotating the complex shear field to align with the principal axes of the shear $\tilde{\gamma}_{EB} = e^{-2i\phi_l} \tilde{\gamma}$, we can express the shear field in terms of the E-mode and B-mode components:

$$\begin{pmatrix} \tilde{\gamma}_E \\ \tilde{\gamma}_B \end{pmatrix} := \begin{pmatrix} \cos 2\phi_l & \sin 2\phi_l \\ -\sin 2\phi_l & \cos 2\phi_l \end{pmatrix} \begin{pmatrix} \tilde{\gamma}_1 \\ \tilde{\gamma}_2 \end{pmatrix} \quad (3.25)$$

where $\tilde{\gamma}_E$ and $\tilde{\gamma}_B$ are the E-mode and B-mode components of the shear, respectively. The E-mode represents the gradient component of the shear field, while the B-mode describes the curl component.

For standard gravitational lensing by density fluctuations, the B-mode component $\tilde{\gamma}_B(\ell)$ is expected to vanish in the absence of systematics or additional physical effects. This implies that all the shear signal is contained within the E-mode such that:

$$\tilde{\gamma}_E(\ell) = \tilde{\kappa}(\ell), \quad \tilde{\gamma}_B(\ell) = 0. \quad (3.26)$$

3.4 Estimating Lensing Fields from Observables

In the case of cosmic shear, not the convergence but the shear is measured from the observed galaxy shapes. Theoretical predictions of the convergence can be related to the observed shear using the relationship in Eq. (3.24). Here, we introduce a concept of the reduced shear. Furthermore, a convergence field can be estimated from the observed galaxy shapes (Kaiser & Squires, 1993) and can be estimated from magnification (Bartelmann & Schneider, 2001).

3.4.1 Galaxy Ellipticity and Reduced Shear

To quantify the shapes of galaxies, we use the second moments of their surface brightness distributions $I(\theta)$. For each galaxy, the second moments Q_{ab} are defined as (Bartelmann & Schneider, 2001):

$$Q_{ab} := \frac{\int d^2\theta I(\theta) \theta_a \theta_b}{\int d^2\theta I(\theta)}, \quad (3.27)$$

where $\theta = (\theta_1, \theta_2)$ is the angular position relative to the galaxy center. The complex ellipticity ϵ of the galaxy is then defined as:

$$\epsilon := \frac{Q_{11} - Q_{22} + 2iQ_{12}}{Q_{11} + Q_{22}}. \quad (3.28)$$

Gravitational lensing transforms the image of a source galaxy via the Jacobian matrix A (see Eq. (3.14)). Assuming that the surface brightness is conserved during lensing, $I^{(s)}(\beta) = I(\theta)$, the second moments in the source plane $Q_{ab}^{(s)}$ are related to those in the image plane by:

$$Q_{ab}^{(s)} = \frac{\int d^2\beta I^{(s)}(\beta) \beta_a \beta_b}{\int d^2\beta I^{(s)}(\beta)} \approx A_{ac} A_{bd} Q_{cd}, \quad (3.29)$$

where we have approximated the size of the galaxy as sufficiently small so that the lensing distortion is constant across the galaxy image.

By expanding the components of $Q_{ab}^{(s)}$ and performing straightforward calculations (see Schneider et al. 1992 for details), we find that the intrinsic ellipticity $\epsilon^{(s)}$ is related to the observed ellipticity ϵ

through:

$$\epsilon^{(s)} = \frac{(1 - \kappa)^2 \epsilon - 2(1 - \kappa)\gamma + \gamma^2 \epsilon^*}{(1 - \kappa)^2 + |\gamma|^2 - 2(1 - \kappa)\text{Re}[\gamma \epsilon^*]}, \quad (3.30)$$

where ϵ^* denotes the complex conjugate of ϵ , and Re denotes the real part.

Introducing the *reduced shear* $g = \gamma/(1 - \kappa)$, the above equation simplifies to (Schneider & Seitz, 1995):

$$\epsilon^{(s)} = \frac{\epsilon - 2g + g^2 \epsilon^*}{1 + |g|^2 - 2\text{Re}[g \epsilon^*]}. \quad (3.31)$$

This relation indicates that weak lensing measurements are sensitive to the reduced shear g rather than the shear γ directly. In the weak lensing regime, where $|\kappa|, |\gamma| \ll 1$, and assuming that the intrinsic ellipticities of galaxies are randomly oriented (i.e., $\langle \epsilon^{(s)} \rangle = 0$), the observed ellipticity becomes an unbiased estimator of the reduced shear:

$$\langle \epsilon \rangle \approx g \approx \gamma. \quad (3.32)$$

However, in the weak-lensing regime, the shear cannot be detected from an individual galaxy due to the dominance of intrinsic shape noise. The typical root mean square (rms) of the intrinsic ellipticity is $\sigma_\epsilon \approx 0.26$ (Euclid Collaboration et al., 2019), which necessitates averaging over a large number of galaxies to measure the shear signal accurately. The noise for the reduced shear estimator is dominated by Poisson noise, which is given by:

$$\sigma_\gamma = \frac{\sigma_\epsilon}{\sqrt{N}}, \quad (3.33)$$

where N is the number of galaxies used in the shear estimation. It is known that the following transformation of convergence does not change the reduced shear,

$$\kappa' = \lambda \kappa + (1 - \lambda), \quad (3.34)$$

where λ is a arbitrary constant. This degree of freedom in the convergence field is known as the mass-sheet degeneracy (Falco et al., 1985).

3.4.2 Inversion Techniques: From Shear to Convergence

As we have seen in Eq. (3.24), shear and convergence are related through the Fourier transform. Following Kaiser & Squires (1993), the relation between the shear and convergence fields in real space can be expressed as:

$$\gamma(\boldsymbol{\theta}) = \frac{1}{\pi} \int d^2 \theta' \mathcal{D}(\boldsymbol{\theta} - \boldsymbol{\theta}') \kappa(\boldsymbol{\theta}'), \quad (3.35)$$

where $\mathcal{D}(\boldsymbol{\theta})$ is a kernel function defined by:

$$\mathcal{D}(\boldsymbol{\theta}) = -\frac{\theta_1^2 - \theta_2^2 + 2i\theta_1\theta_2}{\boldsymbol{\theta}^4} = -\frac{1}{(\theta_1 - i\theta_2)^2}. \quad (3.36)$$

Therefore, the convolution with the kernel function in Fourier space yields:

$$\tilde{\gamma}(\boldsymbol{\ell}) = \frac{1}{\pi} \tilde{\mathcal{D}}(\boldsymbol{\ell}) \tilde{\kappa}(\boldsymbol{\ell}), \quad (3.37)$$

The Fourier transform of the kernel function, $\tilde{\mathcal{D}}(\boldsymbol{\ell})$, is given by:

$$\tilde{\mathcal{D}}(\boldsymbol{\ell}) = \pi \frac{\ell_1^2 - \ell_2^2 + 2i\ell_1\ell_2}{\ell^2} \quad (3.38)$$

It is notable that this kernel function and its conjugate satisfies:

$$\tilde{\mathcal{D}}(\boldsymbol{\ell}) \tilde{\mathcal{D}}^*(\boldsymbol{\ell}) = \pi^2 \left(\frac{\ell_1^2 - \ell_2^2 + 2i\ell_1\ell_2}{\ell^2} \right) \left(\frac{\ell_1^2 - \ell_2^2 - 2i\ell_1\ell_2}{\ell^2} \right) = \pi^2. \quad (3.39)$$

Substituting the expression for $\tilde{\mathcal{D}}(\boldsymbol{\ell})$ into the relation between the shear and convergence fields, we find:

$$\tilde{\kappa}(\boldsymbol{\ell}) = \frac{1}{\pi} \tilde{\gamma}(\boldsymbol{\ell}) \tilde{\mathcal{D}}^*(\boldsymbol{\ell}). \quad (3.40)$$

$$\kappa(\boldsymbol{\theta}) = \kappa_0 + \frac{1}{\pi} \int d^2\theta' \mathcal{D}^*(\boldsymbol{\theta} - \boldsymbol{\theta}') \gamma(\boldsymbol{\theta}'). \quad (3.41)$$

The constant term κ_0 arises from the unknown $\ell = 0$ mode in the Fourier space, which is not constrained by the observed shear field.

Summary Statistics in Weak Lensing

Contents

4.1	Two-point Correlation Function	40
4.2	Convergence Power Spectrum	41
4.3	Convergence Bispectrum	44
4.4	Probability Density Functions	44
4.5	Peak and Minimum Counts	46
4.6	Minkowski Functionals	47

4.1 Two-point Correlation Function

The two-point correlation function (2PCF), denoted by $\xi(\mathbf{r})$, is a fundamental statistical tool in cosmology that characterizes the clustering of matter in the Universe. It quantifies the excess probability of finding a pair of points separated by a distance $\mathbf{r}$ compared to a random distribution ([Bernardeau et al., 2002](#)), defined as:

$$\xi(\mathbf{r}) = \langle \delta(\mathbf{x})\delta(\mathbf{x} + \mathbf{r}) \rangle, \quad (4.1)$$

where $\delta(\mathbf{x})$ represents the matter density contrast at position $\mathbf{x}$, and $\langle \cdot \rangle$ denotes the ensemble average over all positions in the Universe.

In cosmological analyses, the tracers of the underlying matter distribution, such as galaxies or clusters of galaxies, are often biased. This means their spatial distribution does not perfectly follow that of the dark matter. To accurately model the 2PCF of these biased tracers, the concept of halo bias is introduced (see Section 2.7). For linear bias models, the relationship between the biased tracers and the underlying matter density contrast is expressed as:

$$\delta_h(\mathbf{x}) = b_1 \delta(\mathbf{x}), \quad (4.2)$$

where $\delta_h(\mathbf{x})$ is the density contrast of the biased tracers, and b_1 is the linear bias parameter. Substituting Equation (4.2) into Equation (4.1), the biased 2PCF, $\xi_h(\mathbf{r})$, becomes (Mo & White, 1996):

$$\xi_h(\mathbf{r}) = b_1^2 \xi(\mathbf{r}). \quad (4.3)$$

This relation indicates that the clustering strength of biased tracers is amplified by the square of the bias parameter, b_1^2 , compared to the underlying matter distribution. Therefore, understanding and accurately modeling this bias is crucial for interpreting observational data.

4.2 Convergence Power Spectrum

The matter power spectrum, $P(k)$, is a fundamental quantity in cosmology that characterizes the distribution of dark matter density fluctuations in Fourier space. It is defined as the Fourier transform of the two-point correlation function of the dark matter density field, $\delta(\mathbf{x})$ (Bartelmann & Schneider, 2001). Mathematically, the matter power spectrum is expressed as:

$$\langle \tilde{\delta}(\mathbf{k}) \tilde{\delta}(\mathbf{k}') \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{k}') P(k), \quad (4.4)$$

where $\tilde{\delta}(\mathbf{k})$ represents the Fourier transform of the density contrast $\delta(\mathbf{x})$, and $\delta^{(3)}$ is the three-dimensional Dirac delta function ensuring statistical isotropy and homogeneity.

In the context of weak gravitational lensing, the matter power spectrum $P(k)$ is not directly observable. Instead, observations yield the angular power spectrum of the convergence field, $C_\ell^{\kappa\kappa}$, which encapsulates the statistical properties of the convergence $\kappa(\boldsymbol{\theta})$ across the sky (Bartelmann & Schneider, 2001). The convergence power spectrum, $C_\ell^{\kappa\kappa}$, is defined through the relation:

$$\langle \tilde{\kappa}(\boldsymbol{\ell}) \tilde{\kappa}(\boldsymbol{\ell}') \rangle = (2\pi)^2 \delta^{(2)}(\boldsymbol{\ell} + \boldsymbol{\ell}') C_\ell^{\kappa\kappa}, \quad (4.5)$$

where $\tilde{\kappa}(\boldsymbol{\ell})$ is the Fourier transform of the convergence field $\kappa(\boldsymbol{\theta})$, and $\delta^{(2)}$ is the two-dimensional Dirac delta function.

4.2.1 Derivation of the Convergence Power Spectrum

The convergence field $\kappa(\boldsymbol{\theta})$ can be expressed as a weighted projection of the matter density contrast along the line of sight (see Eq. (3.18)):

$$\kappa(\boldsymbol{\theta}) = \int_0^{\chi_s} d\chi W(\chi) \delta_m(\chi \boldsymbol{\theta}, \chi), \quad (4.6)$$

where $W(\chi)$ is the lensing kernel, χ is the comoving radial distance.

Recognizing the Fourier transform of the matter density field $\delta_m(\chi\boldsymbol{\theta}, \chi)$, we write:

$$\delta_m(\chi\boldsymbol{\theta}, \chi) = \int \frac{d^3\mathbf{k}}{(2\pi)^3} \tilde{\delta}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}}, \quad (4.7)$$

where $\mathbf{x} = (\chi\boldsymbol{\theta}, \chi)$ is the comoving position vector. Substituting this into the Fourier transform of the convergence field $\tilde{\kappa}(\boldsymbol{\ell})$, we obtain:

$$\begin{aligned} \tilde{\kappa}(\boldsymbol{\ell}) &= \int_0^{\chi_s} d\chi W(\chi) \int d\boldsymbol{\theta} e^{-i\boldsymbol{\ell}\cdot\boldsymbol{\theta}} \delta_m(\chi\boldsymbol{\theta}, \chi) \\ &= \int_0^{\chi_s} d\chi W(\chi) \int \frac{d^3\mathbf{k}}{(2\pi)^3} \tilde{\delta}(\mathbf{k}) e^{ik_{\parallel}\chi} \int d\boldsymbol{\theta} e^{-i\boldsymbol{\ell}\cdot\boldsymbol{\theta}} e^{i\chi\mathbf{k}_{\perp}\cdot\boldsymbol{\theta}} \\ &= \int_0^{\chi_s} d\chi W(\chi) \int \frac{dk_{\parallel}}{2\pi} \frac{d^2\mathbf{k}_{\perp}}{(2\pi)^2} \tilde{\delta}_m(k_{\parallel}, \mathbf{k}_{\perp}) e^{ik_{\parallel}\chi} \int d\boldsymbol{\theta} e^{-i(\boldsymbol{\ell}-\chi\mathbf{k}_{\perp})\cdot\boldsymbol{\theta}} \\ &= \int_0^{\chi_s} d\chi \frac{W(\chi)}{\chi^2} \int \frac{dk_{\parallel}}{2\pi} \tilde{\delta}_m\left(k_{\parallel}, \frac{\boldsymbol{\ell}}{\chi}\right) e^{ik_{\parallel}\chi}, \end{aligned} \quad (4.8)$$

where $\mathbf{k}_{\perp}$ and $k_{\parallel}$ are the components of $\mathbf{k}$ perpendicular and parallel to the line of sight, respectively.

Next, we evaluate the ensemble average of the Fourier transform of the convergence field. Starting from the definition, we have:

$$\begin{aligned} \langle \tilde{\kappa}(\boldsymbol{\ell}) \tilde{\kappa}(\boldsymbol{\ell}') \rangle &= \int_0^{\chi_s} d\chi \frac{W(\chi)}{\chi^2} \int_0^{\chi_s} d\chi' \frac{W(\chi')}{\chi'^2} \int \frac{dk_{\parallel}}{2\pi} \int \frac{dk'_{\parallel}}{2\pi} \\ &\quad \times \langle \tilde{\delta}_m\left(k_{\parallel}, \frac{\boldsymbol{\ell}}{\chi}\right) \tilde{\delta}_m\left(k'_{\parallel}, \frac{\boldsymbol{\ell}'}{\chi'}\right) \rangle e^{ik_{\parallel}\chi} e^{ik'_{\parallel}\chi'} \\ &= \int_0^{\chi_s} d\chi \frac{W(\chi)}{\chi^2} \int_0^{\chi_s} d\chi' \frac{W(\chi')}{\chi'^2} \int \frac{dk_{\parallel}}{2\pi} e^{ik_{\parallel}(\chi-\chi')} \\ &\quad \times (2\pi)^2 \delta^{(2)}\left(\frac{\boldsymbol{\ell}}{\chi} + \frac{\boldsymbol{\ell}'}{\chi'}\right) P_m\left(\sqrt{k_{\parallel}^2 + \frac{\ell^2}{\chi^2}}\right) \end{aligned}$$

with $P_m(k)$ being the matter power spectrum.

To proceed, we invoke the Limber approximation (Limber, 1954). This approximation assumes that the dominant contributions arise from modes close to the line of sight, implying $k_{\parallel} \ll \ell/\chi$. Physically, this corresponds to the limit of small angular separations (or large multipole number ℓ) where the geometry of the problem simplifies. In this regime, functions that vary slowly along the line of sight can be factored out of integrals, and the highly oscillatory spherical Bessel functions that would appear in a full-sky calculation can be approximated by their limiting behavior.

Under the Limber approximation, the integral simplifies as:

$$\int \frac{dk_{\parallel}}{2\pi} e^{ik_{\parallel}(\chi-\chi')} P_m\left(\sqrt{k_{\parallel}^2 + \frac{\ell^2}{\chi^2}}\right) \approx P_m\left(\frac{\ell}{\chi}\right) \delta(\chi - \chi'), \quad (4.9)$$

where $P_m(k)$ is evaluated at $k = \ell/\chi$. Substituting this into Eq. (4.9), we obtain:

$$\langle \tilde{\kappa}(\ell) \tilde{\kappa}(\ell') \rangle = (2\pi)^2 \delta^{(2)}(\ell + \ell') \int_0^{\chi_s} d\chi \frac{W^2(\chi)}{\chi^2} P_m\left(\frac{\ell}{\chi}; \chi\right), \quad (4.10)$$

where $P_m\left(\frac{\ell}{\chi}; \chi\right)$ denotes the matter power spectrum evaluated at wavenumber $k = \ell/\chi$ and at the comoving distance χ . Finally, equating this result with the definition of the convergence power spectrum in Eq. (4.5), we derive the expression for $C_\ell^{\kappa\kappa}$:

$$C_\ell^{\kappa\kappa} = \int_0^{\chi_s} d\chi \frac{W^2(\chi)}{\chi^2} P_m\left(\frac{\ell}{\chi}; \chi\right). \quad (4.11)$$

This relation demonstrates how the observable convergence angular power spectrum $C_\ell^{\kappa\kappa}$ is sourced by the underlying matter power spectrum $P_m(k; \chi)$ integrated along the line of sight.

4.2.2 Halofit Model

Based on the halo models (discussed in Sec. 2.7), Halofit model is a widely used prescription to compute the non-linear matter power spectrum $P(k)$ from the linear power spectrum $P_L(k)$ (Smith et al., 2003; Takahashi et al., 2012). In the Halofit regime, the power spectrum is consists of two terms:

$$P(k) = P_{1h}(k) + P_{2h}(k), \quad (4.12)$$

where the two-halo term $P_{2h}(k)$ captures the contribution from large-scale linear structures, given by:

$$P_{2h}(k) = P_L(k) \left[\frac{1}{\bar{\rho}} \int dM b(M) n(M) \tilde{\rho}(k, M) \right]^2, \quad (4.13)$$

and one-halo term $P_{1h}(k)$ accounts for the contribution from small-scale non-linear structures, defined as:

$$P_{1h}(k) = \frac{1}{\bar{\rho}^2 (2\pi)^3} \int dM n(M) |\tilde{\rho}(k, M)|^2. \quad (4.14)$$

Here, $\bar{\rho}$ is the mean matter density, $n(M)dM$ is the halo mass function, $b(M)$ is the halo bias, and $\tilde{\rho}(k, M)$ is the Fourier transform of the halo density profile. Those two terms are then approximated into empirical fitting formulae and calibrated against N -body simulations.

The one-halo term resembles a shot noise spectrum on large scales but is progressively reduced on small scales due to the filtering effects of halo profiles and the mass function. Conversely, the two-halo term modifies the relative correlations of halos beyond what is predicted by linear theory and becomes negligible on small scales.

4.3 Convergence Bispectrum

The bispectrum, $B(k)$, serves as the Fourier counterpart to the three-point correlation function and is the lowest-order statistical quantity capable of characterizing non-Gaussianity in the matter distribution (Bernardeau et al., 2002). While the power spectrum effectively captures Gaussian fluctuations through two-point statistics, the bispectrum provides deeper insights by incorporating three-point correlations, thereby unveiling more complex structures in the cosmic density field (Scoccimarro et al., 1999; Takada & Jain, 2004). In a purely Gaussian scenario, the bispectrum vanishes, whereas any non-zero measurement directly signals skewness and reveals the presence of non-Gaussian features in the matter distribution.

Analogous to the angular power spectrum, the convergence bispectrum can be expressed as the ensemble average of three Fourier-transformed convergence modes, $\tilde{\kappa}$ (Dodelson & Zhang, 2005):

$$\langle \tilde{\kappa}(\ell_1) \tilde{\kappa}(\ell_2) \tilde{\kappa}(\ell_3) \rangle = (2\pi)^2 \delta_D(\ell_1 + \ell_2 + \ell_3) B_{\ell_1 \ell_2 \ell_3}^\kappa, \quad (4.15)$$

Building upon the derivations analogous to Equations (4.8) through (4.10), the convergence bispectrum can be expressed as:

$$B_{\ell_1 \ell_2 \ell_3}^\kappa = \int_0^{\chi_s} d\chi \frac{W^3(\chi)}{\chi^4} B_m \left(\frac{\ell_1}{\chi}, \frac{\ell_2}{\chi}, \frac{\ell_3}{\chi}; \chi \right), \quad (4.16)$$

where $B_m(k_1, k_2, k_3, z)$ denotes the matter bispectrum at redshift z , and $W(\chi)$ is the lensing kernel.

The bispectrum depends not only on the magnitudes of the wavevectors but also on the shapes formed by the triplet $(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$, constrained by the condition $\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 = 0$. Different triangle configurations (e.g. equilateral, squeezed, isosceles) probe different physical processes and scales in the Universe (Dodelson & Zhang, 2005). For instance, equilateral triangles are sensitive to primordial non-Gaussianity, while squeezed triangles probe the growth of structure and the impact of gravity on large scales.

4.4 Probability Density Functions

The Probability Density Function (PDF) of the convergence field, κ , provides a fundamental statistical characterization of the field's one-point distribution. By encompassing all moments and cumulants, the PDF captures both Gaussian and non-Gaussian features intrinsic to the convergence field. The PDF is defined as:

$$P(\kappa) d\kappa = \text{Prob}(\kappa \leq \kappa' \leq \kappa + d\kappa), \quad (4.17)$$

where Prob denotes the probability that the convergence κ' lies within the interval $[\kappa, \kappa + d\kappa]$.

4.4.1 Smoothing and Normalization Procedures

To effectively suppress noise and small-scale fluctuations, the convergence map $\kappa(\hat{\mathbf{n}})$ is first smoothed with a Gaussian kernel. The smoothed convergence field, $\kappa_{\text{smooth}}(\hat{\mathbf{n}})$, is defined as:

$$\kappa_{\text{smooth}}(\hat{\mathbf{n}}) = \int_{S^2} \kappa(\hat{\mathbf{n}}') W(\hat{\mathbf{n}} - \hat{\mathbf{n}}') d\hat{\mathbf{n}}', \quad (4.18)$$

where $W(\theta)$ is the Gaussian smoothing kernel given by:

$$W(\theta) = \frac{1}{2\pi\sigma_\theta^2} \exp\left(-\frac{\theta^2}{2\sigma_\theta^2}\right), \quad (4.19)$$

with $\theta = \arccos(\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}')$ representing the angular separation between the points $\hat{\mathbf{n}}$ and $\hat{\mathbf{n}}'$ with $\hat{\mathbf{n}}$, the unitary normal vector over the unit sphere S^2 , and σ_θ is the smoothing scale. To standardize the statistical analysis, the smoothed convergence values are normalized by their standard deviation. The normalized smoothed convergence, $\tilde{\kappa}_{\text{smooth},i}$, is defined as:

$$\nu_i = \frac{\kappa_{\text{smooth},i} - \langle \kappa_{\text{smooth}} \rangle}{\sigma_{\text{smooth}}}, \quad (4.20)$$

where:

$$\langle \kappa_{\text{smooth}} \rangle = \frac{1}{N_{\text{pix}}} \sum_{i=1}^{N_{\text{pix}}} \kappa_{\text{smooth},i}, \quad \sigma_{\text{smooth}}^2 = \sigma_{\text{signal}}^2 + \sigma_{\text{noise}}^2. \quad (4.21)$$

Formally, the PDF $P(\nu)$ is defined such that:

$$P(\nu) d\nu = \text{Prob}(\nu \leq \nu' \leq \nu + d\nu), \quad (4.22)$$

4.4.2 Practical Estimation of the Convergence PDF

For a discrete set of normalized convergence measurements $\{\nu_i\}_{i=1}^{N_{\text{pix}}}$ obtained from N_{pix} pixels, the PDF can be represented using the Dirac delta function δ_D :

$$P(\nu) = \frac{1}{N_{\text{pix}}} \sum_{i=1}^{N_{\text{pix}}} \delta_D(\nu - \nu_i). \quad (4.23)$$

This expression effectively constructs the PDF by summing over all pixel values, assigning a weight to each normalized convergence measurement ν_i at its exact value.

In practical applications, however, the Dirac delta function is not computationally feasible. Instead, we approximate the PDF by discretizing the normalized convergence values into bins of finite

width $\Delta\nu$. This leads to a binned estimator:

$$P(\nu) \approx \frac{1}{N_{\text{pix}}\Delta\nu} \sum_{i=1}^{N_{\text{pix}}} \Theta\left(|\nu_i - \nu| \leq \frac{\Delta\nu}{2}\right), \quad (4.24)$$

where $\Theta(x)$ is the Heaviside step function. This estimator counts the number of normalized convergence ν_i that fall within each bin centered at ν , normalizing by the total number of pixels and the bin width $\Delta\nu$.

4.4.3 hmpdf Model

The hmpdf model (Thiele et al., 2020) presents a halo-model formalism to compute the weak lensing convergence PDF, and its covariance matrix. The one-point $P(\kappa_a)$ and two-point $P(\kappa_a, \kappa_b; \phi)$ PDFs are separated into one-halo and two-halo terms as:

$$P_{1\text{pt}/2\text{pt}} = P_{1\text{pt}/2\text{pt}}^{1h} P_{1\text{pt}/2\text{pt}}^{2h}, \quad (4.25)$$

for exact formulae, see Thiele et al. (2020). Expanding the exponentials to the first order, the p -th order of the one-halo term describes overlaps of p halos along the line of sight. The two-halo term arises from the dependence of halo density on the underlying matter density field.

4.5 Peak and Minimum Counts

Local maxima (peaks) and minima in convergence field correspond to regions of over-densities and under-densities, respectively. Analyzing the statistics of these extrema offers insights into the non-Gaussian features of the matter distribution, providing a powerful tool to constrain cosmological models beyond traditional two-point statistics like the power spectrum (Jain & Van Waerbeke, 2000; Dietrich & Hartlap, 2010).

The i -th pixel in the normalized convergence map, ν is identified as a peak or a minimum by comparing its value with those of its neighboring pixels. Formally, let $\mathcal{N}(i)$ denote the set of neighboring pixels adjacent to pixel i . The conditions for a pixel to be classified as a peak or a minimum are then defined as:

$$\text{Peak Condition: } \kappa_{\text{smooth},i} > \kappa_{\text{smooth},j} \quad \forall j \in \mathcal{N}(i), \quad (4.26)$$

$$\text{Minimum Condition: } \kappa_{\text{smooth},i} < \kappa_{\text{smooth},j} \quad \forall j \in \mathcal{N}(i). \quad (4.27)$$

These conditions ensure that peaks are local maxima and minima are local minima in the convergence field. Figure 4.1 illustrates the identification of peaks (red circles) and minima (blue circles) in the smoothed convergence map $\kappa_{\text{smooth}}(\hat{\mathbf{n}})$.




Figure 4.1: Identification of peaks and minima in a smoothed convergence map. The left panel shows the smoothed convergence field $\kappa_{\text{smooth}}(\hat{\mathbf{n}})$ with peaks (red circles) satisfying the peak condition (Equation (4.26)), and the right panel highlights the minima (blue circles) satisfying the minimum condition (Equation (4.27)).

The noise properties of peaks can be analytically described using Gaussian statistics (Bardeen et al., 1986; Bond & Efstathiou, 1987). Due to the symmetry of a Gaussian field, the expected number density of minima below $-\nu$ is equal to that of peaks above ν , $N_{\text{min}}(-\nu) = N_{\text{peak}}(\nu)$.

Different approaches to analytically predict weak lensing peak counts have been developed over the recent years. One milestone is the model for Gaussian random Field developed by Bardeen et al. (1986), and the extension to the weak lensing field by Maturi et al. (2010). Another popular way to analyze weak lensing data is to compare the measured peak counts to weak lensing maps generated by N-body simulations (Matsubara, 2010; Kacprzak et al., 2016; Martinet et al., 2018; Zürcher et al., 2021).

4.6 Minkowski Functionals

Minkowski functionals provide an algebraic description of the geometrical properties of a field and are widely used to quantify the morphology of cosmic structures (Mecke et al., 1994). In cosmology, they measure the features of the patterns formed by large-scale structure of the Universe (Schmalzing et al., 1996; Schmalzing & Buchert, 1997).

4.6.1 Definition of Minkowski Functionals

For a two-dimensional normalized convergence field $\nu(\hat{\mathbf{n}})$ with zero mean and variance σ_0^2 , the excursion set $\Sigma(\nu_0)$ at threshold ν_0 is defined as the set of points where the field exceeds a certain threshold value:

$$\Sigma(\nu_0) = \left\{ \hat{\mathbf{n}} \in S^2 \mid \nu(\hat{\mathbf{n}}) > \nu_0 \right\}. \quad (4.28)$$

Figure 4.2 shows $\Sigma(\nu_0)$ for increasing threshold values $\nu_0 = 0.5, 1, 1.5$ and 2. As the threshold increases, the size and connectivity of the excursion sets diminish, reflecting the decreasing number of high-value regions in the field.




Figure 4.2: Excursion sets $\Sigma(\nu) = \{\kappa > \nu\sigma_0\}$ for increasing thresholds ($\nu = 0.5, 1, 1.5, 2$). Black regions indicate areas where κ exceeds $\nu\sigma_0$, showing reduced size and connectivity as ν increases.

The Minkowski functionals $V_0(\nu_0)$, $V_1(\nu_0)$, and $V_2(\nu_0)$ quantify the morphological properties of these excursion sets (Matsubara, 2010):

$$V_0(\nu_0) = \frac{1}{A} \int_{\Sigma(\nu_0)} da, \quad (4.29)$$

$$V_1(\nu_0) = \frac{1}{4A} \int_{\partial\Sigma(\nu_0)} dl, \quad (4.30)$$

$$V_2(\nu_0) = \frac{1}{2\pi A} \int_{\partial\Sigma(\nu_0)} \mathcal{K} dl, \quad (4.31)$$

where A is the total area, da and dl are area and length elements, and $\mathcal{K}$ is the geodesic curvature of the boundary $\partial\Sigma(\nu)$. Specifically: $V_0(\nu)$ measures the area fraction of the excursion set, $V_1(\nu)$ measures half the boundary length per unit area, and $V_2(\nu)$ quantifies the Euler characteristic per unit area.

For a pixelized map with ν_{pix} pixels, the continuous integrals in Equations (4.29)–(4.31) are approximated by discrete sums (Kratovich et al., 2012):

$$V_0(\nu_0) \approx \frac{1}{N_{\text{pix}}} \sum_{i=1}^{N_{\text{pix}}} \Theta(\nu_i - \nu_0), \quad (4.32)$$

$$V_1(\nu_0) \approx \frac{1}{4N_{\text{pix}}} \sum_{i=1}^{N_{\text{pix}}} \sum_{j \in \mathcal{N}(i)} |\Theta(\nu_i - \nu_0) - \Theta(\nu_j - \nu_0)|, \quad (4.33)$$

$$V_2(\nu_0) \approx \frac{1}{2\pi N_{\text{pix}}} \sum_{i=1}^{N_{\text{pix}}} \delta_D(\nu_i - \nu_0) \left(\frac{\nu_{,xx}\nu_{,yy} - \nu_{,xy}^2}{\nu_{,x}^2 + \nu_{,y}^2} \right), \quad (4.34)$$

where Θ is the Heaviside function, δ_D the Dirac delta function, $\mathcal{N}(i)$ the neighboring pixels of i , and derivatives are estimated via finite differences.

4.6.2 Analytical Expressions in Gaussian Random Fields

For a two-dimensional GRF $\kappa(\hat{\mathbf{n}})$ with zero mean and unit variance (after normalization), the Minkowski functionals are ([Matsubara, 2010](#)):

$$V_0(\nu) = \frac{1}{2} \operatorname{erf} \left(\frac{\nu}{\sqrt{2}} \right), \quad (4.35)$$

$$V_1(\nu) = \frac{\sigma_1}{8\sqrt{2}\sigma_0} e^{-\nu^2/2}, \quad (4.36)$$

$$V_2(\nu) = \frac{\sigma_1^2}{2\pi\sigma_0^3} \nu e^{-\nu^2/2}, \quad (4.37)$$

where erf is the error function, and $\sigma_1^2 = \langle |\nabla \nu|^2 \rangle = \langle \kappa_{,x}^2 + \kappa_{,y}^2 \rangle$. These expressions provide a Gaussian benchmark for identifying non-Gaussian features in the data.

Chapter 5

Estimation of Covariance Matrix for Weak Lensing Statistics

Contents

5.1	The Role of the Covariance Matrix in Parameter Inference	51
5.2	Covariance of the Matter Power Spectrum	52
5.3	Covariance of the Angular Convergence Power Spectrum	54
5.4	Covariance of Higher-Order Weak Lensing Statistics	55

The covariance matrix encapsulates the uncertainties and correlations between different measurements. It plays a critical role in parameter estimation techniques, including maximum likelihood analyses and Bayesian inference, and is foundational in forecasting the capabilities of future surveys through the Fisher information matrix.

The covariance matrix between two observables $\mathcal{O}_i$ and $\mathcal{O}_j$ is defined as:

$$\text{Cov}(\mathcal{O}_i, \mathcal{O}_j) = \langle (\mathcal{O}_i - \langle \mathcal{O}_i \rangle)(\mathcal{O}_j - \langle \mathcal{O}_j \rangle) \rangle, \quad (5.1)$$

where $\langle \cdot \rangle$ denotes the ensemble average over multiple realizations. For an unbiased estimator, the covariance matrix is given by:

$$\text{Cov}(\mathcal{O}_i, \mathcal{O}_j) = \frac{1}{N_{\text{sim}} - 1} \sum_{n=1}^{N_{\text{sim}}} (\mathcal{O}_i^{(n)} - \langle \mathcal{O}_i \rangle)(\mathcal{O}_j^{(n)} - \langle \mathcal{O}_j \rangle), \quad (5.2)$$

where N_{sim} is the number of simulations, and $\mathcal{O}_i^{(n)}$ is the i -th realization of the statistic in the n -th simulation.

5.1 The Role of the Covariance Matrix in Parameter Inference

The study of the covariance matrix is essential for Fisher information analyses, but we will defer a comprehensive discussion to future work. Nonetheless, it is instructive to briefly discuss the importance of the covariance matrix in the context of weak lensing statistics (Takada & Jain, 2004; Kilbinger & Schneider, 2005). In particular, the covariance matrix serves as a fundamental ingredient in quantifying the uncertainties, correlations, and overall statistical properties of the data.

5.1.1 Likelihood Functions and the Gaussian Approximation

In cosmology, as in many areas of science, we want to determine the values of underlying model parameters $\mathbf{p}$ that best describe the data $\mathbf{d}$. This process is typically done within a Bayesian framework:

$$P(\mathbf{p}|\mathbf{d}) = \frac{\mathcal{L}(\mathbf{d}|\mathbf{p})\pi(\mathbf{p})}{\epsilon(\mathbf{d})}, \quad (5.3)$$

where $P(\mathbf{p}|\mathbf{d})$ is the posterior distribution, $\pi(\mathbf{p})$ is the prior distribution of the parameters, and $\epsilon(\mathbf{d})$ is the evidence, which normalizes the posterior distribution:

$$\epsilon(\mathbf{d}) = \int \mathcal{L}(\mathbf{d}|\mathbf{p})\pi(\mathbf{p}) d\mathbf{p}. \quad (5.4)$$

The likelihood function $\mathcal{L}(\mathbf{d}|\mathbf{p})$ quantifies the probability of observing the data $\mathbf{d}$ given the model parameters $\mathbf{p}$. Suppose we have a set of data points $\mathbf{d} = \{d_1, d_2, \dots, d_N\}$, the joint probability of observing the data is obtained as:

$$\mathcal{L}(\mathbf{d}|\mathbf{p}) = \prod_{i=1}^N \mathcal{L}(d_i|\mathbf{p}), \quad (5.5)$$

For Gaussian-distributed data, it can be simply combine into a multivariate Gaussian distribution, and thus the likelihood function can be expressed as:

$$\ln \mathcal{L}(\mathbf{d}|\mathbf{p}) = -\frac{1}{2}(\mathbf{d} - \mathbf{m}(\mathbf{p}))^T \mathbf{C}^{-1}(\mathbf{d} - \mathbf{m}(\mathbf{p})) + \text{const}, \quad (5.6)$$

where $\mathbf{m}(\mathbf{p})$ is the model prediction for the data vector $\mathbf{d}$. The covariance matrix $\mathbf{C}$ quantifies the uncertainties and correlations between different data points.

5.1.2 Fisher Information and Parameter Constraints

Based on the likelihood function, we can construct the Fisher information matrix $\mathcal{F}_{\alpha\beta}$, which quantifies the sensitivity of the likelihood function to changes in the model parameters $\mathbf{p}$. The Fisher information matrix is defined as:

$$\mathcal{F}_{\alpha\beta} = -\left\langle \frac{\partial^2 \ln \mathcal{L}}{\partial p_\alpha \partial p_\beta} \right\rangle, \quad (5.7)$$

In the case of Gaussian likelihoods, the Fisher information matrix is simplified to:

$$\mathcal{F}_{\alpha\beta} = \frac{1}{2} \text{Tr} \left[\mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial p_\alpha} \mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial p_\beta} \right] \bigg|_{p_\alpha = \mu_\alpha} + \left(\frac{\partial \mu}{\partial p_\alpha} \right)^T \mathbf{C}^{-1} \frac{\partial \mu}{\partial p_\beta}, \quad (5.8)$$

where μ is the mean of the data vector $\mathbf{d}$. As assumed in most studies, the covariance matrix $\mathbf{C}$ is model independent, so that the Fisher information matrix can be reduced to:

$$\mathcal{F}_{\alpha\beta} = \left(\frac{\partial \mu}{\partial p_\alpha} \right)^T \mathbf{C}^{-1} \frac{\partial \mu}{\partial p_\beta}, \quad (5.9)$$

The Fisher matrix allows us to forecast the expected uncertainties on the parameters via the Cramér-Rao bound (Rao, 1952):

$$\langle (\Delta p_\alpha)^2 \rangle \geq (\mathcal{F}^{-1})_{\alpha\alpha}, \quad (5.10)$$

where Δp_α is the uncertainty on the α -th parameter. Note that this condition is marginalized over all other parameters, p_β ($\beta \neq \alpha$). The correlation coefficient between two parameters p_α and p_β is given by:

$$\rho_{\alpha\beta} = \frac{(\mathcal{F}^{-1})_{\alpha\beta}}{\sqrt{(\mathcal{F}^{-1})_{\alpha\alpha}(\mathcal{F}^{-1})_{\beta\beta}}}. \quad (5.11)$$

Therefore, the prior information $\pi(\mathbf{p})$ together with the likelihood function $\mathcal{L}(\mathbf{d}|\mathbf{p})$ can be used to infer the posterior distribution $P(\mathbf{p}|\mathbf{d})$.

5.2 Covariance of the Matter Power Spectrum

Understanding the covariance matrix of the matter power spectrum $P_m(k)$ is crucial before delving into two-dimensional weak lensing statistics. The covariance matrix for the matter power spectrum is defined as:

$$\text{Cov}(k, k') = \langle \hat{P}_m(k) \hat{P}_m(k') \rangle - \langle \hat{P}_m(k) \rangle \langle \hat{P}_m(k') \rangle, \quad (5.12)$$

where $\hat{P}_m(k)$ is an estimator of the matter power spectrum obtained from a finite volume V . An estimator for the matter power spectrum in a finite survey volume is given by (Feldman et al., 1994):

$$\hat{P}_m(k) = (2\pi)^3 \delta_D(\mathbf{k} + \mathbf{k}') \tilde{\delta}(\mathbf{k}) \tilde{\delta}(-\mathbf{k}) = V_f \int_{V_s(k)} \frac{d^3\mathbf{k}}{V_s(k)} \tilde{\delta}(\mathbf{k}) \tilde{\delta}(-\mathbf{k}), \quad (5.13)$$

where $V_f = (2\pi)^3/V$ is the volume of a Fourier cell where V is the total survey volume, and $V_s(k) = 4\pi k^2 \Delta k$ is the volume of the shell in Fourier space corresponding to wavenumber k .

To derive the covariance matrix, we substitute the estimator $\hat{P}_m(k)$ into the covariance definition:

$$\text{Cov}(\mathbf{k}, \mathbf{k}') = V_f^2 \left(\langle \tilde{\delta}(\mathbf{k}) \tilde{\delta}(-\mathbf{k}) \tilde{\delta}(\mathbf{k}') \tilde{\delta}(-\mathbf{k}') \rangle - \langle \tilde{\delta}(\mathbf{k}) \tilde{\delta}(-\mathbf{k}) \rangle \langle \tilde{\delta}(\mathbf{k}') \tilde{\delta}(-\mathbf{k}') \rangle \right) \quad (5.14)$$

The four-point correlation function $\langle \tilde{\delta}(\mathbf{k}_1)\tilde{\delta}(\mathbf{k}_2)\tilde{\delta}(\mathbf{k}_3)\tilde{\delta}(\mathbf{k}_4) \rangle$ can be decomposed using Wick's theorem into products of two-point functions (Wick, 1950):

$$\begin{aligned} \langle \tilde{\delta}(\mathbf{k})\tilde{\delta}(-\mathbf{k})\tilde{\delta}(\mathbf{k}')\tilde{\delta}(-\mathbf{k}') \rangle &= \langle \tilde{\delta}(\mathbf{k})\tilde{\delta}(-\mathbf{k}) \rangle \langle \tilde{\delta}(\mathbf{k}')\tilde{\delta}(-\mathbf{k}') \rangle \\ &+ \langle \tilde{\delta}(\mathbf{k})\tilde{\delta}(\mathbf{k}') \rangle \langle \tilde{\delta}(-\mathbf{k})\tilde{\delta}(-\mathbf{k}') \rangle \\ &+ \langle \tilde{\delta}(\mathbf{k})\tilde{\delta}(-\mathbf{k}') \rangle \langle \tilde{\delta}(-\mathbf{k})\tilde{\delta}(\mathbf{k}') \rangle \\ &+ \langle \tilde{\delta}(\mathbf{k})\tilde{\delta}(-\mathbf{k})\tilde{\delta}(\mathbf{k}')\tilde{\delta}(-\mathbf{k}') \rangle_c, \end{aligned} \quad (5.15)$$

where the last term represents the connected (non-Gaussian) part of the four-point function, known as the trispectrum $T(\mathbf{k}, -\mathbf{k}, \mathbf{k}', -\mathbf{k}')$.

Using the properties of the Dirac delta function and assuming statistical isotropy, the covariance matrix simplifies to:

$$\text{Cov}(k, k') = \frac{2P_m(k)^2}{N(k)}\delta_{k,k'} + \frac{T(k, k')}{V} \quad (5.16)$$

where $N(k) = V_s(k)/V_f$ is the number of independent modes in the shell at wavenumber k . The first term represents the Gaussian (disconnected) contribution, and the second term accounts for the non-Gaussian (connected) contribution from the trispectrum. The first term indicates the unavoidable cosmic variance because we are only able to measure from only one realization of the Universe.

5.2.1 Shot Noise and Super-Sample Covariance

In practice, we can only observe from discrete tracers, such as galaxies, which introduce shot noise into the power spectrum. The shot noise term can be included in the matter power spectrum as:

$$P_{\text{obs}}(k) = P_m(k) + \frac{1}{\bar{n}}, \quad (5.17)$$

where $\bar{n}$ is the number density of galaxies.

In the presence of a finite survey volume, super-sample covariance arises due to modes larger than the survey size influencing the observed modes (Takada & Hu, 2013). This effect adds an additional term to the covariance matrix:

$$\text{Cov}(k, k') = \frac{2P_m^2(k)}{N(k)}\delta_{kk'} + \frac{1}{V}T(k, k') + \left(\frac{\partial P_m(k)}{\partial \delta_b} \right) \left(\frac{\partial P_m(k')}{\partial \delta_b} \right) \sigma_b^2, \quad (5.18)$$

where δ_b represents the large-scale (background) density fluctuation, and σ_b^2 is its variance:

$$\sigma_b^2 = \int \frac{d^3k}{(2\pi)^3} P_m(k) |\tilde{W}(\mathbf{k})|^2, \quad (5.19)$$

with $\tilde{W}(\mathbf{k})$ being the Fourier transform of the survey window function $W(\mathbf{x})$. The derivatives $\partial P_m(k)/\partial \delta_b$

quantify the response of the power spectrum to changes in the background density and can be related to the concept of the response function or integrated perturbation theory (Li et al., 2014).

5.3 Covariance of the Angular Convergence Power Spectrum

We consider a cosmological survey characterized by a window function $\mathcal{W}(\boldsymbol{\theta})$ and a total survey area $\Omega_{\mathcal{W}}$, defined as the integral of the window function over the sky (Takada & Hu, 2013):

$$\Omega_{\mathcal{W}} = \int d^2\theta \mathcal{W}(\boldsymbol{\theta}). \quad (5.20)$$

The window function $\mathcal{W}(\boldsymbol{\theta})$, and its Fourier transform, $\tilde{\mathcal{W}}(\boldsymbol{\ell})$, accounts for the survey geometry and selection effects. Therefore, we can defined the observed convergence field $\kappa_{\mathcal{W}}(\boldsymbol{\theta})$ as:

$$\kappa_{\mathcal{W}}(\boldsymbol{\theta}) = \mathcal{W}(\boldsymbol{\theta}) \kappa(\boldsymbol{\theta}). \quad (5.21)$$

where $\kappa(\boldsymbol{\theta})$ is the true convergence field. The Fourier transform of the observed convergence field is given by:

$$\tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}) = \int \frac{d^2\ell'}{(2\pi)^2} \tilde{\mathcal{W}}(\boldsymbol{\ell}') \tilde{\kappa}(\boldsymbol{\ell} - \boldsymbol{\ell}'). \quad (5.22)$$

In the presence of the window function, the estimator for the angular power spectrum C_{ℓ} is given by (Takada & Hu, 2013):

$$\hat{C}_{\ell} = \frac{1}{\Omega_{\mathcal{W}}} \tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}) \tilde{\kappa}_{\mathcal{W}}(-\boldsymbol{\ell}). \quad (5.23)$$

The covariance matrix of the angular power spectrum C_{ℓ} is defined as:

$$\text{Cov}(\ell_1, \ell_2) = \langle \hat{C}_{\ell_1} \hat{C}_{\ell_2} \rangle - \langle \hat{C}_{\ell_1} \rangle \langle \hat{C}_{\ell_2} \rangle, \quad (5.24)$$

which measures the statistical correlation between estimates of C_{ℓ_1} and C_{ℓ_2} .

Substituting the estimator $\hat{C}_{\ell}$ into the covariance definition and expanding the resulting expression leads to terms involving two-point and four-point correlation functions of the convergence field $\kappa(\boldsymbol{\theta})$. Specifically, the covariance can be expressed as:

$$\begin{aligned} \text{Cov}(\ell_1, \ell_2) &= \frac{1}{\Omega_{\mathcal{W}}^2} [\langle \tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}_1) \tilde{\kappa}_{\mathcal{W}}(-\boldsymbol{\ell}_1) \tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}_2) \tilde{\kappa}_{\mathcal{W}}(-\boldsymbol{\ell}_2) \rangle - \langle \tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}_1) \tilde{\kappa}_{\mathcal{W}}(-\boldsymbol{\ell}_1) \rangle \langle \tilde{\kappa}_{\mathcal{W}}(\boldsymbol{\ell}_2) \tilde{\kappa}_{\mathcal{W}}(-\boldsymbol{\ell}_2) \rangle] \\ &= \frac{1}{\Omega_{\mathcal{W}}^2} [\langle \tilde{\kappa}(\boldsymbol{\ell}_1) \tilde{\kappa}(-\boldsymbol{\ell}_1) \tilde{\kappa}(\boldsymbol{\ell}_2) \tilde{\kappa}(-\boldsymbol{\ell}_2) \rangle + (\ell_2 \leftrightarrow -\ell_2)] \\ &+ \frac{1}{\Omega_{\mathcal{W}}^2} \langle \tilde{\kappa}(\boldsymbol{\ell}_1) \tilde{\kappa}(-\boldsymbol{\ell}_1) \tilde{\kappa}(\boldsymbol{\ell}_2) \tilde{\kappa}(-\boldsymbol{\ell}_2) \rangle_c \end{aligned} \quad (5.25)$$

By a straightforward calculation, this expression can be simplified to:

$$\begin{aligned} \text{Cov}(\ell_1, \ell_2) &= \frac{1}{\Omega_{\mathcal{W}}^2} [C_{\ell_1}]^2 \left[\left| \tilde{\mathcal{W}}(\ell_1 + \ell_2) \right|^2 + \left| \tilde{\mathcal{W}}(\ell_1 - \ell_2) \right|^2 \right] \\ &+ \frac{1}{\Omega_{\mathcal{W}}^2} \int \frac{d^2 \ell'}{(2\pi)^2} \mathcal{T}(\ell'_1, -\ell'_1 + \ell', \ell'_2, -\ell'_2 - \ell'), \end{aligned} \quad (5.26)$$

where $\mathcal{T}(\ell_1, -\ell_1, \ell_2, -\ell_2)$ is the trispectrum of the convergence field.

Using the Limber approximation (Limber, 1954), which simplifies the projection of three-dimensional quantities into two dimensions, the covariance matrix can be related to the matter power spectrum.

$$C_\ell = \int d\chi \frac{W^2(\chi)}{\chi^2} P_m\left(\frac{\ell}{\chi}, \chi\right), \quad (5.27)$$

$$\mathcal{T}(\ell_1, -\ell_1, \ell_2, -\ell_2) = \int d\chi \frac{W^4(\chi)}{\chi^6} T_m\left(\frac{\ell_1}{\chi}, -\frac{\ell_1}{\chi}, \frac{\ell_2}{\chi}, -\frac{\ell_2}{\chi}\right), \quad (5.28)$$

Similar to the matter power spectrum, the covariance matrix for the angular power spectrum can be decomposed into:

$$\text{Cov}(\ell_1, \ell_2) = \text{Cov}^G(\ell_1, \ell_2) + \text{Cov}^{\text{cNG}}(\ell_1, \ell_2) + \text{Cov}^{\text{SSC}}(\ell_1, \ell_2), \quad (5.29)$$

where:

$$\text{Cov}^G(\ell_1, \ell_2) = \frac{1}{\Omega_{\mathcal{W}}^2} [C_{\ell_1}]^2 \left[\left| \tilde{\mathcal{W}}(\ell_1 + \ell_2) \right|^2 + \left| \tilde{\mathcal{W}}(\ell_1 - \ell_2) \right|^2 \right], \quad (5.30)$$

$$\text{Cov}^{\text{cNG}}(\ell_1, \ell_2) = \frac{1}{\Omega_{\mathcal{W}}^2} \int \frac{d^2 \ell'}{(2\pi)^2} \mathcal{T}^{\text{cNG}}(\ell'_1, -\ell'_1 + \ell', \ell'_2, -\ell'_2 - \ell'), \quad (5.31)$$

$$\text{Cov}^{\text{SSC}}(\ell_1, \ell_2) = \frac{1}{\Omega_{\mathcal{W}}^2} \int \frac{d^2 \ell'}{(2\pi)^2} \left| \tilde{\mathcal{W}}(\ell') \right|^2 \sigma_{\ell_1, \ell_2}^{\ell'}, \quad (5.32)$$

with:

$$\sigma_{\ell_1, \ell_2}^{\ell'} = \int d\chi \frac{W^4(\chi)}{\chi^6} \left(\frac{\partial P_m\left(\frac{\ell_1}{\chi}, \chi\right)}{\partial \delta_b} \right) \left(\frac{\partial P_m\left(\frac{\ell_2}{\chi}, \chi\right)}{\partial \delta_b} \right) P_m\left(\frac{\ell_1}{\chi}, \chi\right) P_m\left(\frac{\ell_2}{\chi}, \chi\right) P_L\left(\frac{\ell'}{\chi}\right), \quad (5.33)$$

where $P_L(k)$ is the linear matter power spectrum. Notably, the SSC term $\text{Cov}^{\text{SSC}}(\ell_1, \ell_2)$ arises from the large-scale density fluctuations modulating the observed power spectrum within the survey area (Takada & Hu, 2013).

5.4 Covariance of Higher-Order Weak Lensing Statistics

Despite some successes in analytical modeling (Chan et al. 2018; Lacasa et al. 2018; Lacasa & Grain 2019; Uhlemann et al. 2023), computing the covariance matrices for higher-order statistics still need to rely on simulations. Drawing an analogy with the matter power spectrum, the covariance matrix

for higher-order statistics can be expressed as:

$$\text{Cov}(\mathcal{O}_i, \mathcal{O}_j) = \text{Cov}^{\text{noSSC}}(\mathcal{O}_i, \mathcal{O}_j) + \text{Cov}^{\text{SSC}}(\mathcal{O}_i, \mathcal{O}_j), \quad (5.34)$$

A rigorous super-sample covariance for line-of-sight integrated observable $\mathcal{O}_i$, where $\mathcal{O}_i = \int dV_i \mathbf{o}_i = \int \chi_i^2 d\chi \mathbf{o}_i$, is given by (Lacasa & Rosenfeld, 2016):

$$\text{Cov}^{\text{SSC}}(\mathcal{O}_i, \mathcal{O}_j) = \iint dV_i dV_j \left(\frac{\partial \mathbf{o}_i}{\partial \delta_b} \right) \left(\frac{\partial \mathbf{o}_j}{\partial \delta_b} \right) \sigma_b^2, \quad (5.35)$$

In most cases, the accurate estimation of such covariance matrices still necessitates averaging over numerous realizations of N-body simulations. These simulations capture the relevant mode-coupling and environmental dependencies that are difficult to model analytically, ensuring that the resulting covariance matrices are both realistic and robust. This approach is exactly what we will adopt in this study to quantify the uncertainties due to super-sample covariance in higher-order statistics.

Numerical Simulations for Large Scale Structure and Weak Lensing

Contents

6.1	Introduction to Numerical Simulations in Cosmology	57
6.2	Initial Condition Generation	58
6.3	N-Body Simulation Methods	60
6.4	Computational Tools and Optimization Techniques	63
6.5	Advanced Codes and Methods: FASTPM	66
6.6	Generating Weak Lensing Maps from Simulations	68

6.1 Introduction to Numerical Simulations in Cosmology

Numerical simulations are crucial in physics and astronomy for studying complex systems of interacting particles, such as galaxies and the Universe’s large-scale structure, where analytical solutions are often impractical due to complexity and nonlinearity ([Hockney & Eastwood, 1981](#); [Efstathiou et al., 1985](#); [Springel et al., 2005](#); [Springel, 2005](#)). This section provides an overview of N -body simulations commonly used in cosmology.

6.1.1 Dark Matter-Only vs. Hydrodynamical Simulations

In cosmology and astrophysics, simulation can be grouped into two main categories: dark matter-only simulations and hydrodynamical simulations.

Dark matter simulations are used to study the large-scale structure of the universe, the formation of dark matter halos, and the evolution of dark matter particles under the influence of gravity. Since dark matter is effectively collisionless and interacts primarily through gravity, it can be modeled as a collection of particles that evolve under the influence of gravitational forces ([Efstathiou et al., 1985](#)).

Hydrodynamical simulations, on the other hand, include the effects of gas dynamics, star formation, and feedback processes from supernovae and active galactic nuclei (AGN) (Hernquist & Katz, 1989; Springel et al., 2005; Springel, 2005). These simulations are more computationally expensive and require additional physics beyond gravity, such as hydrodynamics, radiative transfer, and chemistry.

6.1.2 Historical Evolution and Growth in Particle Number

Since the 1980s, numerical cosmology has developed algorithms to mitigate the computational challenges posed by long-range gravitational interactions by reducing global communication across the computational domain. Key algorithms include mesh-based methods, tree codes, and multipole expansions (Hockney & Eastwood, 1981). Figure 6.1 displays the number of particles used in selected N -body simulations employing these techniques. Symbols and colors indicate the gravitational solvers: particle-particle-particle-mesh (P^3M) and adaptive P^3M (AP^3M); parallel or vectorized P^3M ; Tree codes; TreePM; and particle-mesh methods with adaptive mesh refinement (PM AMR).

Advancements in algorithms and software optimization have increased the number of particles in cosmological simulations beyond what direct summation methods allow. Since 1990, gravity-only simulations have exhibited a super-exponential growth trend, indicated by the quadratic regression in Figure 6.1, reflecting significant methodological innovations beyond hardware improvements (Leclercq, 2020).

6.2 Initial Condition Generation

As we have seen in Section 2.4, the primordial power spectrum $P(k)$ is a key ingredient in generating initial conditions for cosmological simulations. Based on the linear power spectrum, we will review the process of generating initial conditions for N -body simulations.

6.2.1 Initial Density Field

To generate the initial density field for the simulations, we express the density contrast $\delta(\mathbf{x})$ in terms of its Fourier components:

$$\delta(\mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \tilde{\delta}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}}. \quad (6.1)$$




Figure 6.1: Evolution of the number of particles used in N -body simulations as a function of the year of publication (Leclercq, 2020). The symbols and colors indicate the gravitational solver employed: P³M and adaptive P³M (AP³M); parallel or vectorized P³M; Tree codes; TreePM; and particle-mesh methods with adaptive mesh refinement (PM AMR). Hydrodynamic simulations are represented by black squares.

Assuming a Gaussian random field, each Fourier mode $\tilde{\delta}(\mathbf{k})$ is a complex Gaussian random variable with zero mean and variance $P(k)$:

$$\tilde{\delta}(\mathbf{k}) = A(\mathbf{k}) + iB(\mathbf{k}), \quad (6.2)$$

$$\langle A(\mathbf{k}) \rangle = \langle B(\mathbf{k}) \rangle = 0, \quad (6.3)$$

$$\langle A(\mathbf{k})A(\mathbf{k}') \rangle = \langle B(\mathbf{k})B(\mathbf{k}') \rangle = \frac{1}{2}P(k)\delta_D(\mathbf{k} - \mathbf{k}'), \quad (6.4)$$

$$\langle A(\mathbf{k})B(\mathbf{k}') \rangle = 0, \quad (6.5)$$

where $A(\mathbf{k})$ and $B(\mathbf{k})$ are real Gaussian random variables, and δ_D is the Dirac delta function.

6.2.2 Initial Displacement Field

The initial displacement field $\Psi(\mathbf{q})$ relates the Lagrangian coordinates $\mathbf{q}$ to the Eulerian coordinates $\mathbf{x}$:

$$\mathbf{x}(\mathbf{q}) = \mathbf{q} + \Psi(\mathbf{q}). \quad (6.6)$$

The displacement field is proportional to the gradient of the gravitational potential $\Phi(\mathbf{q})$:

$$\Psi(\mathbf{q}) = -\nabla\Phi(\mathbf{q}), \quad (6.7)$$

where the potential satisfies Poisson's equation:

$$\nabla^2 \Phi(\mathbf{q}) = \delta(\mathbf{q}). \quad (6.8)$$

The first order solution to the displacement field is given by the Zel'dovich approximation (Zel'dovich, 1970):

$$-k^2 \tilde{\Phi}(\mathbf{k}) = \tilde{\delta}(\mathbf{k}), \quad (6.9)$$

$$\tilde{\Psi}(\mathbf{k}) = i\mathbf{k} \tilde{\Phi}(\mathbf{k}) = i\mathbf{k} \frac{\tilde{\delta}(\mathbf{k})}{k^2}, \quad (6.10)$$

$$\Psi(\mathbf{q}) = \int \frac{d^3k}{(2\pi)^3} i\mathbf{k} \frac{\tilde{\delta}(\mathbf{k})}{k^2} e^{i\mathbf{k} \cdot \mathbf{q}}. \quad (6.11)$$

6.2.3 Initial Velocities

The initial velocities of particles are derived from the time derivative of the displacement field. The velocities are given by (Efstathiou et al., 1985):

$$\mathbf{v}(\mathbf{q}) = aHf(a)\Psi(\mathbf{q}), \quad (6.12)$$

$$\tilde{\mathbf{v}}(\mathbf{k}) = aHf(a)\tilde{\Psi}(\mathbf{k}) = aHf(a)i\mathbf{k} \frac{\tilde{\delta}(\mathbf{k})}{k^2}, \quad (6.13)$$

$$\mathbf{v}(\mathbf{q}) = iaHf(a) \int \frac{d^3k}{(2\pi)^3} \frac{\mathbf{k}}{k^2} \tilde{\delta}(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{q}}, \quad (6.14)$$

where a is the scale factor, H is the Hubble parameter, and $f(a)$ is the growth rate defined as:

$$f(a) = \frac{d \ln D}{d \ln a}, \quad (6.15)$$

6.3 N-Body Simulation Methods: An Overview

We outline the fundamental concepts and algorithms used in N -body simulations, including direct summation, particle-mesh methods, particle-particle particle-mesh (P3M) methods, and tree-particle-mesh (Tree-PM) methods.

Table 6.1 summarizes the key features of these methods, including computational complexity and example objectives they are best suited for.

6.3.1 Direct Summation

Direct Summation calculates gravitational forces between all particle pairs directly, scaling as $\mathcal{O}(N^2)$ and becoming computationally intensive for large N . Therefore, direct summation is typically used for small systems like globular clusters, where accuracy is paramount (Wang et al., 2015; Panamarev

Method	Complexity	Objective	Key Features
Direct Summation	$\mathcal{O}(N^2)$	Globular Clusters	Accurate, computationally intensive
PM	$\mathcal{O}(N + M \log M)$	Large-Scale Structure	Fast, smooths small-scale forces
P3M	$\mathcal{O}(N \log N)$	Large-Scale Structure	Combines direct summation and PM
Tree-PM	$\mathcal{O}(N \log N)$	Large-Scale Structure	Tree algorithm for short-range forces

 Table 6.1: Comparison of N -body simulation methods

et al., 2019). Each particle i has position $\mathbf{r}_i$, velocity $\mathbf{v}_i$, and mass m_i . At each time step t :

1. **Compute Forces:**

$$\mathbf{F}_i = Gm_i \sum_{\substack{j=1 \\ j \neq i}}^N \frac{m_j(\mathbf{r}_j - \mathbf{r}_i)}{\|\mathbf{r}_j - \mathbf{r}_i\|^3}$$

2. **Update Particle States:**

$$\mathbf{v}_i(t + \Delta t) = \mathbf{v}_i(t) + \frac{\mathbf{F}_i}{m_i} \Delta t, \quad \mathbf{r}_i(t + \Delta t) = \mathbf{r}_i(t) + \mathbf{v}_i(t + \Delta t) \Delta t$$

3. **Advance Time:**

$$t \leftarrow t + \Delta t$$

6.3.2 Particle-Mesh (PM) Method

The PM method approximates gravitational forces by mapping particles onto a grid and solving for the gravitational potential, reducing computational cost to $\mathcal{O}(N + M \log M)$, where M is the number of grid points. Combining with Adaptive Mesh Refinement (AMR), PM simulations can achieve high resolution in regions of interest while maintaining efficiency so that it is used in Hydrodynamical simulations recently (Springel et al., 2018; Trebitsch et al., 2021). The main difference between the PM method and direct summation is the grid-based force calculation:

1. **Assign Particles to Grid:** See Section 6.4.2.

2. **Compute Density Field:**

$$\rho(\mathbf{x}) = \sum_i m_i W(\mathbf{x} - \mathbf{r}_i) \quad (\text{where } W: \text{Interpolation Kernel})$$

3. **Solve Poisson's Equation:**

$$\nabla^2 \Phi = 4\pi G \rho$$

4. **Compute Force due to Potential:**

$$\mathbf{E} = -\nabla \Phi$$

6.3.3 Particle-Particle Particle-Mesh (P3M) Method

The P³M method combines direct summation for short-range forces with the PM approach for long-range interactions, achieving $\mathcal{O}(N \log N)$ complexity while enhancing accuracy for nearby particles. Due to its balance between accuracy and efficiency, the P³M (or the Adaptive P³M (Couchman, 1991)) method is widely used in hydrodynamical simulations (Couchman et al., 1995; Teyssier, 2002).

Key parameters include mesh size, softening parameter ϵ , and force resolution.

The difference between the P³M method and the PM method lies in the force calculation:

1. Long-Range Forces (PM):

$$\mathbf{F}_{\text{long},i} = m_i \mathbf{E}_{\text{long}}(\mathbf{r}_i)$$

2. Short-Range Forces (Direct Summation):

(a) **Neighbor Search:** Identify particles j within cutoff radius r_{cut} of particle i .

(b) **Force Calculation:**

$$\mathbf{F}_{\text{short},i} = -Gm_i \sum_{j \in \text{neighbors}} \frac{m_j (\mathbf{r}_i - \mathbf{r}_j)}{(\|\mathbf{r}_i - \mathbf{r}_j\|^2 + \epsilon^2)^{3/2}}$$

3. Combine Forces:

$$\mathbf{F}_i = \mathbf{F}_{\text{long},i} + \mathbf{F}_{\text{short},i}$$

6.3.4 Tree-Particle-Mesh (Tree-PM) Method

The Tree-PM method integrates the PM approach for long-range forces with a tree algorithm for short-range interactions, reducing complexity to $\mathcal{O}(N \log N)$ (Barnes & Hut, 1986). Several popular simulations such as Illustris (Vogelsberger et al., 2014) and EAGLE (Schaye et al., 2015; Crain et al., 2015; The EAGLE team, 2017) use the Tree-PM method. Proper tuning of parameters like grid size, softening length ϵ , and opening angle θ_{max} is essential.

The main updates in the Tree-PM method compared to the P³M method are in the tree construction when calculating short-range forces:

1. Tree Construction:

(a) **Build Spatial Cells:** Partition the simulation volume into spatial cells (e.g., octree) and assign particles to nodes.

(b) **Multipole Moments:** For each node j , calculate mass M_j and center of mass $\mathbf{r}_{\text{cm},j}$.

2. **Force Calculation:** For each particle i , traverse the tree to compute the short-range gravitational force:

$$\mathbf{F}_{\text{short},i} = -Gm_i \sum_{\text{nodes}} \frac{M_j (\mathbf{r}_i - \mathbf{r}_j)}{(\|\mathbf{r}_i - \mathbf{r}_j\|^2 + \epsilon^2)^{3/2}}$$

using the opening angle criterion:

$$\theta = \frac{l_j}{\|\mathbf{r}_i - \mathbf{r}_j\|} < \theta_{\max}$$

where l_j is the size of node j and $\theta_{\max}$ is the maximum allowed opening angle.

6.4 Computational Tools and Optimization Techniques

Efficient computational tools are crucial for large-scale simulations and data analysis in scientific and engineering applications. This section overviews key computational techniques and algorithms used in N -body simulations and large-scale structure studies.

6.4.1 Fast Fourier Transforms (FFT)

The Fast Fourier Transform (FFT) is a highly efficient algorithm for computing the Discrete Fourier Transform (DFT) of a sequence. Given a sequence of N complex numbers $\{x_n\}_{n=0}^{N-1}$, the DFT is defined as:

$$X_k = \sum_{n=0}^{N-1} x_n e^{-2\pi i k n / N}, \quad k = 0, 1, \dots, N-1. \quad (6.16)$$

The naive computation of the DFT requires $\mathcal{O}(N^2)$ operations. The FFT reduces this complexity to $\mathcal{O}(N \log N)$ by exploiting the symmetry and periodicity properties of the exponential kernel. The most common FFT algorithm is the Cooley-Tukey radix-2 FFT (Cooley & Tukey, 1965), which recursively decomposes the DFT into smaller DFTs of even and odd-indexed elements:

$$X_k = \sum_{n=0}^{N/2-1} x_{2n} e^{-2\pi i k (2n) / N} + \sum_{n=0}^{N/2-1} x_{2n+1} e^{-2\pi i k (2n+1) / N} \quad (6.17)$$

$$= X_k^{\text{even}} + e^{-2\pi i k / N} X_k^{\text{odd}}, \quad (6.18)$$

where X_k^{even} and X_k^{odd} are the DFTs of the even and odd subsequences, respectively.

6.4.2 Mass Assignment Schemes

Mass assignment schemes map particle masses onto a computational grid to compute density fields and gravitational forces, ensuring mass conservation and minimizing aliasing errors. Common schemes include:

- **Nearest Grid Point (NGP):** Each particle is assigned entirely to the nearest grid point.
- **Cloud-In-Cell (CIC):** Mass is linearly interpolated to the nearest $2^3 = 8$ surrounding grid points.

- **Triangular-Shaped Cloud (TSC):** Mass is distributed to the nearest $3^3 = 27$ grid points using a quadratic interpolation function.

In Fourier space, these mass assignment window functions are represented as:

$$W(\mathbf{k}) = \prod_{i=1}^3 W(k_i), \quad (6.19)$$

where

$$W(k_i) = \left[\frac{\sin(\pi k_i / (2k_N))}{\pi k_i / (2k_N)} \right]^p, \quad (6.20)$$

with k_N being the Nyquist wavenumber, k_i the i -th component of the wavevector $\mathbf{k}$, and $p = 1$ for NGP, $p = 2$ for CIC, and $p = 3$ for TSC (Hockney & Eastwood, 1981; Efstathiou et al., 1985).

Figure 6.2 illustrates the mass assignment process for a particle distribution on a 1D grid using different schemes.




Figure 6.2: Illustration of three mass assignment schemes—Nearest Grid Point (NGP), Cloud-In-Cell (CIC), and Triangular-Shaped Cloud (TSC)—used to map a particle's mass onto a 1D grid.

6.4.3 Parallelization Techniques

Parallelization accelerates computations in large-scale simulations by leveraging multiple processors or computing nodes. Key strategies include:

- **Domain Decomposition:** The computational domain is partitioned into smaller subdomains, each assigned to a separate processor (Barnes & Hut, 1986).
- **Task Parallelism:** Distributing independent tasks across multiple processors.
- **Data Parallelism:** Performing identical operations concurrently on different data elements, enabling SIMD (Single Instruction, Multiple Data) execution.




Figure 6.3: Illustration of adaptive mesh refinement (AMR) applied to a 2D image with two Gaussian kernels. The left panel shows the initial coarse grid structure over the image. The middle and right panels demonstrate progressively finer levels of mesh refinement in regions of higher intensity, where the Gaussian kernels are located. The red grid outlines indicate the adaptively refined mesh hierarchy, ensuring higher resolution where needed while maintaining computational efficiency in lower-intensity regions.

6.4.4 Adaptive Mesh Refinement (AMR)

Adaptive Mesh Refinement (AMR) dynamically adjusts grid resolution, refining the mesh where higher accuracy is needed (e.g., regions with high density gradients) and coarsening it elsewhere (Berger & Colella, 1989). This technique is typically used in hydrodynamical simulations to capture complex fluid dynamics and shock fronts accurately. Because N-body simulations with discrete particles do not require continuous field solution and have more efficient methods like Tree-PM, AMR is not commonly used in N-body simulations.

This creates a hierarchy of grids with increasing resolution and optimizes computational resources. Refinement is typically triggered when:

$$|\nabla\phi(\mathbf{x})| > \theta, \quad (6.21)$$

with θ being a predefined threshold.

Figure 6.3 demonstrates the application of Adaptive Mesh Refinement (AMR) to a two-dimensional image containing two Gaussian kernels. Initially, a uniformly coarse grid overlays the entire image (left panel). As the refinement process progresses, the mesh becomes increasingly finer in regions with higher intensity, specifically around the Gaussian kernels (middle and right panels). The red grid lines represent the hierarchy of the refined meshes, enabling higher resolution where it is most needed and optimizing computational resources by keeping a coarser grid in less significant areas.

6.4.5 Tree Construction

Tree-based data structures efficiently organize hierarchical spatial data. The Barnes-Hut algorithm employs an octree to partition space, reducing computational complexity from $\mathcal{O}(N^2)$ to $\mathcal{O}(N \log N)$

by approximating distant particle clusters as single mass points. This approximation is controlled by the opening angle θ :

$$\frac{s}{d} < \theta, \quad (6.22)$$

where s is the node size and d is the distance from the particle to the node's center of mass.

One of the popular algorithms for tree construction is the Barnes-Hut Octree (Barnes & Hut, 1986), which recursively subdivides the simulation volume into hierarchical grid cells. Figure 6.4 illustrates the Octree decomposition for a 3D volume containing four particles.




Figure 6.4: Illustration of an Octree decomposition for a 3D volume containing four particles. The left panel showcases the spatial subdivision of the volume into hierarchical grid cells, with color-coding indicating different levels of refinement. The right panel presents the corresponding Octree data structure, highlighting the hierarchical relationships between nodes. Credit by Powell et al. (2023)

Parallel tree construction involves building local trees within each subdomain and integrating them for global computations (Dubinski, 1996). Efficient parallelization enhances scalability and performance in large-scale simulations.

6.5 Advanced Codes and Methods: FASTPM

FASTPM (Fast Particle Mesh; Feng et al. 2016) is an advanced N-body simulation code tailored for efficiently modeling the evolution of dark matter and halo structures on cosmological scales. Building upon the foundational PM approach, FASTPM integrates modified kick and drift factors derived from the Zel'dovich Approximation (ZA). This enhancement allows FASTPM to achieve high accuracy in large-scale structure formation while significantly reducing computational overhead. This subsection delineates the core methodology of FASTPM, incorporating the mathematical formalism of its modified kick and drift factors.

6.5.1 Modified Kick and Drift Factors

The cornerstone of FASTPM's enhanced performance lies in its **modified kick** (K_{FASTPM}) and **drift** (D_{FASTPM}) factors. These factors are meticulously derived from the Zel'dovich Approximation (ZA), a first-order Lagrangian perturbation theory (1LPT), to rectify inaccuracies in large-scale growth inherent in standard PM solvers, especially when operating with a limited number of time steps.

First, the Zel'dovich equation of motion to the first order is defined as:

$$\begin{aligned}\mathbf{x}_{\text{ZA}}(a) &= \mathbf{q} + D(a)\mathbf{s}_1, \\ \mathbf{p}_{\text{ZA}}(a) &= a^3 E(a) g_p(a) \mathbf{s}_1, \\ \mathbf{f}_{\text{ZA}}(a) &= a^2 E(a) g_f(a) \mathbf{s}_1,\end{aligned}\tag{6.23}$$

where $E(a) = \frac{H(a)}{H(a=1)}$ is the dimensionless Hubble parameter, and $g_p(a)$ and $g_f(a)$ are auxiliary factors defined as:

$$g_p(a) = \frac{dD}{da}, G_p(a) = D(a) \tag{6.24}$$

$$g_f(a) = \frac{d(a^3 E g_p)}{da}, G_f(a) = a^3 E g_p(a) \tag{6.25}$$

The ZA equations of motion are reformulated in terms of drift and kick operators by integrating over a time step from a_0 to a_1 and eliminating the ZA displacement $\mathbf{s}_1$:

$$\begin{aligned}\Delta \mathbf{x}_{\text{ZA}} &= \mathbf{x}_{\text{ZA}}(a_1) - \mathbf{x}_{\text{ZA}}(a_0) \\ &= [D(a)]_{a_0}^{a_1} \mathbf{s}_1 \\ &= \frac{\mathbf{p}_{\text{ZA}}(a_r)}{a_r^3 E(a_r)} \left(\frac{\Delta G_p}{g_p(a_r)} \right),\end{aligned}\tag{6.26}$$

$$\begin{aligned}\Delta \mathbf{p}_{\text{ZA}} &= \mathbf{p}_{\text{ZA}}(a_1) - \mathbf{p}_{\text{ZA}}(a_0) \\ &= \frac{\mathbf{f}_{\text{ZA}}(a_r)}{a_r^2 E(a_r)} \left(\frac{\Delta G_f}{g_f(a_r)} \right),\end{aligned}\tag{6.27}$$

where $\Delta \mathbf{x}_{\text{ZA}}$ is the change in displacement over the time step, $\Delta \mathbf{p}_{\text{ZA}}$ is the change in momentum over the time step, a_r is a reference scale factor within the time step, $\Delta G_p = G_p(a_1) - G_p(a_0)$, and $\Delta G_f = G_f(a_1) - G_f(a_0)$. Therefore, the modified kick and drift factors in FASTPM are defined as:

$$\mathcal{D}_{\text{FASTPM}} = \frac{\Delta \mathbf{x}_{\text{ZA}}}{\mathbf{p}_{\text{ZA}}} = \frac{1}{a_r^3 E(a_r)} \left(\frac{\Delta G_p}{g_p(a_r)} \right) \tag{6.28}$$

$$\mathcal{K}_{\text{FASTPM}} = \frac{\Delta \mathbf{p}_{\text{ZA}}}{\mathbf{f}_{\text{ZA}}} = \frac{1}{a_r^2 E(a_r)} \left(\frac{\Delta G_f}{g_f(a_r)} \right) \tag{6.29}$$

These operators ensure the exact integration of the ZA equations of motion, thereby accurately capturing the linear growth of structures within each time step.

6.5.2 Algorithm Steps

The main steps of the FASTPM algorithm follow the standard PM method discussed in Section 6.3.2, with the addition of the modified kick and drift operators to ensure accurate linear growth.

1. **Apply Modified Operators:** Utilize the modified kick (K_{FASTPM}) and drift (D_{FASTPM}) factors to update particle velocities and positions. These factors, derived from the ZA, ensure accurate linear growth:

- (a) **Kick Step:** Update particle velocities by applying the gravitational acceleration scaled by the modified kick factor:

$$\mathbf{v}_i \left(t + \frac{\Delta t}{2} \right) = \mathbf{v}_i(t) + \mathbf{g}_i(t) \cdot K_{\text{FASTPM}} \cdot \Delta t$$

- (b) **Drift Step:** Update particle positions using the updated velocities and the modified drift factor:

$$\mathbf{r}_i(t + \Delta t) = \mathbf{r}_i(t) + \mathbf{v}_i \left(t + \frac{\Delta t}{2} \right) \cdot D_{\text{FASTPM}} \cdot \Delta t$$

- (c) **Second Kick Step:** Apply another kick to update velocities to the full time step:

$$\mathbf{v}_i(t + \Delta t) = \mathbf{v}_i \left(t + \frac{\Delta t}{2} \right) + \mathbf{g}_i(t + \Delta t) \cdot K_{\text{FASTPM}} \cdot \Delta t$$

2. **Update Particle States:** Finalize the update of particle velocities and positions after applying the modified kick and drift operators:

$$\mathbf{v}_i(t + \Delta t) = \mathbf{v}_i \left(t + \frac{\Delta t}{2} \right) + \mathbf{g}_i(t + \Delta t) \cdot K_{\text{FASTPM}} \cdot \Delta t$$

$$\mathbf{r}_i(t + \Delta t) = \mathbf{r}_i(t) + \mathbf{v}_i \left(t + \frac{\Delta t}{2} \right) \cdot D_{\text{FASTPM}} \cdot \Delta t$$

3. **Advance Time:** Increment the simulation time by the time step Δt :

$$t \leftarrow t + \Delta t$$

6.6 Generating Weak Lensing Maps from Simulations

To extract statistics from N-body simulations, it is necessary to generate detailed WL maps by simulating the propagation of a vast number of virtual light rays from source galaxies to the observer. This

process involves calculating the distortions and magnifications of the source images caused by the cumulative gravitational deflections from the intervening matter distribution along each line of sight.

6.6.1 Conventional Ray-Tracing Algorithm

Conventional ray-tracing algorithms are fundamental tools for generating WL maps from N-body simulations. These algorithms simplify the complex three-dimensional matter distribution by projecting it onto a series of two-dimensional lens planes. The numerical implementation of the ray-tracing algorithm involves several key computational steps, outlined below (Das & Bode, 2008; Teyssier et al., 2009; Shirasaki et al., 2015):

1. **Lens Plane Generation:** The first step involves constructing the surface mass density $\Delta_{\Sigma}^j(\boldsymbol{\theta})$ for each lens plane j . This is achieved by projecting the three-dimensional matter density $\rho(\chi, \chi\boldsymbol{\theta})$ within a spherical shell bounded by comoving distances χ_j and χ_{j+1} onto a two-dimensional plane:

$$\Delta_{\Sigma}^j(\boldsymbol{\theta}) = \int_{\chi_j}^{\chi_{j+1}} [\rho(\chi, \chi\boldsymbol{\theta}) - \bar{\rho}(\chi)] \chi^2 d\chi, \quad (6.30)$$

where $\bar{\rho}(\chi)$ is the mean matter density at distance χ . The effective comoving distance to the center of the j -th shell, χ^j , is calculated as (Shirasaki et al., 2015):

$$\chi^j = \frac{\int_{\chi_{\min}}^{\chi_{\max}} \chi^3 d\chi}{\int_{\chi_{\min}}^{\chi_{\max}} \chi^2 d\chi} = \frac{3}{4} \frac{\chi_{\max}^4 - \chi_{\min}^4}{\chi_{\max}^3 - \chi_{\min}^3}, \quad (6.31)$$

where $\chi_{\min}$ and $\chi_{\max}$ are the minimum and maximum comoving distances of the shell, respectively. The convergence field $\kappa_j(\boldsymbol{\theta})$ for the j -th lens plane is then computed as:

$$\kappa_j(\boldsymbol{\theta}) = \frac{4\pi G}{c^2} \frac{\Delta_{\Sigma}^j(\boldsymbol{\theta})}{a_j \chi_j}, \quad (6.32)$$

with a_j being the scale factor and χ_j the comoving distance to the j -th lens plane.

To express $\kappa_j(\boldsymbol{\theta})$ in terms of the simulation parameters, we consider the following quantities:

- V_{sim} : The simulation volume.
- N_{part} : The total number of particles in the simulation.
- N_{pix} : The number of pixels on each lens plane.
- n_{part}^j : The number of particles in the j -th shell.
- $\bar{n}_{\text{part}}^j$: The mean number of particles per pixel in the j -th shell.

The convergence field is then given by:

$$\kappa_j(\boldsymbol{\theta}) = \frac{3H_0^2 \Omega_m}{2c^2 a_j \chi_j} \frac{V_{\text{sim}}}{N_{\text{part}}} \frac{N_{\text{pix}}}{4\pi} \left(n_{\text{part}}^j - \bar{n}_{\text{part}}^j \right), \quad (6.33)$$

2. **Potential Calculation:** To facilitate the computation of lensing effects, the lensing potential $\psi^j(\boldsymbol{\theta})$ for each lens plane is derived from the convergence field $\kappa_j(\boldsymbol{\theta})$. This is efficiently done using spherical harmonics:

$$\psi_{lm}^j = \frac{2}{l(l+1)} \kappa_{lm}^j \quad \text{for } l \neq 0, \quad (6.34)$$

and $\psi_{lm}^j = 0$ for $l = 0$. Subsequently, the lensing potential in real space is reconstructed, allowing the calculation of the deflection field $\boldsymbol{\alpha}^j(\boldsymbol{\theta})$ and the optical tidal matrix $U_{ik}^j(\boldsymbol{\theta})$ through:

$$\alpha_i^j = -\nabla_i \psi^j, \quad U_{ik}^j = \nabla_i \nabla_k \psi^j. \quad (6.35)$$

3. **Deflection Angle Determination:** The deflection field $\boldsymbol{\alpha}^j(\boldsymbol{\theta})$ and the optical tidal matrix $U_{ik}^j(\boldsymbol{\theta})$ are interpolated to arbitrary positions on each lens plane. The lensing matrix $\mathcal{A}_{ik}^j$ at each plane is then updated using the recurrence relation:

$$\begin{aligned} \mathcal{A}_{ik}^{j+1} = & \left(1 - \frac{\chi_j}{\chi_{j+1}} \frac{\chi_{j+1} - \chi_{j-1}}{\chi_j - \chi_{j-1}} \right) \mathcal{A}_{ik}^{j-1} + \frac{\chi_j}{\chi_{j+1}} \frac{\chi_{j+1} - \chi_{j-1}}{\chi_j - \chi_{j-1}} \mathcal{A}_{ik}^j \\ & - \frac{\chi_{j+1} - \chi_j}{\chi_{j-1}} U_{im}^j \mathcal{A}_{mk}^j, \end{aligned} \quad (6.36)$$

with the initial conditions:

$$\mathcal{A}_{ik}^1 = \delta_{ik}, \quad \mathcal{A}_{ik}^0 = \delta_{ik}, \quad (6.37)$$

for $j \geq 1$. The closest lens plane to the observer is designated as $j = 0$.

4. **Ray Propagation:** Light rays are propagated from the source galaxies to the observer through the sequence of lens planes. At each plane, the accumulated deflection angles are updated based on the lensing matrix $\mathcal{A}_{ik}^j$.
5. **Map Assembly:** After propagating all light rays through the lens planes, the convergence $\kappa(\boldsymbol{\theta})$ and shear $\gamma(\boldsymbol{\theta})$ fields are compiled into full-sky maps, using Eq. (3.14). These maps are typically represented on the HEALPix grid.

6.6.2 Born-approximated Ray-tracing

The Born approximation offers a simplified approach to weak lensing map generation by assuming that light rays travel along their unperturbed, straight-line paths from the source galaxies to the observer. This approximation neglects the bending of light rays due to gravitational deflections between lens planes, thereby reducing the computational complexity of the ray-tracing process (Schneider, 2006).

While this simplification can lead to faster computations, it introduces certain limitations in accurately capturing multiple deflections and non-linear lensing effects. To overcome these limitations, researchers have developed Post-Born corrections (Cooray & Hu, 2002; Dodelson et al., 2005) that account for the deflection during ray propagation and for the so-called lens-lens coupling, which describes how gravitational lenses at different redshifts can interact to generate rotational modes in the observable fields. It is shown that the Post-Born corrections can impact the higher-order moments or peak statistics of galaxy weak lensing convergence maps (Petri et al., 2017; Fabbian et al., 2019). Figure 6.5 (Petri et al., 2017) illustrates the parameter bias induced by the Born approximation in the convergence power spectrum and moments of the convergence field. The Born approximation is accurate in predicting the convergence power spectrum, but it leads to significant biases in the moments of the convergence field.




Figure 6.5: Parameter bias induced by the Born approximation in the convergence power spectrum (Left Panel) and moments of the convergence field (Right Panel). While the Born approximation is accurate in predicting the convergence power spectrum, it leads to significant biases in the moments of the convergence field. credit by Petri et al. (2017)

Nonetheless, the Born-approximation remains a valuable tool for generating weak lensing maps in large-scale structure studies (Fosalba et al., 2008; Hilbert et al., 2009; Fosalba et al., 2015a). The Born-approximated ray-tracing algorithm comprises the following key steps:

1. **Lens Plane Calculation:** For each lens plane j , the convergence contribution $\kappa_j(\boldsymbol{\theta})$ is computed independently, incorporating the lensing efficiency function $W(\chi_j, \chi_s)$, where χ_j is the comoving distance to the j -th lens plane and χ_s is the comoving distance to the source galaxy. The convergence on the j -th plane is given by

$$\kappa_j(\boldsymbol{\theta}) = W(\chi_j, \chi_s) \delta_j(\boldsymbol{\theta}) \Delta\chi_j, \quad (6.38)$$

where $\delta_j(\boldsymbol{\theta}) = n_{\text{part}}^j(\boldsymbol{\theta}) / \bar{n}_{\text{part}}^j - 1$ represents the projected matter density contrast on the j -th lens plane, and $\Delta\chi_j$ is the comoving thickness of the j -th lens plane. The lensing efficiency

function, previously defined in Eq. (3.20), is discretized for each lens plane as:

$$W(\chi_j, \chi_s) = \frac{3H_0^2\Omega_m}{2c^2} \frac{(\chi_s - \chi_j)}{\chi_s} \frac{\chi_j}{a_j}, \quad (6.39)$$

2. **Convergence Field Assembly:** The total convergence field $\kappa(\boldsymbol{\theta})$ is obtained by summing the contributions from all individual lens planes:

$$\kappa(\boldsymbol{\theta}) = \sum_j \kappa_j(\boldsymbol{\theta}). \quad (6.40)$$

This linear superposition is a direct consequence of the Born approximation, which assumes that each lens plane contributes independently to the total convergence without accounting for the altered path of the light ray due to previous deflections.

Simulation-Based Methodology for Assessing Higher-Order Weak Lensing Covariance

Contents

7.1	Constructing the BIGBOX and TILED Datasets	73
7.2	Producing Weak Lensing Convergence Maps at Multiple Redshifts	75
7.3	Incorporating Galaxy Shape Noise into Convergence Maps	76
7.4	Patch Selection and Projection	77
7.5	Applying Gaussian Kernels to Convergence Fields	80
7.6	Extracting Weak Lensing Statistics from Convergence Maps	81
7.7	Comparing Covariances from BIGBOX and TILED	84

In this work, we quantified the effect of the super-sample effect on the covariance matrix of higher-order statistics for a weak lensing survey. To achieve this, we conducted a series of N-body simulations and analyzed the resulting convergence maps. These maps enable the computation of covariance matrices for various summary statistics. Thus, we could investigate the impact of the super-sample effect on these matrices.

In the following sections, we discuss the methodology used to generate the convergence maps, extract patches for analysis, incorporate noise, apply Gaussian smoothing, and measure the summary statistics from the simulated maps. We also outline the process for estimating the covariance matrices and comparing the results between the BIGBOX and TILED simulations.

7.1 Constructing the BIGBOX and TILED Datasets

We employed the publicly available particle-mesh simulation code, FASTPM ([Feng et al., 2016](#)), to generate the simulations used in this study. As discussed in Section 6.5, FASTPM was chosen to achieve high accuracy while minimizing computational time.

The two simulations used in this work are **BIGBOX** and **TILED**. For both simulations, we adopted the same cosmological parameters as those used in the IllustrisTNG project (Nelson et al., 2019). The parameters are listed in Table 7.1.

Parameter	Symbol	Value
Hubble constant	H_0	67.74 [km s ⁻¹ Mpc ⁻¹]
Matter density	Ω_m	0.3089
Baryon density	Ω_b	0.0486
Amplitude of fluctuations	σ_8	0.8159
Spectral index	n_s	0.9667
Sum of neutrino masses	M_ν	0.0 [eV]

Table 7.1: Cosmological parameters used in the N-body simulations.

The BIGBOX simulation, conducted as part of the HalfDome project (Bayer et al., 2024), uses 6144^3 particles within a box of side length 3750 Mpc/ h , employing periodic boundary conditions. The simulation volume is replicated approximately 2.6 times per dimension to cover the redshift range $z = 0$ to $z = 4$, resulting in a total volume of about 10 (Gpc/ h)³. At the maximum redshift considered in this work, $z = 2.5$, the simulation volume is replicated approximately 1.2 times per dimension.

The TILED simulation covers a smaller volume with a side length of $L = 625$ Mpc/ h , populated with 1024^3 particles and also employing periodic boundary conditions. This combination of box size and particle number was chosen to match the resolution of the BIGBOX simulation. To cover the same redshift range as the BIGBOX simulation, the TILED box was replicated 10 times along each axis, although 6 replications are sufficient for the range $z = 0$ to $z = 2.5$. Figure 7.1 illustrates the spatial and redshift setup for the BIGBOX and TILED simulations employed in this cosmological study.

Both simulations commenced at an initial redshift of $z = 9$, utilizing an initial linear matter power spectrum at $z = 0$ generated via the CLASS code (Blas et al., 2011). The simulations were evolved over 60 time steps from $z = 9$ to the present day ($z = 0$). The resulting particle distributions were output in 80 shells spanning scale factors from $a = 0.2$ to $a = 1.0$, with a uniform spacing of $\Delta a = 0.01$. At each scale factor a_i , particles within the shells were projected onto a full-sky HEALPix grid (Górski et al., 2005) with $N_{\text{side}} = 8192$, providing an angular resolution of approximately 0.43 arcminutes, to generate mass maps.

The BIGBOX and TILED simulations were executed on the TACC (Texas Advanced Computing Center) cluster. The BIGBOX simulations required approximately 4 hours per realization, utilizing 2048 nodes, while the TILED simulations were completed in 2 hours per realization using 64 nodes.

In total, we generated 11 realizations of the BIGBOX simulation and 20 realizations of the TILED simulation, each initialized with distinct random seeds. This ensemble of realizations enables us to




Figure 7.1: Spatial and redshift setup for the BIGBOX and TILED simulations. Dashed blue grids partition the overall simulation volume into smaller, manageable tiling regions, with each tile representing a replication of the TILED simulation. The lower horizontal axis indicates the line-of-sight distance in Mpc/h , while the corresponding redshift values are displayed on the top axis.

robustly sample cosmic variance and ensures that our statistical analyses are not biased by the initial conditions of any single simulation.

It is important to note that the observer is positioned at the corner point shared by 8 replicated boxes in both the TILED and BIGBOX simulations. This placement induces the Box Replication Effect, characterized by a kaleidoscope-like pattern of heavily tiled regions along the line of sight, particularly near the equatorial regions and in directions parallel to the box edges (see Figure 7.2 for $z = 1.5$). These replicated regions can introduce artificial correlations and anisotropies in the mass maps, potentially biasing our statistical measurements. To mitigate this effect, we exclude the most heavily tiled regions from our analysis, ensuring that the Box Replication Effect does not significantly impact our results. A detailed discussion of this phenomenon and its implications is provided in Section 8.5.

7.2 Producing Weak Lensing Convergence Maps at Multiple Redshifts

Since we already have the projected mass map at each scale factor, we calculate the convergence map at each redshift following the discussion in Sections 3.2 and 6.6. As the difference between the TILED and BIGBOX simulations due to the super-sample effect becomes apparent around redshift $z \approx 1$, it is worthwhile to examine the contribution from each redshift to the convergence map.

Figure 7.3 shows the normalized lensing efficiency as a function of comoving distance (in Mpc/h)




Figure 7.2: Illustration of a 5×5 patch around $(\theta, \phi) = (\pi/2, 0)$ extracted from the TILED and BIGBOX simulations at $z = 1.5$. The TILED simulation displays a distinct kaleidoscope-like pattern caused by box replication, resulting in heavily tiled regions along the line of sight and near the equator. This pattern becomes more pronounced at higher redshifts.

for multiple source redshifts (z). The lensing efficiency curves exhibit distinct peaks at intermediate comoving distances, highlighting the regions where the matter distribution along the line of sight most significantly enhances the gravitational lensing signal

We considered source redshifts $z_s = [0.5, 1.0, 1.5, 2.0, 2.5]$, covering the range of distances relevant for both current and upcoming galaxy surveys. At low redshifts ($z < 1$), both the BIGBOX and TILED simulations are affected by super-sample effects. However, at higher redshifts ($z > 1$), these effects become more pronounced in the BIGBOX simulation. This divergence occurs because, at approximately $z = 1$, the light cone in the TILED simulation begins to extend tangentially across multiple replicated boxes. As a result, no additional super-survey modes arise within the TILED simulation beyond this redshift, suffering from the difficulty of accurately simulating the lensing mass map.

Figure 7.4 shows the convergence maps generated from both the BIGBOX and TILED simulations for a source redshift of $z_s = 1.5$. These maps are depicted on Healpix grids with $N_{\text{side}} = 8192$, consistent with the methodology described.

7.3 Incorporating Galaxy Shape Noise into Convergence Maps

In actual observations, measurements of the lensing signal are affected by noise arising from the intrinsic shapes of galaxies and inaccuracies in shape measurements. This type of noise, known as shape noise, is a major source of uncertainty, especially on small angular scales. For example, shape




Figure 7.3: Normalized lensing efficiency as a function of comoving distance for multiple source redshifts. The efficiency curves peak at intermediate comoving distances, indicating regions where the matter distribution enhances the gravitational lensing signal.

noise can differ the number densities by approximately 12% in the HSC shear catalog (See Section 5.2 of [Mandelbaum et al. 2018a](#)).

We considered four different surveys with varying galaxy number densities, as detailed in Table 1.1. The variance of the shape noise per pixel was calculated using the following expression:

$$\sigma_{\kappa, \text{noise}}^2 = \frac{\sigma_{\epsilon}^2}{2n_{\text{gal}}A_{\text{pix}}}, \quad (7.1)$$

where σ_{ϵ} is the intrinsic ellipticity dispersion of galaxies, set to $\sigma_{\epsilon} = 0.26$ ([Euclid Collaboration et al., 2019](#)), n_{gal} is the galaxy number density per square arcminute, and A_{pix} is the solid angle of a pixel, set to 0.43^2 arcminutes². Using this calculated variance, we generated a Gaussian random field, $n(\hat{\mathbf{n}})$, and added it to the convergence maps to simulate the observed signal:

$$\kappa_{\text{obs}}(\hat{\mathbf{n}}) = \kappa(\hat{\mathbf{n}}) + n(\hat{\mathbf{n}}). \quad (7.2)$$

7.4 Patch Selection and Projection

To simplify the analysis onto a flat patch, we extracted $10^\circ \times 10^\circ$ patches from the full-sky convergence maps. To maximize the number of uniformly distributed patches without repetition, we employed a Fibonacci grid ([Swinbank & James Purser, 2006](#); [Ferlito et al., 2023](#)) for extracting patches from the full-sky map. The center of each patch was positioned at the vertices of the Fibonacci grid, defined




Figure 7.4: Convergence maps from the BIGBOX and TILED simulations for source redshift $z_s = 1.5$. Yellow regions indicate positive convergence, while blue regions represent negative convergence. The maps exhibit similar large-scale structures, highlighting the agreement in the overall distribution of matter.

by golden ratio spirals:

$$\sin \theta_i = \frac{2i}{2N+1}, \quad \phi_i = \frac{2\pi i}{\varphi}, \quad -N \leq i \leq N, \quad -\frac{\pi}{2} \leq \theta_i \leq \frac{\pi}{2}, \quad (7.3)$$

where N is the number of patches and $\varphi = (1 + \sqrt{5})/2$ is the golden ratio. The visualization of the Fibonacci grid is shown in Figure 7.5. Following [Ferlito et al. \(2023\)](#), the maximum number of




Figure 7.5: Visualization of the Fibonacci grid with $N_{\text{patches}} = 273$, where each patch covers approximately $10^\circ \times 10^\circ$. After optimization and masking, the number of patches is reduced to $N_{\text{patches}} = 194$, effectively covering 47% of the sky. Each panel shows the patch distribution from a front view, top view, and the final front view after masking.

patches can be obtained by aligning the diagonals of square patches with the longitude lines. However, due to programming constraints, we extracted patches with their sides aligned to the latitude lines.

To achieve the desired $10^\circ \times 10^\circ$ shape, we first extracted patches with a larger side length and then rotated and cropped them accordingly.

Before extracting the patches, we optimized the number of patches to be extracted from the full-sky map. The number of patches, denoted N_{patches} , was determined to ensure that individual patches do not overlap, except in regions near the poles, where overlapping patches were subsequently discarded. Each patch on the full-sky map is defined as:

$$\begin{aligned} &(\theta_i - R_{\text{patch}}, \phi_i + R_{\text{patch}} \sin \theta_i), \quad (\theta_i + R_{\text{patch}}, \phi_i + R_{\text{patch}} \sin \theta_i), \\ &(\theta_i - R_{\text{patch}}, \phi_i - R_{\text{patch}} \sin \theta_i), \quad (\theta_i + R_{\text{patch}}, \phi_i - R_{\text{patch}} \sin \theta_i) \end{aligned} \quad (7.4)$$

with $R_{\text{patch}} = 5\sqrt{2} \text{ deg}$, the half diagonal length of the patch. The optimization process began with an initial count of $N_{\text{patches}} = 400$ and involved iteratively reducing this number until a configuration was achieved in which patches remained non-overlapping, except for centers located within $2R_{\text{patch}}$ of the poles. Specifically, the thresholds $|\theta_i| \geq 2R_{\text{patch}}$ and $|\phi_i| \leq \pi - 2R_{\text{patch}}$ were applied to ensure that the patches did not include the poles.

After optimization and masking, the number of patches was reduced to $N_{\text{patches}} = 273$, further decreasing to $N_{\text{patches}} = 265$ to effectively cover 64% of the sky. Additionally, patches containing points heavily tiled along the line of sight were excluded to mitigate the severe Box Replication Effect (see Section 8.5 for further details). Consequently, the final number of patches used for analysis was $N_{\text{patches}} = 194$, effectively covering 47% of the sky.

For a Fibonacci grid center characterized by coordinates (θ_i, ϕ_i) , we first employed the `gnomview` function from the `healpy` library (Zonca et al., 2019) to project each spherical patch onto a flat plane using a gnomonic projection. Each patch was then rotated by 45° around its center to align the diagonal of the patch with the longitude lines. Finally, the patch was cropped according to the corresponding vertices to achieve the desired shape. Figure 7.6 illustrates the process of extracting a patch from the full-sky convergence map and the additional steps taken to obtain the final patch.

Each patch is represented by a 2048×2048 grid of pixels, resulting in a pixel size of:

$$\Delta\theta = \frac{10^\circ}{2048} \approx 0.00488^\circ \approx 0.293' \quad \text{per pixel.} \quad (7.5)$$

For each realization, the covariance was computed using 194 patches extracted from the full-sky map of each simulation. This resulted in a total of 2134 patches from the BIGBOX simulation and 3880 patches from the TILED simulation. This ensemble of patches enables us to robustly sample both cosmic variance and shot noise, ensuring that our statistical analyses are sufficient for the power spectrum at $\ell = 5283$ and peak counts in $\kappa \in [-0.06, 0.45]$ (Petri et al., 2016).




Figure 7.6: Extraction of a patch from the full-sky convergence map using a Fibonacci grid. The patch covers an area of $10^\circ \times 10^\circ$ and is centered at a vertex of the Fibonacci grid. From left to right, the panels show the originally extracted patch, the patch rotated 45° around its center, and the final patch after a second rotation. The red-shaded region represents the final patch used for analysis.

7.5 Applying Gaussian Kernels to Convergence Fields

Shape noise primarily affects small angular scales, which can hinder the extraction of the underlying lensing signal. To mitigate this effect, we apply a Gaussian smoothing filter to the noisy convergence maps. Owing to its straightforward implementation and reliance on a single parameter, the Gaussian filter is commonly adopted for handling noisy lensing data. A discussion of alternative smoothing approaches can be found in Appendix A. The Gaussian filter is defined as:

$$W(\theta) = \frac{1}{\pi\theta_G^2} \exp\left(-\frac{\theta^2}{\theta_G^2}\right), \quad (7.6)$$

where θ is the angular distance from the center of the filter, and θ_G is the smoothing scale. For our analysis, we selected $\theta_G = 2', 5', 8',$ and $10'$.

By convolving the noisy convergence map with the Gaussian filter, we obtained the smoothed convergence map:

$$\kappa_{\text{smoothed}}(\hat{\mathbf{n}}) = \int d\Omega' W(|\hat{\mathbf{n}} - \hat{\mathbf{n}}'|) \kappa_{\text{obs}}(\hat{\mathbf{n}}'). \quad (7.7)$$

Figure 7.7 illustrates the impact of Gaussian smoothing on a noisy convergence map, demonstrating the progressive suppression of small-scale fluctuations. The figure consists of four panels, each corresponding to a different smoothing scale: $\theta_G = 2', 5', 8',$ and $10'$. As the smoothing scale increases, the Gaussian kernel increasingly attenuates small-scale noise, reducing the amplitude of fluctuations in the map. Although this process naturally diminishes the power spectrum at small angular scales, smoothing scales below $10'$ are sufficient to mitigate shape noise while retaining the non-Gaussian structures of the convergence field.

For the analysis of non-Gaussian statistics, we employ a Gaussian smoothing procedure, after which the statistics are computed from the resulting smoothed convergence maps. The application




Figure 7.7: Effect of Gaussian smoothing on a noisy convergence map. Each panel shows the result of applying a Gaussian filter with a smoothing scale of $\theta_G = 2', 5', 8',$ or $10'$. As the smoothing scale increases, small-scale noise is progressively suppressed, making large-scale structures more prominent. This demonstrates how Gaussian smoothing effectively reduces shape noise while preserving the underlying lensing signal.

of a smoothing kernel effectively suppresses small-scale structures, thereby modifying the range of κ values. To avoid the smoothing issues and the complex binning to consider, we normalize κ values by the standard deviation of each patch’s convergence map, σ_κ .

Figure 7.8 shows the mean standard deviation of the noiseless convergence maps obtained from both the BIGBOX and TILED simulations. While there is no significant difference between the two simulations, the standard deviation increases with both redshift and smoothing scale. The observed increase with redshift reflects the inclusion of additional shells along the line of sight, leading to greater variations in density contrasts. Similarly, the increase with smoothing scale arises from the smoothing process itself, which suppresses small-scale structures and amplifies the variations of larger-scale fluctuations.

7.6 Extracting Weak Lensing Statistics from Convergence Maps

To characterize the influence of super-sample effect on higher-order statistics, this study focuses on the bispectrum, probability distribution function (PDF), peak counts, minima counts, and Minkowski functionals.

Table 7.2 summarizes the range of values and the computational subroutines employed for each summary statistic.

7.6.1 Angular Power Spectrum

The angular power spectrum ($C_\ell^{\kappa\kappa}$) is estimated directly from the simulated convergence maps using `lenstools` (Petri, 2016), without any preliminary smoothing. The considered multipole range spans from $\ell = 300$ to $\ell = 3000$, partitioned into eight logarithmically spaced bins. This binning scheme is chosen to maintain consistency with the multipole selection used in the Hyper Suprime-Cam Year 3 (HSC Y3) cosmic shear analysis (Dalal et al., 2023).




Figure 7.8: Average standard deviation of the noiseless convergence maps for the BIGBOX and TILED simulations. The standard deviation increases with both redshift and smoothing scale, with only subtle differences observed between the BIGBOX and TILED simulations.

Statistic	Range	Subroutine (Sky Patch)
Angular Power Spectrum	$300 \leq \ell \leq 3000$	<code>lenstools.powerSpectrum</code>
Bispectrum	$300 \leq \ell \leq 3000$	<code>lenstools.bispectrum</code>
Peak Counts	$-4 \leq \kappa/\sigma_\kappa \leq 4$	<code>lenstools.peakCount</code>
Minima Counts	$-4 \leq \kappa/\sigma_\kappa \leq 4$	<code>lenstools.peakCount</code>
Probability Distribution Function (PDF)	$-4 \leq \kappa/\sigma_\kappa \leq 4$	<code>lenstools.pdf</code>
Minkowski Functionals	$-4 \leq \kappa/\sigma_\kappa \leq 4$	Custom implementation

Table 7.2: Summary of the summary statistics employed in this study, including their respective value ranges and the computational subroutines used for analysis.

The lower limit of $\ell = 300$ excludes large-scale modes, as the primary focus of this study is on higher-order statistics, which are most sensitive to intermediate and small angular scales. Conversely, the upper limit of $\ell = 3000$ ensures the inclusion of small-scale modes where the super-sample effect becomes increasingly significant. Furthermore, this multipole range is robust given the sampling of simulated maps, as discussed in Section 7.4.

For benchmarking, we compute the theoretical angular power spectrum using the `Halofit` model (Takahashi et al., 2012) and compare it with the measured angular power spectrum. The theoretical prediction is generated using the same set of cosmological parameters as those employed in the simulations.

7.6.2 Bispectrum

The bispectrum (B_ℓ) is computed from the unsmoothed convergence maps using `lenstools`. We consider three distinct configurations: equilateral ($\ell_1 = \ell_2 = \ell_3$), squeezed ($\ell_1 = \ell_2 = 10\ell_3$), and isosceles ($\ell_1 = \ell_2 = 2\ell_3$). These configurations are chosen to capture different shapes of the bispectrum, providing complementary information about the underlying matter distribution.

The bispectrum computations are confined to the same multipole range as the angular power spectrum, specifically $\ell \in [300, 3000]$, divided into eight logarithmically spaced bins. Although the bispectrum is more sensitive to noise and small-scale structures, maintaining an identical multipole range facilitates direct comparison with the angular power spectrum results.

7.6.3 PDF

We compute the probability distribution function (PDF) of the convergence field using `lenstools`. Each PDF is derived from the smoothed convergence maps over a normalized range of $-4 \leq \kappa/\sigma_\kappa \leq 4$, divided into eight equally spaced bins. Here, σ_κ represents the standard deviation of the convergence field within each individual patch. This range is chosen to align with the study by [Thiele et al. \(2023\)](#), while the binning is optimized to ensure that each bin contains a sufficient number of data points.

The theoretical prediction for the PDF is generated using the `hmpdf` code ([Thiele et al., 2020](#)), employing the same linear matter power spectrum as that used in the simulations. The theoretically predicted PDF is then compared with the measured PDF to evaluate the consistency and accuracy of the underlying cosmological model and its assumptions.

7.6.4 Peak/Minima Counts

Peak and minima counts are computed from the smoothed convergence maps using `lenstools`, utilizing the same normalized range as the PDF: $-4 \leq \kappa/\sigma_\kappa \leq 4$, divided into eight evenly spaced bins. While the highest and lowest peaks are represented by relatively few data points, these extreme bins are retained to ensure consistency with the binning scheme adopted for the PDF.

7.6.5 Minkowski Functionals

The Minkowski functionals are calculated from the smoothed convergence maps using a custom implementation that adheres closely to the definitions outlined in Section 4.6. As with the PDF and peak/minima counts, these functionals are computed over the normalized range $-4 \leq \kappa/\sigma_\kappa \leq 4$, divided into eight linearly spaced bins.

Figure 7.9 illustrates the variance of the gradient field, $\sigma_1 = \sqrt{\kappa_x^2 + \kappa_y^2}$, which is a critical input for calculating the Minkowski functionals. As the smoothing scale increases, σ_1 decreases, reflect-

ing the reduced influence of small-scale structures. Differences between the BIGBOX and TILED simulations, as well as across varying redshifts, remain subtle, highlighting the robustness of these measures and their relative insensitivity to large-scale environmental variations.




Figure 7.9: Average $\sigma_1 = \sqrt{\kappa_x^2 + \kappa_y^2}$ of the noiseless convergence maps for the BIGBOX and TILED simulations. These values are calculated for the smoothing scales $\theta_G = 2', 5', 8',$ and $10'$. σ_1 decreases as the smoothing scale increases, while differences between the BIGBOX and TILED simulations, or across redshifts, remain subtle.

7.7 Comparing Covariances from BIGBOX and TILED

After the measurement phase, this study investigates the impact of super-sample effect on the covariance matrices of the summary statistics discussed earlier. To facilitate this analysis, we use an unbiased estimator for the covariance matrix, as defined in Equation 7.8.

$$\text{Cov}(\mathcal{O}_i, \mathcal{O}_j) = \frac{1}{N_{\text{sim}} - 1} \sum_{n=1}^{N_{\text{sim}}} (\mathcal{O}_i^{(n)} - \langle \mathcal{O}_i \rangle) (\mathcal{O}_j^{(n)} - \langle \mathcal{O}_j \rangle), \quad (7.8)$$

Additionally, we compute the correlation matrix for each summary statistic to analyze the interdependence between different scales and configurations. The correlation matrix is defined as:

$$\rho_{ij} = \frac{\text{Cov}(\mathcal{O}_i, \mathcal{O}_j)}{\sqrt{\text{Cov}(\mathcal{O}_i, \mathcal{O}_i) \text{Cov}(\mathcal{O}_j, \mathcal{O}_j)}}, \quad (7.9)$$

where $\mathcal{O}_i$ and $\mathcal{O}_j$ represent the i -th and j -th summary statistics, respectively. Correlation matrix normalize off-diagonal elements by the scale dependent variance, thereby enabling a direct comparison between different simulations and summary statistics.

After computing the covariance and correlation matrices for both the BIGBOX and TILED simulations, we assess the impact of super-sample effect by comparing the two sets of matrices. The comparison is performed by calculating the ratio of the BIGBOX to TILED matrices, defined as:

$$R_{ij}^{\text{Cov}} = \frac{\text{Cov}_{ij}^{\text{BIGBOX}}}{\text{Cov}_{ij}^{\text{TILED}}}, \quad R_{ij}^{\rho} = \frac{\rho_{ij}^{\text{BIGBOX}}}{\rho_{ij}^{\text{TILED}}}, \quad (7.10)$$

where $\text{Cov}^{\text{BIGBOX}}$ and $\text{Cov}^{\text{TILED}}$ denote the covariance matrices derived from the BIGBOX and TILED simulations, respectively. Similarly, ρ^{BIGBOX} and ρ^{TILED} represent the corresponding correlation matrices.

To quantitatively evaluate the overall influence of super-sample effect, we compute the mean values of the ratios R_{ij}^{Cov} and R_{ij}^{ρ} across the respective covariance and correlation matrices. Specifically, the average of R^{ρ} is calculated by excluding the diagonal elements, which are equal to unity by definition. In contrast, the average of R^{Cov} utilizes all elements of the covariance matrix, including the diagonal.

For ℓ -binned statistics, such as the angular power spectrum ($C_{\ell}^{\kappa\kappa}$) and the bispectrum, the average ratio is determined across all available multipole bins. Conversely, for ν -binned statistics, such as the PDF, peak/minima counts, and Minkowski functionals, the first and last bins are excluded from the ratio calculation to avoid biases introduced by limited data points and the inherent unreliability of these extreme bins. The average ratios are defined as follows:

$$\langle R^{\text{Cov}} \rangle = \frac{1}{N_{\text{bins}}^2} \sum_{i,j} R_{ij}^{\text{Cov}}, \quad \langle R^{\rho} \rangle = \frac{1}{N_{\text{bins}}^2 - N_{\text{bins}}} \sum_{i \neq j} R_{ij}^{\rho}, \quad (7.11)$$

where $N_{\text{bins}} = 8$ for ℓ -binned statistics and $N_{\text{bins}} = 6$ for ν -binned statistics.

Comparative Analysis of Statistical Measures and Covariance Structures in BIGBOX and TILED

Contents

8.1	Comparison of Mean and Variance for Summary Statistics	86
8.2	Analysis of Covariance and Correlation Matrices	90
8.3	Effects of Noise on Summary Statistics	96
8.4	Influence of Smoothing Scale	99
8.5	Systematic Effects: Box Replication Artifact	102

This chapter presents a comprehensive comparison of weak lensing statistics derived from two distinct simulation sets: BIGBOX and TILED. The primary objective is to quantify the differences in mean values, covariance matrices, and correlation matrices between these simulations to assess the impacts of super-sample covariance and other systematic effects on the measurements. Additionally, the chapter investigates the influence of shape noise, varying smoothing scales, and box replication artifacts on the results. Unless otherwise specified, the main findings are based on noiseless simulations with a smoothing scale of 2 arcminutes. Furthermore, the power spectrum (C_ℓ) and bispectrum results are normalized by factors of $\ell(\ell + 1)/2\pi$ and ℓ^4 , respectively, to facilitate meaningful comparisons.

8.1 Comparison of Mean and Variance for Summary Statistics

In this section, we analyze and compare the mean (μ) and variance (σ^2) of summary statistics derived from the BIGBOX and TILED simulations. Figures 8.1 and 8.2 illustrate these comparisons for the angular power spectrum, bispectrum, and other higher-order statistics, highlighting the differences arising from simulation methodologies and super-sample effect.

The mean values, represented by the ratio R_μ , demonstrate remarkable consistency in capturing statistical distributions across the two simulation sets. For most summary statistics, deviations in R_μ

are less than 1%, underscoring the robustness of both simulation pipelines in reproducing the underlying statistics. Moreover, comparisons with theoretical predictions (`Halofit` and `hmpdf`) reveal no significant conflicts between the theoretical models and the simulation results.

However, systematic deviations between TILED and BIGBOX simulations emerge at low ν values for peak counts, minima, and Minkowski Functionals, as well as at high ν values for minima (see the left panels in Figure 8.2). These deviations likely result from the limited resolution of the TILED simulations in capturing super high-density and low-density regions, leading to fewer data points in the extreme tails of the distribution.

In contrast, the variance ratios R_{σ^2} exhibit more pronounced differences, highlighting the critical impact of super-sample effect on variance measurements. Specifically, the variance of the angular power spectrum (top-right panel in Figure 8.1) shows a significant increase in the BIGBOX simulations at high multipole moments (ℓ). This increase is consistent with the expected amplification due to the inclusion of larger-scale modes in the BIGBOX setup, which are absent in the TILED simulations. Furthermore, discrepancies in variance ratios grow with increasing source redshift, reflecting the progressive loss of large-scale modes in the TILED simulations. For ν -binned statistics (right panels in Figure 8.2), the BIGBOX simulations consistently yield higher variances across nearly all ν bins, with this trend becoming more pronounced at higher source redshifts. These findings emphasize the necessity of accounting for super-sample effect and other systematic effects to ensure accurate cosmological parameter estimation from weak lensing statistics.




Figure 8.1: Comparison of the mean and variance for the angular power spectrum (C_ℓ^{KK}) and the bispectrum components across three configurations: equilateral ($B_\ell^{(eq)}$), isosceles ($B_\ell^{(iso)}$), and squeezed ($B_\ell^{(sq)}$). Results are presented for the BIGBOX and TILED simulations across source redshifts $z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5 . The figure highlights how different simulation methodologies and super-sample effect influence the summary statistics of these cosmological observables.




Figure 8.2: Comparison of the mean and variance for ν -binned statistics, including the Minkowski Functionals (V_0, V_1, V_2), the probability density function (PDF), and peak and minima counts. Results are derived from the BIGBOX and TILED simulations across source redshifts $z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5 , highlighting their sensitivity to super-sample effect. The figure illustrates how super-sample effect and other systematic effects influence the distribution and variance of these higher-order weak lensing statistics.

8.2 Analysis of Covariance and Correlation Matrices

In this section, we quantify the impact of super-sample effect on the covariance and correlation matrices of various summary statistics derived from the BIGBOX and TILED simulations. Figures 8.4 and 8.5 present the covariance matrices for the BIGBOX and TILED simulations, respectively, revealing that while the general covariance structures are consistent between the two simulation sets, the BIGBOX simulations exhibit higher covariance values on average. Figure 8.6 illustrates the corresponding correlation matrices, where the upper-left triangle represents the BIGBOX results and the lower-right triangle shows the TILED results. The color scale indicates that the correlation structures are almost consistent between the two simulation sets.

Figures 8.7 and 8.8 depict the ratios of the covariance and correlation matrices between BIGBOX and TILED, denoted as R_{Cov} and R_ρ , respectively. These ratios provide a quantitative assessment of super-sample effect's influence on both the diagonal and off-diagonal elements of the covariance structure. The covariance ratios R_{Cov} generally exceed unity, ranging from 10% to 30% above the baseline values for most summary statistics, underscoring super-sample effect's significant contribution to the overall covariance matrix. Conversely, the correlation ratios R_ρ typically deviate by less than 5%, suggesting that while super-sample effect elevates the covariance, it has a minimal impact on the correlation coefficients. However, notable exceptions are observed in the angular power spectrum and peak/minima counts. For the angular power spectrum, larger deviations in R_ρ are consistent with super-sample effect, which introduce strong correlations through shared large-scale modes, particularly at small multipole moments (ℓ). In the case of peak and minima counts, the deviations in correlation ratios likely arise from systematic shifts in peak positions within the TILED simulations, affecting correlations near the peak bins. For a summary of the average ratios of covariance and correlation matrices, refer to Figure 8.3.




Figure 8.3: Average ratios of covariance matrices (left) and correlation matrices (right) between the BIGBOX and TILED simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0, \text{ and } 2.5$). The increasing trend in the covariance ratios (R_{Cov}) highlights the significant contribution of super-sample effect to the covariance structure. Conversely, the correlation ratios (R_ρ) remain relatively stable, indicating minimal impact on the correlation coefficients.




Figure 8.4: Covariance matrices of various summary statistics in the BIGBOX simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5). The matrices illustrate the covariance structure influenced by super-sample effect, with higher covariance values observed on average compared to the TILED simulations.




Figure 8.5: Covariance matrices of various summary statistics in the TILED simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5), using a color scale consistent with Figure 8.4. These matrices demonstrate generally lower covariance values compared to the BIGBOX simulations, highlighting the reduced influence of super-sample effect due to the absence of larger-scale modes.




Figure 8.6: Correlation matrices of various summary statistics in the BIGBOX and TILED simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5). The upper-left triangle of each panel displays the correlation coefficients from the BIGBOX simulations, while the lower-right triangle shows the corresponding values from the TILED simulations. The similar color scales indicate that the correlation structures are almost consistent between the two simulation sets.




Figure 8.7: Ratios of covariance matrices (R_{Cov}) between the BIGBOX and TILED simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5). The ratios consistently range from 10% to 30% above unity for most measured statistics, indicating the substantial impact of super-sample effect on the covariance structure.




Figure 8.8: Ratios of correlation matrices (R_p) between the BIGBOX and TILED simulations across multiple source redshifts ($z_s = 0.5, 1.0, 1.5, 2.0$, and 2.5). The ratios generally deviate by less than 5%, indicating that while super-sample effect elevates the entire covariance structure, its impact on the correlation coefficients remains minimal.

8.3 Effects of Noise on Summary Statistics

The presence of shape noise in convergence maps can influence the covariance and correlation structures of weak lensing summary statistics. To evaluate the interaction between shape noise and super-sample effect, we analyzed multiple noise levels corresponding to those expected in current and future surveys (see Table 1.1). This analysis aims to quantify the extent to which super-sample effect dominates over the contribution from shape noise. Since the bispectrums exhibit no clear trends in the noiseless case (as shown in Figures 8.7 and 8.8), we focus our analysis on the angular power spectrum and select higher-order statistics.

Figures 8.9 and 8.10 present the covariance matrix ratios and their averages across summary statistics for the BIGBOX and TILED simulations at various noise levels. The angular power spectrum and peak/minima counts exhibit greater sensitivity to noise, particularly at high noise levels (e.g., DES and HSC surveys), with covariance ratios consistently increasing across most statistics. This trend suggests that in future surveys, super-sample effect is expected to dominate over noise contributions.

Conversely, Figures 8.11 and 8.12 display the correlation matrix ratios and their averages across summary statistics at different noise levels. These ratios generally remain below 10% for most statistics, indicating that noise has a minimal impact on the overall covariance structure. However, higher values in the correlation ratios are concentrated near the boundaries of each summary statistic, arising from regions where data points are sparse and measurements are more sensitive to noise amplification. These findings underscore the necessity of accounting for super-sample effect in weak lensing analyses to ensure accurate cosmological parameter estimation, especially as survey sensitivities improve and noise levels vary.




Figure 8.9: Average ratio of covariance matrices for summary statistics between the BIGBOX and TILED simulations at different shape noise levels (see Table 1.1). The increasing trend in covariance ratios (R_{Cov}) remains consistent across most summary statistics.




Figure 8.10: Ratio of covariance matrices (R_{Cov}) for summary statistics between the BIGBOX and TILED simulations at varying shape noise levels. The covariance ratios exhibit a consistent increasing trend, highlighting the persistent influence of super-sample covariance across different noise conditions.




Figure 8.11: Average ratio of correlation matrices for summary statistics between the BIGBOX and TILED simulations at varying shape noise levels (see Table 1.1). The correlation ratios (R_ρ) remain below 10% for most summary statistics.




Figure 8.12: Ratios of correlation matrices (R_ρ) between the BIGBOX and TILED simulations across multiple shape noise levels (see Table 1.1). Higher correlation ratios are concentrated near the boundaries of summary statistics, resulting from sparse data points and amplified noise effects in these regions.

8.4 Influence of Smoothing Scale

In this section, we analyze the impact of varying Gaussian smoothing scales on higher-order summary statistics derived from convergence maps, focusing on how smoothing deviates their underlying structures. Increasing the smoothing scale progressively suppresses small-scale features, redistributing signal intensities across different scales. Notably, smoothed convergence maps were not utilized for the calculation of the angular power spectrum and bispectrum; therefore, the results presented herein correspond exclusively to other higher-order statistics.

Figures 8.13 and 8.14 illustrate the changes in covariance matrix ratios and their averages as a function of the Gaussian smoothing scale. Across all smoothing scales, the average covariance ratios between the BIGBOX and TILED simulations exhibit a consistent upward trend for most summary statistics. However, notable variations occur in the peak and minima counts, particularly at larger smoothing scales. These variations are attributed to the averaging methodology, where contributions from the edges of ν bins and peak bins become increasingly significant at higher smoothing levels.

Similarly, Figures 8.15 and 8.16 present the ratios of correlation matrices and their averages across different Gaussian smoothing scales. The correlation ratios generally remain below 5% for most summary statistics, indicating that the overall structure of the covariance matrix is robust to variations in smoothing scale. However, consistent with the covariance trends, peak and minima counts exhibit elevated correlation ratios, especially at larger smoothing scales. These elevated ratios arise from regions with sparse data points, making measurements more susceptible to noise amplification.

In summary, these findings suggest that while Gaussian smoothing scales can influence individual summary statistics, the overall structural integrity of the covariance and correlation matrices remains largely unaffected by changes in the smoothing scale. This robustness underscores the reliability of the covariance structures in weak lensing analyses, even when different smoothing scales are applied to the convergence maps.




Figure 8.13: Average ratio of covariance matrices (R_{Cov}) for various summary statistics between the BIGBOX and TILED simulations at different Gaussian smoothing scales, illustrating the effect of smoothing on covariance ratios. The overall increasing trend persists across all smoothing scales, with pronounced variations in peak and minima counts becoming more evident at larger smoothing scales due to the increased significance of edge contributions in ν bins.




Figure 8.14: Ratio of covariance matrices (R_{Cov}) for various summary statistics between the BIGBOX and TILED simulations across different Gaussian smoothing scales. The figure highlights consistent trends in covariance ratios regardless of the applied smoothing scale, demonstrating the persistent influence of super-sample effect across varying smoothing conditions.




Figure 8.15: Average ratio of correlation matrices (R_ρ) for various summary statistics between the BIGBOX and TILED simulations at different Gaussian smoothing scales. While the angular power spectrum, peak counts, and minima counts exhibit greater sensitivity to noise, the correlation ratios remain below 5% for most summary statistics. Elevated correlation ratios in peak and minima counts at higher smoothing scales are attributed to edge effects in ν bins.




Figure 8.16: Ratios of correlation matrices (R_ρ) for various summary statistics between the BIGBOX and TILED simulations across different Gaussian smoothing scales. The correlation ratios generally remain consistently modest across all smoothing scales, except near the boundaries of ν bins. Elevated ratios in these regions are due to sparse data points and amplified noise effects.

8.5 Systematic Effects: Box Replication Artifact

In weak lensing simulations, extending the simulated volume often necessitates replicating finite simulation boxes to create a larger effective volume. This box replication process can introduce artificial correlations in the data, particularly near the boundaries or along preferred replication directions. Such artifacts may compromise the reliability of higher-order summary statistics. To systematically investigate this issue, we categorize simulation regions into two distinct groups: regions significantly affected by artifacts, termed **Replication-Influenced Patches (RIP)**, and regions minimally affected by artifacts, termed **Replication-Minimal Patches (RMP)**.

The criteria for categorizing patches are as follows:

- **Replication-Influenced Patches (RIP):**

- *Patches near the equator:*

$$\left| \theta_i - \frac{\pi}{2} \right| \leq R_{\text{patch}}$$

- *Patches near the edges of octants:*

$$\left| \phi_i - \frac{k\pi}{2} \right| \leq R_{\text{patch}} \quad \text{for } k = 0, 1, 2, 3$$

- **Replication-Minimal Patches (RMP):**

- All other patches not meeting the above criteria.

where (θ_i, ϕ_i) denote the center of patch i , and $R_{\text{patch}} = 5\sqrt{2} \text{ deg}$ is the half-diagonal of the patch.




Figure 8.17: Angular power spectrum for the Replication-Influenced Patches (RIPs) heavily influenced by box replication artifacts. The systematic biases introduced by these artifacts are evident in the deviations from the values observed in other patches.

Our analysis indicates that patches containing the point $(\theta_i, \phi_i) = (\pi/2, 0)$ exhibit significant deviations in both the mean and variance of summary statistics compared to other RIPs. To prevent these outliers from skewing our results, we exclude this specific point from subsequent analyses.

Figure 8.17 presents the angular power spectrum for the heavily influenced RIPs, highlighting the systematic biases introduced by box replication artifacts.

After these adjustments, we obtained 70 RMPs per realization, resulting in a total of 1,400 RMPs for the TILED simulations and 770 RMPs for the BIGBOX simulations across all realizations.

Figure 8.20 presents the ratios of mean and variance for various summary statistics between Replication-Influenced Patches (RIPs) and Replication-Minimal Patches (RMPs). The analysis reveals that the mean angular power spectra in RIPs are systematically underestimated by approximately 0.5%. Additionally, the ν -binned statistics exhibit elevated mean values in low ν bins and reduced values in high ν bins. This pattern suggests that box replication intensifies both overdense and underdense regions, leading to more extreme convergence values. Specifically, in the TILED simulations, the limited resolution in underdense regions causes a shift of the ν -binned statistics toward lower ν bins. Conversely, the variance ratios approach unity, indicating that the box replication effect amplifies variance to levels comparable to those observed in the BIGBOX simulations. Notably, the dependence on source redshift is effectively nullified in RIPs, implying that box replication obscures the super-sample variance.




Figure 8.18: Average BIGBOX/TILED ratios of covariance matrices for Replication-Influenced Patches (RIP) and Replication-Minimal Patches (RMP). The ratios for RIPs (right panel) are consistently lower than those for RMPs (left panel), indicating that box replication artifacts reduce covariance ratios in replication-influenced regions.




Figure 8.19: Average BIGBOX/TILED ratios of correlation matrices for Replication-Influenced Patches (RIP) and Replication-Minimal Patches (RMP). The ratios for RIPs (right panel) are consistently lower than those for RMPs (left panel), except at $z_s = 0.5$, where super-sample effects impact both TILED and BIGBOX simulations similarly.

Figures 8.21 and 8.22 display the ratios of covariance and correlation matrices between TILED and BIGBOX simulations for RIPs. Additionally, Figures 8.18 and 8.19 present the average ratios for both RIPs and RMPs. The main observations are that RIPs consistently exhibit lower covariance ratios compared to RMPs. Similarly, the correlation ratios for RIPs are generally lower than those for RMPs, except at $z_s = 0.5$, where the super-sample effect is present in both TILED and BIGBOX simulations. Furthermore, distinct covariance structures, such as those for C_ℓ and V_0 , emerge due to box replication artifacts. Structural transitions observed in previous analyses occur at higher redshifts ($z_s = 2.0 \sim 2.5$) in RIPs, aligning with RMPs as higher redshift regions become dominated by RMPs.




Figure 8.20: Ratios of the mean and variance of summary statistics between Replication-Influenced Patches (RIP) and Replication-Minimal Patches (RMP). The ν -binned statistics exhibit elevated mean values in low ν bins and reduced mean values in high ν bins. Variance ratios are generally close to unity, indicating that box replication amplifies variance to levels comparable to those in BIGBOX simulations.




Figure 8.21: BIGBOX/TILED ratios of covariance matrices specifically for Replication-Influenced Patches (RIP). Compared to Figure 8.8, RIP ratios are consistently lower than those for RMPs. Covariance structures unique to RIPs, such as C_l and V_0 , indicate that box replication artifacts can generate distinct covariance structures.




Figure 8.22: BIGBOX/TILED ratios of correlation matrices specifically for Replication-Influenced Patches (RIP). Similar to covariance ratios, RIP correlation ratios are consistently lower than those for RMPs. Off-diagonal structures differ from those in Figure 8.8 at lower redshifts but are likely to become more similar at higher redshifts.

Conclusions

In this study, we performed a set of simulations: TILED and BIGBOX, to investigate the impacts of super-sample effect on the covariance and correlation matrices of the power spectrum and higher-order statistics. We utilized 11 realizations of state-of-the-art simulations, using 6144^3 particles within $3750 \text{ Mpc}/h$ from the HalfDome project (Bayer et al., 2024) as BIGBOX, and accordingly computed 20 realizations of the same volume tiled with smaller boxes with the same cosmology, where each box component uses 1024^3 particles within $625 \text{ Mpc}/h$, as TILED. We considered the convergence field propagated under the Born approximation. We extracted $10^\circ \times 10^\circ$ patches from full-sky convergence maps via a Fibonacci grid, achieving uniform coverage with minimal overlap.

To probe non-Gaussian features of the convergence field, we employ a suite of higher-order statistics, including the angular power spectrum, bispectrum, probability distribution function (PDF), peak and minima counts, and Minkowski functionals. These measures are computed using both established tools and custom implementations, with validation against theoretical models such as `Halofit` and `hmpdf`. By examining a wide range of scales, from ℓ -binned statistics ($\ell \in [300, 3000]$) to ν -binned statistics ($-4 \leq \kappa/\sigma_\kappa \leq 4$), we capture the broad impact of the super-sample effect across different observational regimes.

Below, we summarize our key findings and their implications, based on a comparative analysis of statistical measures and their covariance and correlation matrices in BIGBOX and TILED simulations:

- Comparing mean values of statistical measures between BIGBOX and TILED simulations revealed consistent results within 1% across all statistics, redshifts, and multipoles. This agreement underscores the reliability of the simulations in capturing the underlying cosmological model.
- Despite similarity in mean values, variance ratios differ by 10%-30%, with the largest deviations occurring at high multipoles and redshifts. These discrepancies arise from the super-sample effect, which is driven by large-scale modes present in BIGBOX but absent in TILED.
- shape noise notably affects certain observables, particularly the angular power spectrum and

peak/minima counts, with deviations becoming more pronounced at higher noise levels. Nonetheless, the super-sample effect remains dominant, and correlation matrices exhibit minimal structural changes ($< 5\%$), indicating robust covariance estimation even in the presence of noise.

- Varying the Gaussian smoothing scale primarily modifies small-scale map features without substantially altering covariance or correlation matrices. Higher-order statistics also show only modest changes, supporting the notion that the super-sample effect is largely independent of smoothing scales.
- Box replication artifacts introduce systematic challenges, leading to $\sim 2\%-5\%$ biases in mean summary statistics within Replication-Influenced Patches (RIP). Covariance and correlation ratios in these patches deviate by up to 10% relative to BIGBOX, indicating higher covariance levels and potentially obscuring redshift-dependent trends associated with the super-sample variance. These findings emphasize the need to mitigate box replication artifacts for unbiased simulation-based analyses.

These results have significant implications for weak lensing surveys such as LSST, Euclid, and Roman. The substantial influence of super-sample variance underscores the importance of including large-scale modes in simulations to ensure accurate variance modeling. The demonstrated robustness of the super-sample effect to shape noise and smoothing scales further reinforces its relevance across diverse observational settings. However, mitigating systematic effects, particularly box replication artifacts, remains critical to prevent biased interpretations of higher-order statistics.

A dimensional analysis suggests that $R_{\sigma^2} \propto (L^2/\Omega_W)^\alpha$, where L is the simulation box size, Ω_W is the survey area, and α is an undetermined exponent. Accordingly, super-sample variance becomes more pronounced for larger surveys when box sizes are relatively small. For instance, in a survey with an area of $18,000 \text{ deg}^2$ (e.g., LSST), achieving super-sample variance levels comparable to TILED would require a simulation box size of $625 \text{ Mpc}/h \times \sqrt{18,000/100} \approx 8 \text{ Gpc}/h$.

Future work will focus on the following aspects:

- Developing methods to mitigate box replication artifacts, especially in simulations utilizing simply replicated volumes.
- Extending the analysis to incorporate additional cosmological models and baryonic effects to enhance simulation realism.
- Integrating these insights into survey design and data analysis pipelines to improve the precision of cosmological parameter inference.

By refining simulation techniques and observational approaches, weak lensing remains a powerful method for probing fundamental cosmic properties and constraining cosmological models with increasingly stringent accuracy.

Chapter A

Filtering Techniques in Cosmological Lensing

A.1 Gaussian Filter

A straightforward approach to smoothing noisy lensing data is the Gaussian filter. Gaussian filtering is simple to implement, requiring only the choice of σ . Larger σ effectively suppresses high-frequency noise, though it also blurs small-scale structures, such as strong peaks and void edges, limiting its effectiveness for extracting higher-order cosmological information. In two dimensions, a Gaussian kernel G with standard deviation σ is given by:

$$G(\mathbf{x}; \sigma) = \frac{1}{2\pi \sigma^2} \exp\left(-\frac{\|\mathbf{x}\|^2}{2\sigma^2}\right). \quad (\text{A.1})$$

If $\kappa(\mathbf{x})$ represents the unsmoothed convergence map, then its Gaussian-smoothed counterpart is:

$$\kappa_{\text{smooth}}(\mathbf{x}) = [\kappa(\mathbf{x}) * G(\mathbf{x}; \sigma)], \quad (\text{A.2})$$

where $*$ denotes a 2D convolution.

A.2 Wiener Filter

A more sophisticated filtering technique for reconstructing cosmological signals in the presence of noise is the Wiener filter (Wiener, 1949; Starck et al., 2021). The Wiener filter aims to minimize the mean-square error between an estimated signal $\hat{\kappa}$ and the true signal κ . If both signal and noise can be approximated as Gaussian random fields, the Wiener filter is the optimal linear filter in a least-squares sense. The Wiener filter is highly relevant in lensing studies, particularly for kaiser-Squires inversion (Kaiser & Squires, 1993). In Fourier or harmonic space, one commonly encounters:

$$\hat{\kappa}(\ell) = \frac{C_{\ell}^{\kappa\kappa}}{C_{\ell}^{\kappa\kappa} + N_{\ell}} \tilde{\kappa}(\ell), \quad (\text{A.3})$$

where $\kappa(\ell)$ is the convergence field in harmonic space, $\gamma(\ell)$ is the observed shear field in harmonic space, $C_\ell^{\kappa\kappa}$ is the theoretical convergence power spectrum, N_ℓ is the noise power spectrum, and $\tilde{\kappa}(\ell)$ is the Fourier transform of the observed noisy convergence map.

The ratio $\frac{C_\ell^{\kappa\kappa}}{C_\ell^{\kappa\kappa} + N_\ell}$ acts as a mode-by-mode weighting: modes with high signal-to-noise ($C_\ell^{\kappa\kappa} \gg N_\ell$) pass through unattenuated, while low signal-to-noise modes ($C_\ell^{\kappa\kappa} \ll N_\ell$) are strongly suppressed.

A.3 Starlet Filter

The Starlet filter is a *wavelet transform* that satisfies the admissibility condition for analyzing data across multiple scales simultaneously. Let $I(x, y)$ be the original 2D map (e.g., a convergence map in cosmological lensing). The Starlet transform decomposes I into a coarse or residual approximation $c_J(x, y)$ and $j_{\max}$ wavelet (detail) images $w_j(x, y)$ at different resolution scales:

$$I(x, y) = c_J(x, y) + \sum_{j=1}^{j_{\max}} w_j(x, y), \quad (\text{A.4})$$

where each wavelet scale w_j captures structures corresponding to a dyadic scale 2^j in spatial size, and $J = j_{\max} + 1$. The Starlet wavelet function is derived from a cubic B-spline, making the transform particularly smooth. In one dimension, the generating function for the Starlet can be written in terms of a scaling function ϕ and its dilations. Specifically, the Starlet wavelet is obtained by applying:

$$\psi(t_1, t_2) = 4[\phi(2t_1) - \phi(2t_2)], \quad (\text{A.5})$$

where ϕ is the cubic B-spline function of order 3. A common 1D representation of the B-spline function $\phi(t)$ is (for $t \geq 0$):

$$\phi(t) = \frac{1}{12} \left(t^3 - 4t^2 + 6t - 4 \right)_+ \quad (\text{appropriately shifted and scaled}), \quad (\text{A.6})$$

with ‘ $(\cdot)_+$ ’ indicating that $\phi(t)$ is zero outside a finite support. In two dimensions, the Starlet basis is separable, defined by products of the 1D scaling function or its wavelet function:

$$\Phi(x, y) = \phi(x) \phi(y), \quad \Psi(x, y) = \psi(x) \psi(y). \quad (\text{A.7})$$

For a detailed derivation of these functions and the full Starlet transform algorithm, see [Starck et al. \(2010\)](#).

Bibliography

- Abbott B. P., et al., 2016, [Phys. Rev. Lett.](#), **116**, 061102
- Abbott T. M. C., et al., 2018, [ApJS](#), **239**, 18
- Abbott T. M. C., et al., 2021, [ApJS](#), **255**, 20
- Abbott T. M. C., et al., 2022, [Phys. Rev. D](#), **105**, 023520
- Aihara H., et al., 2018, [PASJ](#), **70**, S4
- Albrecht A., et al., 2006, [arXiv e-prints](#), pp astro-ph/0609591
- Alpher R. A., Herman R., 1948, [Nature](#), **162**, 774
- Amon A., et al., 2022, [Phys. Rev. D](#), **105**, 023514
- Angulo R. E., Springel V., White S. D. M., Jenkins A., Baugh C. M., Frenk C. S., 2012, [MNRAS](#), **426**, 2046
- Armijo J., Marques G. A., Novaes C. P., Thiele L., Cowell J. A., Grandón D., Shirasaki M., Liu J., 2024, [arXiv e-prints](#), p. arXiv:2410.00401
- Asgari M., et al., 2021, [A&A](#), **645**, A104
- Bardeen J. M., Bond J. R., Kaiser N., Szalay A. S., 1986, [ApJ](#), **304**, 15
- Barnes J., Hut P., 1986, [Nature](#), **324**, 446
- Barreira A., Schmidt F., 2017, [J. Cosmology Astropart. Phys.](#), **2017**, 051
- Barreira A., Krause E., Schmidt F., 2018, [J. Cosmology Astropart. Phys.](#), **2018**, 015
- Bartelmann M., 2010, [Classical and Quantum Gravity](#), **27**, 233001
- Bartelmann M., Schneider P., 2001, [Phys. Rep.](#), **340**, 291
- Bayer A. E., Liu J., Terasawa R., Barreira A., Zhong Y., Feng Y., 2023, [Phys. Rev. D](#), **108**, 043521
- Bayer A. E., Zhong Y., Li Z., DeRose J., Feng Y., Liu J., 2024, [arXiv e-prints](#), p. arXiv:2407.17462

Bennett C. L., et al., 2003, [ApJS](#), **148**, 97

Berger M. J., Colella P., 1989, [Journal of Computational Physics](#), **82**, 64

Bernardeau F., Colombi S., Gaztañaga E., Scoccimarro R., 2002, [Phys. Rep.](#), **367**, 1

Bernstein G. M., 2010, [MNRAS](#), **406**, 2793

Blas D., Lesgourgues J., Tram T., 2011, [J. Cosmology Astropart. Phys.](#), **2011**, 034

Bond J. R., Efstathiou G., 1987, [MNRAS](#), **226**, 655

Born M., 1926, [Zeitschrift fur Physik](#), **38**, 803

Boyle A., Uhlemann C., Friedrich O., Barthelemy A., Codis S., Bernardeau F., Giocoli C., Baldi M., 2021, [MNRAS](#), **505**, 2886

Carroll S. M., Press W. H., Turner E. L., 1992, [ARA&A](#), **30**, 499

Chan K. C., Moradinezhad Dizgah A., Noreña J., 2018, [Phys. Rev. D](#), **97**, 043532

Chen Z., Yu Y., 2024, [MNRAS](#), **534**, 1205

Cheng S., Yu H.-R., Inman D., Liao Q., Wu Q., Lin J., 2020, [arXiv e-prints](#), p. [arXiv:2003.03931](#)

Chisari N. E., et al., 2018, [MNRAS](#), **480**, 3962

Cooley J. W., Tukey J. W., 1965, [Mathematics of Computation](#), **19**, 297

Cooray A., Hu W., 2002, [ApJ](#), **574**, 19

Couchman H. M. P., 1991, [ApJ](#), **368**, L23

Couchman H. M. P., Thomas P. A., Pearce F. R., 1995, [ApJ](#), **452**, 797

Crain R. A., et al., 2015, [MNRAS](#), **450**, 1937

Crocce M., Castander F. J., Gaztañaga E., Fosalba P., Carretero J., 2015, [MNRAS](#), **453**, 1513

DESI Collaboration et al., 2016, [arXiv e-prints](#), p. [arXiv:1611.00036](#)

Dalal R., et al., 2023, [Phys. Rev. D](#), **108**, 123519

Das S., Bode P., 2008, [ApJ](#), **682**, 1

Dawson K. S., et al., 2013, [AJ](#), **145**, 10

DeRose J., et al., 2019, [ApJ](#), **875**, 69

Di Valentino E., et al., 2021, [Classical and Quantum Gravity](#), **38**, 153001

Dietrich J. P., Hartlap J., 2010, [MNRAS](#), **402**, 1049

Dodelson S., 2003, *Modern Cosmology*

Dodelson S., Zhang P., 2005, [Phys. Rev. D](#), **72**, 083001

Dodelson S., Kolb E. W., Matarrese S., Riotto A., Zhang P., 2005, [Phys. Rev. D](#), **72**, 103004

Doux C., et al., 2022, [MNRAS](#), **515**, 1942

Dubinski J., 1996, [New Astronomy](#), **1**, 133

Efstathiou G., Davis M., White S. D. M., Frenk C. S., 1985, [ApJS](#), **57**, 241

Einasto J., 1965, *Trudy Astrofizicheskogo Instituta Alma-Ata*, **5**, 87

Einstein A., 1915, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, pp 844–847

Eisenstein D. J., et al., 2005, [ApJ](#), **633**, 560

Euclid Collaboration et al., 2019, [A&A](#), **627**, A59

Euclid Collaboration et al., 2023, [A&A](#), **675**, A120

Euclid Collaboration et al., 2024, [arXiv e-prints](#), p. arXiv:2405.13495

Fabbian G., Lewis A., Beck D., 2019, *J. Cosmology Astropart. Phys.*, **2019**, 057

Falco E. E., Gorenstein M. V., Shapiro I. I., 1985, [ApJ](#), **289**, L1

Feldman H. A., Kaiser N., Peacock J. A., 1994, [ApJ](#), **426**, 23

Feng Y., Chu M.-Y., Seljak U., McDonald P., 2016, [MNRAS](#), **463**, 2273

Ferlito F., et al., 2023, [MNRAS](#), **524**, 5591

Flaugher B., et al., 2015, [AJ](#), **150**, 150

Fluri J., Kacprzak T., Sgier R., Refregier A., Amara A., 2018, *J. Cosmology Astropart. Phys.*, **2018**, 051

Fluri J., Kacprzak T., Lucchi A., Refregier A., Amara A., Hofmann T., Schneider A., 2019, [Phys. Rev. D](#), **100**, 063514

Fluri J., Kacprzak T., Lucchi A., Schneider A., Refregier A., Hofmann T., 2022, [Phys. Rev. D](#), **105**, 083518

- Fosalba P., Gaztañaga E., Castander F. J., Manera M., 2008, [MNRAS](#), **391**, 435
- Fosalba P., Gaztañaga E., Castander F. J., Crocce M., 2015a, [MNRAS](#), **447**, 1319
- Fosalba P., Crocce M., Gaztañaga E., Castander F. J., 2015b, [MNRAS](#), **448**, 2987
- Freedman W. L., et al., 2001, [ApJ](#), **553**, 47
- Friedmann A., 1922, [Zeitschrift für Physik](#), **10**, 377
- Fu L., et al., 2014, [MNRAS](#), **441**, 2725
- Gamow G., 1948, [Nature](#), **162**, 680
- García-García C., Ruiz-Zapatero J., Alonso D., Bellini E., Ferreira P. G., Mueller E.-M., Nicola A., Ruiz-Lapuente P., 2021, [J. Cosmology Astropart. Phys.](#), **2021**, 030
- Gatti M., et al., 2020, [MNRAS](#), **498**, 4060
- Grewal N., Zuntz J., Tröster T., Amon A., 2022, [The Open Journal of Astrophysics](#), **5**, 13
- Gunn J. E., Gott J. Richard I., 1972, [ApJ](#), **176**, 1
- Guth A. H., 1981, [Phys. Rev. D](#), **23**, 347
- Górski K. M., Hivon E., Banday A. J., Wandelt B. D., Hansen F. K., Reinecke M., Bartelmann M., 2005, [The Astrophysical Journal](#), **622**, 759
- Heitmann K., et al., 2019, [ApJS](#), **245**, 16
- Hernquist L., Katz N., 1989, [ApJS](#), **70**, 419
- Hikage C., et al., 2019, [PASJ](#), **71**, 43
- Hilbert S., Hartlap J., White S. D. M., Schneider P., 2009, [A&A](#), **499**, 31
- Hockney R. W., Eastwood J. W., 1981, *Computer Simulation Using Particles*. CRC Press
- Hogg D. W., 1999, [arXiv e-prints](#), pp astro-ph/9905116
- Hubble E. P., 1925, *The Observatory*, **48**, 139
- Hubble E. P., 1929, [ApJ](#), **69**, 103
- Ivezić Ž., et al., 2019, [ApJ](#), **873**, 111
- Jain B., Van Waerbeke L., 2000, [ApJ](#), **530**, L1
- Jeans J. H., 1902, [Philosophical Transactions of the Royal Society of London Series A](#), **199**, 1

Joachimi B., et al., 2015, [Space Sci. Rev.](#), 193, 1

Kacprzak T., et al., 2016, [MNRAS](#), 463, 3653

Kaiser N., 2000, [ApJ](#), 537, 555

Kaiser N., Squires G., 1993, [ApJ](#), 404, 441

Kiessling A., et al., 2015, [Space Sci. Rev.](#), 193, 67

Kilbinger M., 2015, [Reports on Progress in Physics](#), 78, 086901

Kilbinger M., Schneider P., 2005, [A&A](#), 442, 69

Kirk D., et al., 2015, [Space Sci. Rev.](#), 193, 139

Kollmeier J., et al., 2019, in Bulletin of the American Astronomical Society. p. 274

Kratochvil J. M., Lim E. A., Wang S., Haiman Z., May M., Huffenberger K., 2012, [Phys. Rev. D](#), 85, 103513

LSST Dark Energy Science Collaboration 2012, [arXiv e-prints](#), p. arXiv:1211.0310

LSST Science Collaboration et al., 2009, [arXiv e-prints](#), p. arXiv:0912.0201

Lacasa F., Grain J., 2019, [A&A](#), 624, A61

Lacasa F., Rosenfeld R., 2016, [J. Cosmology Astropart. Phys.](#), 2016, 005

Lacasa F., Lima M., Aguena M., 2018, [A&A](#), 611, A83

Leclercq F., 2020, Evolution of cosmological simulations over the last 50 years, <http://florent-leclercq.eu/blog.php?page=2>

Lemaître G., 1931, [MNRAS](#), 91, 483

Lewis A., Challinor A., Lasenby A., 2000, [ApJ](#), 538, 473

Li Y., Hu W., Takada M., 2014, [Phys. Rev. D](#), 89, 083519

Li X., et al., 2023, [Phys. Rev. D](#), 108, 123518

Limber D. N., 1954, [ApJ](#), 119, 655

Linde A. D., 1982, [Physics Letters B](#), 108, 389

Linke L., Heydenreich S., Burger P. A., Schneider P., 2023, [A&A](#), 672, A185

Liu J., Bird S., Zorrilla Matilla J. M., Hill J. C., Haiman Z., Madhavacheril M. S., Petri A., Spergel D. N., 2018, [J. Cosmology Astropart. Phys.](#), 2018, 049

Loureiro A., et al., 2022, [A&A](#), 665, A56

Madhavacheril M. S., et al., 2024, [ApJ](#), 962, 113

Maksimova N. A., Garrison L. H., Eisenstein D. J., Hadzhiyska B., Bose S., Satterthwaite T. P., 2021, [MNRAS](#), 508, 4017

Mandelbaum R., 2018, [ARA&A](#), 56, 393

Mandelbaum R., Slosar A., Baldauf T., Seljak U., Hirata C. M., Nakajima R., Reyes R., Smith R. E., 2013, [MNRAS](#), 432, 1544

Mandelbaum R., et al., 2018a, [PASJ](#), 70, S25

Mandelbaum R., et al., 2018b, [MNRAS](#), 481, 3170

Marques G. A., Liu J., Zorrilla Matilla J. M., Haiman Z., Bernui A., Novaes C. P., 2019, [J. Cosmology Astropart. Phys.](#), 2019, 019

Marques G. A., et al., 2024, [MNRAS](#), 528, 4513

Martinet N., et al., 2018, [MNRAS](#), 474, 712

Matsubara T., 2010, [Phys. Rev. D](#), 81, 083505

Maturi M., Angrick C., Pace F., Bartelmann M., 2010, [A&A](#), 519, A23

Mead A. J., Heymans C., Lombriser L., Peacock J. A., Steele O. I., Winther H. A., 2016, [MNRAS](#), 459, 1468

Mecke K. R., Buchert T., Wagner H., 1994, [A&A](#), 288, 697

Miyazaki S., et al., 2018, [PASJ](#), 70, S1

Mo H. J., White S. D. M., 1996, [MNRAS](#), 282, 347

Navarro J. F., Frenk C. S., White S. D. M., 1996, [ApJ](#), 462, 563

Navarro J. F., Frenk C. S., White S. D. M., 1997, [ApJ](#), 490, 493

Nelson D., et al., 2019, [Computational Astrophysics and Cosmology](#), 6, 2

Oguri M., Hamana T., 2011, [MNRAS](#), 414, 1851

Omori Y., 2024, [MNRAS](#), 530, 5030

Panamarev T., Just A., Spurzem R., Berczik P., Wang L., Arca Sedda M., 2019, [MNRAS](#), **484**, 3279

Peebles P. J. E., 1968, [ApJ](#), **153**, 1

Peebles P. J. E., Yu J. T., 1970, [ApJ](#), **162**, 815

Penzias A. A., Wilson R. W., 1965, [ApJ](#), **142**, 419

Perlmutter S., et al., 1999, [ApJ](#), **517**, 565

Petri A., 2016, [Astronomy and Computing](#), **17**, 73

Petri A., Liu J., Haiman Z., May M., Hui L., Kratochvil J. M., 2015, [Phys. Rev. D](#), **91**, 103511

Petri A., Haiman Z., May M., 2016, [Phys. Rev. D](#), **93**, 063524

Petri A., Haiman Z., May M., 2017, [Phys. Rev. D](#), **95**, 123503

Planck Collaboration et al., 2014, [A&A](#), **571**, A15

Planck Collaboration et al., 2016, [A&A](#), **594**, A13

Planck Collaboration et al., 2020, [A&A](#), **641**, A6

Powell J., Caudill L., Young O., 2023, [American Journal of Physics](#), **91**, 478

Press W. H., Schechter P., 1974, [ApJ](#), **187**, 425

Rao C. R., 1952, *Advanced Statistical Methods in Biometric Research*. John Wiley & Sons, Inc., New York, N.Y.

Refregier A., Amara A., Kitching T. D., Rassat A., Scaramella R., Weller J., 2010, [arXiv e-prints](#), p. [arXiv:1001.0061](#)

Refregier A., Kacprzak T., Amara A., Bridle S., Rowe B., 2012, [MNRAS](#), **425**, 1951

Riess A. G., et al., 1998, [AJ](#), **116**, 1009

Riess A. G., Casertano S., Yuan W., Macri L. M., Scolnic D., 2019, [ApJ](#), **876**, 85

Riess A. G., et al., 2022, [ApJ](#), **934**, L7

Rubin V. C., Ford W. Kent J., 1970, [ApJ](#), **159**, 379

Rubin V. C., Ford W. K. J., Thonnard N., 1980, [ApJ](#), **238**, 471

Sato M., Hamana T., Takahashi R., Takada M., Yoshida N., Matsubara T., Sugiyama N., 2009, [The Astrophysical Journal](#), **701**, 945

Schaye J., et al., 2015, [MNRAS](#), **446**, 521

Schmalzing J., Buchert T., 1997, [ApJ](#), **482**, L1

Schmalzing J., Kerscher M., Buchert T., 1996, in Bonometto S., Primack J. R., Provenzale A., eds, *Dark Matter in the Universe*. p. 281 ([arXiv:astro-ph/9508154](#)), [doi:10.48550/arXiv.astro-ph/9508154](#)

Schneider P., 2006, in Meylan G., Jetzer P., North P., Schneider P., Kochanek C. S., Wambsganss J., eds, *Saas-Fee Advanced Course 33: Gravitational Lensing: Strong, Weak and Micro*. pp 269–451

Schneider P., Seitz C., 1995, [A&A](#), **294**, 411

Schneider P., Ehlers J., Falco E. E., 1992, *Gravitational Lenses*, [doi:10.1007/978-3-662-03758-4](#).

Scoccimarro R., Couchman H. M. P., Frieman J. A., 1999, [ApJ](#), **517**, 531

Secco L. F., et al., 2022, [Phys. Rev. D](#), **105**, 023515

Sehgal N., Bode P., Das S., Hernandez-Monteagudo C., Huffenberger K., Lin Y.-T., Ostriker J. P., Trac H., 2010, [ApJ](#), **709**, 920

Seljak U. c. v., 2009, [Phys. Rev. Lett.](#), **102**, 021302

Sheldon E. S., Becker M. R., MacCrann N., Jarvis M., 2020, [ApJ](#), **902**, 138

Sheth R. K., Tormen G., 1999, [MNRAS](#), **308**, 119

Sheth R. K., Mo H. J., Tormen G., 2001, [MNRAS](#), **323**, 1

Shirasaki M., Hamana T., Yoshida N., 2015, [MNRAS](#), **453**, 3043

Shirasaki M., Hamana T., Takada M., Takahashi R., Miyatake H., 2019, [MNRAS](#), **486**, 52

Skillman S. W., Warren M. S., Turk M. J., Wechsler R. H., Holz D. E., Sutter P. M., 2014, [arXiv e-prints](#), p. [arXiv:1407.2600](#)

Smith R. E., et al., 2003, [MNRAS](#), **341**, 1311

Smoot G. F., et al., 1992, [ApJ](#), **396**, L1

Spergel D., et al., 2015, [arXiv e-prints](#), p. [arXiv:1503.03757](#)

Springel V., 2005, [MNRAS](#), **364**, 1105

Springel V., et al., 2005, [Nature](#), **435**, 629

Springel V., et al., 2018, [MNRAS](#), **475**, 676

- Starck J.-L., Murtagh F., Fadili J. M., 2010, *Sparse Image and Signal Processing: Wavelets, Curvelets, Morphological Diversity*. Cambridge University Press
- Starck J. L., Themelis K. E., Jeffrey N., Peel A., Lanusse F., 2021, [A&A](#), **649**, A99
- Stein G., Alvarez M. A., Bond J. R., van Engelen A., Battaglia N., 2020, [J. Cosmology Astropart. Phys.](#), **2020**, 012
- Sunyaev R. A., Zeldovich Y. B., 1970, [Ap&SS](#), **7**, 3
- Swinbank R., James Purser R., 2006, [Quarterly Journal of the Royal Meteorological Society](#), **132**, 1769
- Takada M., Hu W., 2013, [Phys. Rev. D](#), **87**, 123504
- Takada M., Jain B., 2004, [MNRAS](#), **348**, 897
- Takahashi R., Sato M., Nishimichi T., Taruya A., Oguri M., 2012, [ApJ](#), **761**, 152
- Takahashi R., Hamana T., Shirasaki M., Namikawa T., Nishimichi T., Osato K., Shiroyama K., 2017, [ApJ](#), **850**, 24
- Tamura N., et al., 2016, in Evans C. J., Simard L., Takami H., eds, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 9908, Ground-based and Airborne Instrumentation for Astronomy VI*. p. 99081M ([arXiv:1608.01075](#)), [doi:10.1117/12.2232103](#)
- Tegmark M., 1997, [Phys. Rev. Lett.](#), **79**, 3806
- Teyssier R., 2002, [A&A](#), **385**, 337
- Teyssier R., et al., 2009, [A&A](#), **497**, 335
- The Dark Energy Survey Collaboration 2005, [arXiv e-prints](#), pp astro-ph/0510346
- The EAGLE team 2017, [arXiv e-prints](#), p. [arXiv:1706.09899](#)
- Thiele L., Hill J. C., Smith K. M., 2020, [Phys. Rev. D](#), **102**, 123545
- Thiele L., Marques G. A., Liu J., Shirasaki M., 2023, [arXiv e-prints](#), p. [arXiv:2304.05928](#)
- Trebitsch M., et al., 2021, [A&A](#), **653**, A154
- Uhlemann C., Friedrich O., Boyle A., Gough A., Barthelemy A., Bernardeau F., Codis S., 2023, [The Open Journal of Astrophysics](#), **6**, 1
- Upadhye A., et al., 2024, [MNRAS](#), **529**, 1862

- Vogelsberger M., et al., 2014, [MNRAS](#), **444**, 1518
- Wang L., Spurzem R., Aarseth S., Nitadori K., Berczik P., Kouwenhoven M. B. N., Naab T., 2015, [MNRAS](#), **450**, 4070
- Weinberg S., 1972, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*
- Weinberg S., 2008, *Cosmology*
- Wick G. C., 1950, [Phys. Rev.](#), **80**, 268
- Wiener N., 1949, *Extrapolation, Interpolation, and Smoothing of Stationary Time Series: With Engineering Applications*. The MIT Press (https://direct.mit.edu/book-pdf/2313079/book_9780262257190.pdf), [doi:10.7551/mitpress/2946.001.0001](https://doi.org/10.7551/mitpress/2946.001.0001), <https://doi.org/10.7551/mitpress/2946.001.0001>
- Wu H.-Y., Weinberg D. H., Salcedo A. N., Wibking B. D., Zu Y., 2019, [MNRAS](#), **490**, 2606
- Zel'dovich Y. B., 1970, [A&A](#), **5**, 84
- Zeldovich Y. B., Sunyaev R. A., 1969, [Ap&SS](#), **4**, 301
- Zhang T., Rau M. M., Mandelbaum R., Li X., Moews B., 2023a, [MNRAS](#), **518**, 709
- Zhang T., et al., 2023b, [MNRAS](#), **525**, 2441
- Zonca A., Singer L., Lenz D., Reinecke M., Rosset C., Hivon E., Gorski K., 2019, [Journal of Open Source Software](#), **4**, 1298
- Zürcher D., Fluri J., Sgier R., Kacprzak T., Refregier A., 2021, [J. Cosmology Astropart. Phys.](#), **2021**, 028
- de Jong J. T. A., et al., 2013, *The Messenger*, **154**, 44