

# Two Experiments

Aharonov-Bohm interferometry and Wilson lines

Sun Ning

Institute for Advance Study, Tsinghua University

February 20, 2016

# Outline

## Aharonov-Bohm interferometry

- Experimental setup and theoretical preparation

- The experiment

## Wilson lines

- Experimental setup and theoretical preparation

- Measuring Wilson lines

- Reconstructing band eigenstates

- Determining Wilson line eigenvalue

- Accessing the dispersion relation

# Outline

## Aharonov-Bohm interferometry

- Experimental setup and theoretical preparation

- The experiment

## Wilson lines

- Experimental setup and theoretical preparation

- Measuring Wilson lines


- Reconstructing band eigenstates

- Determining Wilson line eigenvalues

- Accessing the dispersion relation

# An Aharonov-Bohm interferometer for determining Bloch band topology

# An Aharonov-Bohm interferometer for determining Bloch band topology



# Berry connection and Berry curvature

# Berry connection and Berry curvature

- Bloch state

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$



# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

$$\mathbf{A}_n(\mathbf{k}) = i\langle u_{\mathbf{k}}^n | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^n \rangle$$

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

$$\mathbf{A}_n(\mathbf{k}) = i\langle u_{\mathbf{k}}^n | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^n \rangle$$

- Berry curvature

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

$$\mathbf{A}_n(\mathbf{k}) = i\langle u_{\mathbf{k}}^n | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^n \rangle$$

- Berry curvature

$$\Omega_n = \nabla_{\mathbf{k}} \times \mathbf{A}_n(\mathbf{k})$$

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

$$\mathbf{A}_n(\mathbf{k}) = i\langle u_{\mathbf{k}}^n | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^n \rangle$$

- Berry curvature

$$\Omega_n = \nabla_{\mathbf{k}} \times \mathbf{A}_n(\mathbf{k})$$

- Berry phase

# Berry connection and Berry curvature

- Bloch state

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

- Berry connection

$$\mathbf{A}_n(\mathbf{k}) = i\langle u_{\mathbf{k}}^n | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^n \rangle$$

- Berry curvature

$$\Omega_n = \nabla_{\mathbf{k}} \times \mathbf{A}_n(\mathbf{k})$$


- Berry phase

$$\phi_{\text{Berry}} = \oint_C \mathbf{A}_n(\mathbf{k}) d\mathbf{k} = \int_S \Omega_n(\mathbf{k}) d^2\mathbf{k}$$

# Hexagonal lattice in real space

# Hexagonal lattice in real space

Magnetic field  $B = B_0 + \mathbf{r} \cdot \nabla B$  combined with an orthogonal acceleration  $\mathbf{a} \perp \nabla B$  of the lattice:



The Hamiltonian is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V[\mathbf{r} - \mathbf{R}] - \mu \mathbf{r} \cdot \nabla B - \mu B_0$$

in co-moving frame

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t)$$



# How arbitrary path in reciprocal space comes about

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left( + \frac{1}{2} m |\mathbf{a}t|^2 \right)$$

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left( + \frac{1}{2} m |\mathbf{a}t|^2 \right)$$

external force inducing moving in reciprocal space

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left( + \frac{1}{2} m |\mathbf{a}t|^2 \right)$$

external force inducing moving in reciprocal space

$$\mathcal{F}_\mu = \mu \nabla B - m \mathbf{a}$$

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left(+\frac{1}{2}m|\mathbf{a}t|^2\right)$$

external force inducing moving in reciprocal space

$$\mathcal{F}_\mu = \mu \nabla B - m\mathbf{a}$$

Zeeman energy

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left( + \frac{1}{2} m |\mathbf{a}t|^2 \right)$$

external force inducing moving in reciprocal space

$$\mathcal{F}_\mu = \mu \nabla B - m \mathbf{a}$$

Zeeman energy  $\varepsilon_\mu(t) = -\mu[\mathbf{R}(t) \cdot \nabla B + B_0]$

# How arbitrary path in reciprocal space comes about

The Hamiltonian in co-moving frame is

$$H(t) = \frac{\mathbf{p}^2}{2m} + V(\mathbf{r}) - \mathcal{F}_\mu \cdot \mathbf{r} + \varepsilon_\mu(t) \quad \left(+\frac{1}{2}m|\mathbf{a}t|^2\right)$$

external force inducing moving in reciprocal space

$$\mathcal{F}_\mu = \mu \nabla B - m\mathbf{a}$$

Zeeman energy  $\varepsilon_\mu(t) = -\mu[\mathbf{R}(t) \cdot \nabla B + B_0]$

$$\mathcal{F}_\mu : \quad \mathbf{k} \rightarrow \mathbf{k} + \mathcal{F}_\mu t$$



# Evolution of a state in such a lattice

# Evolution of a state in such a lattice

Hamiltonian

$$H = H_0 - \mathcal{F}_\mu \cdot \mathbf{r}$$

# Evolution of a state in such a lattice

Hamiltonian

$$H = H_0 - \mathcal{F}_\mu \cdot \mathbf{r}$$

Bloch state

$$H_0 \psi_{\mathbf{k}}^n(\mathbf{r}) = E_n(\mathbf{k}) \psi_{\mathbf{k}}^n(\mathbf{r})$$

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

# Evolution of a state in such a lattice

Hamiltonian

$$H = H_0 - \mathcal{F}_\mu \cdot \mathbf{r}$$

Bloch state

$$H_0 \psi_{\mathbf{k}}^n(\mathbf{r}) = E_n(\mathbf{k}) \psi_{\mathbf{k}}^n(\mathbf{r})$$

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

Ansatz

$$\tilde{\Psi}(t) = e^{i\eta(t)} \psi_{k_0 + \mathcal{F}_\mu t}^n$$

$$\eta = \phi_{\text{dyn}} + \phi_{\text{Berry}}$$

# Evolution of a state in such a lattice

## Hamiltonian

$$H = H_0 - \mathcal{F}_\mu \cdot \mathbf{r}$$

## Bloch state

$$H_0 \psi_{\mathbf{k}}^n(\mathbf{r}) = E_n(\mathbf{k}) \psi_{\mathbf{k}}^n(\mathbf{r})$$

$$\psi_{\mathbf{k}}^n(\mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_{\mathbf{k}}^n(\mathbf{r})$$

## Ansatz

$$\tilde{\Psi}(t) = e^{i\eta(t)} \psi_{\mathbf{k}_0 + \mathcal{F}_\mu t}^n$$

$$\eta = \phi_{\text{dyn}} + \phi_{\text{Berry}}$$


## Phase


$$\phi_{\text{dyn}} = \int_0^T [E_1(\mathbf{k} + \mathcal{F}_\mu t) + \varepsilon_\mu t] dt$$

$$\phi_{\text{Berry}} = i \int_C \langle u_{\mathbf{k}}^1 | \nabla_{\mathbf{k}} | u_{\mathbf{k}}^1 \rangle d\mathbf{k}$$


# Experiment procedure


# Experiment procedure




**B**




**B**

**B**


1.  $^{87}\text{Rb}$  BEC initial state  
 $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  
 $\pi/2$ -microwave pulse;

**B**

1.  $^{87}\text{Rb}$  BEC initial state  
 $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  
 $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;

**B**


1.  $^{87}\text{Rb}$  BEC initial state  
 $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  
 $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;

**B**

1.  $^{87}\text{Rb}$  BEC initial state  
 $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  
 $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;
4.  $\tau$  evolution;




1.  $^{87}\text{Rb}$  BEC initial state  
 $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  
 $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;
4.  $\tau$  evolution;
5.  $\pi/2$  pulse;

**B**

1.  $^{87}\text{Rb}$  BEC initial state  $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;
4.  $\tau$  evolution;
5.  $\pi/2$  pulse;

zero-area reference:  
V-shape path.


**B**

1.  $^{87}\text{Rb}$  BEC initial state  $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;
4.  $\tau$  evolution;
5.  $\pi/2$  pulse;

zero-area reference:  
V-shape path.

Reversing the lattice  
acceleration after  $\pi$  pulse.



**B**


1.  $^{87}\text{Rb}$  BEC initial state  $|\uparrow\rangle = |F=2, m_F=1\rangle$ ;  $\pi/2$ -microwave pulse;
2.  $\tau$  evolution;
3. microwave  $\pi$  pulse;
4.  $\tau$  evolution;
5.  $\pi/2$  pulse;

zero-area reference:  
V-shape path.


Reversing the lattice  
acceleration after  $\pi$  pulse.

$$n_{\uparrow,\downarrow} \propto 1 \pm \cos(\varphi + \varphi_{MW})$$

# Main result I




# Main result I



- Broadening of the edges — caused by momentum spread.


## Main result I



- Broadening of the edges — caused by momentum spread.
- Systematic errors.


# Self-referenced interferometry at Dirac point

# Self-referenced interferometry at Dirac point




## Main result II

B



## Main result II

B




- Contrast:  $(n_{\downarrow}^{\text{max}} - n_{\downarrow}^{\text{min}}) / (n_{\downarrow}^{\text{max}} + n_{\downarrow}^{\text{min}})$



## Main result II

B




- Contrast:  $(n_{\downarrow}^{\text{max}} - n_{\downarrow}^{\text{min}}) / (n_{\downarrow}^{\text{max}} + n_{\downarrow}^{\text{min}})$
- $\varphi = (\varphi_L + \varphi_R)/2 - \varphi_B$



# Main result II


**B**



- Contrast:  $(n_{\downarrow}^{\text{max}} - n_{\downarrow}^{\text{min}}) / (n_{\downarrow}^{\text{max}} + n_{\downarrow}^{\text{min}})$
- $\varphi = (\varphi_L + \varphi_R) / 2 - \varphi_B = 0.95(10)\pi$
- Berry curvature localization

## Main result II


B



- Contrast:  $(n_{\downarrow}^{\text{max}} - n_{\downarrow}^{\text{min}}) / (n_{\downarrow}^{\text{max}} + n_{\downarrow}^{\text{min}})$
- $\varphi = (\varphi_L + \varphi_R) / 2 - \varphi_B = 0.95(10)\pi$
- Berry curvature localization  $\delta k_w \simeq 10^{-4} k_L (\Delta \simeq h \times 3\text{Hz})$

# Imbalance lattice mapping

# Imbalance lattice mapping




# Imbalance lattice mapping

## Self-referenced phase

# Imbalance lattice mapping


## Self-referenced phase






# *ab initio* calculation of imbalanced lattice

# *ab initio* calculation of imbalanced lattice



# *ab initio* calculation of imbalanced lattice



Seeing Dirac points annihilating clear.

# Outline

## Aharonov-Bohm interferometry

Experimental setup and theoretical preparation

The experiment

## Wilson lines

Experimental setup and theoretical preparation


Measuring Wilson lines

Reconstructing band eigenstates

Determining Wilson line eigenvalues



Accessing the dispersion relation

# Experimental reconstruction of Wilson lines in Bloch bands




# Wilson line regime in honeycomb lattice

# Wilson line regime in honeycomb lattice




# Wilson line regime in honeycomb lattice



- force  $\mathbf{F}$  :  $\mathbf{q}(t) = \mathbf{q}(0) + \mathbf{F}t$



# Wilson line regime in honeycomb lattice



- force  $\mathbf{F}$  :  $\mathbf{q}(t) = \mathbf{q}(0) + \mathbf{F}t$
- unitary time-evolution operator (Wilson line matrix)

$$W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp(i \int_C A_{\mathbf{q}} d\mathbf{q})$$

$A_{\mathbf{q}}$ : Wilczek-Zee connection

$\mathcal{P}$ : Path-ordering (non-Abelian)

# Wilson line

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$
- Wilczek-Zee connection

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$
- Wilczek-Zee connection  $A_{\mathbf{q}}^{n,n'} = i \langle u_{\mathbf{q}}^n | \nabla_{\mathbf{q}} | u_{\mathbf{q}}^{n'} \rangle$

# Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$
- Wilczek-Zee connection  $A_{\mathbf{q}}^{n,n'} = i \langle u_{\mathbf{q}}^n | \nabla_{\mathbf{q}} | u_{\mathbf{q}}^{n'} \rangle$
- $n = n'$ : Berry connections of individual Bloch bands





## Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$
- Wilczek-Zee connection  $A_{\mathbf{q}}^{n,n'} = i \langle u_{\mathbf{q}}^n | \nabla_{\mathbf{q}} | u_{\mathbf{q}}^{n'} \rangle$
- $n = n'$ : Berry connections of individual Bloch bands yields Berry phase along a closed path
- $n \neq n'$ : inter-band Berry connections

## Wilson line

- $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$
- Bloch state (presence of lattice)  $|\Phi_{\mathbf{q}}^n\rangle = e^{i\mathbf{q} \cdot \mathbf{r}} |u_{\mathbf{q}}^n\rangle$
- Wilczek-Zee connection  $A_{\mathbf{q}}^{n,n'} = i \langle u_{\mathbf{q}}^n | \nabla_{\mathbf{q}} | u_{\mathbf{q}}^{n'} \rangle$
- $n = n'$ : Berry connections of individual Bloch bands yields Berry phase along a closed path
- $n \neq n'$ : inter-band Berry connections induce inter-band transition

# Dynamics in combined lattice with external force

- total Hamiltonian:

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$
- lattice

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$
- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$
- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force



# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$
- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$

- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

- initial state

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$

- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

- initial state  $|\psi(0)\rangle = \sum_n \alpha^n(0) |\Phi_{\mathbf{q}_0}^n\rangle$

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$
- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

- initial state  $|\psi(0)\rangle = \sum_n \alpha^n(0) |\Phi_{\mathbf{q}_0}^n\rangle$   
 $|\alpha^n(0)|^2$  gives the population in the  $n^{\text{th}}$  band at  $t = 0$

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$

- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

- initial state  $|\psi(0)\rangle = \sum_n \alpha^n(0) |\Phi_{\mathbf{q}_0}^n\rangle$   
 $|\alpha^n(0)|^2$  gives the population in the  $n^{\text{th}}$  band at  $t = 0$
- ansatz

# Dynamics in combined lattice with external force

- total Hamiltonian:  $H = H_0 - \mathbf{F} \cdot \mathbf{r}$

- lattice

$$H_0 = \sum_{\mathbf{q}, n} E_{\mathbf{q}}^n |\Phi_{\mathbf{q}}^n\rangle \langle \Phi_{\mathbf{q}}^n|$$

constant force  $\mathbf{F}$

- initial state  $|\psi(0)\rangle = \sum_n \alpha^n(0) |\Phi_{\mathbf{q}_0}^n\rangle$   
 $|\alpha^n(0)|^2$  gives the population in the  $n^{\text{th}}$  band at  $t = 0$
- ansatz

$$|\psi(t)\rangle = \sum_n \alpha^n(t) |\Phi_{\mathbf{q}(t)}^n\rangle$$

$$\mathbf{q}(t) = \mathbf{q}_0 + \mathbf{F}t$$

# Wilson lines and Wilczek-Zee connections

# Wilson lines and Wilczek-Zee connections

## Example (two-band system)



# Wilson lines and Wilczek-Zee connections

## Example (two-band system)

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} E_{\mathbf{q}(t)}^1 - \tilde{\zeta}_{\mathbf{q}(t)}^{1,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{1,2} \\ -\tilde{\zeta}_{\mathbf{q}(t)}^{2,1} & E_{\mathbf{q}(t)}^1 - \tilde{\zeta}_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

# Wilson lines and Wilczek-Zee connections

## Example (two-band system)

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} E_{\mathbf{q}(t)}^1 - \tilde{\zeta}_{\mathbf{q}(t)}^{1,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{1,2} \\ -\tilde{\zeta}_{\mathbf{q}(t)}^{2,1} & E_{\mathbf{q}(t)}^1 - \tilde{\zeta}_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

where

# Wilson lines and Wilczek-Zee connections

## Example (two-band system)

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{1,1} & -\zeta_{\mathbf{q}(t)}^{1,2} \\ -\zeta_{\mathbf{q}(t)}^{2,1} & E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

where

$$\zeta_{\mathbf{q}(t)}^{n,n'} = A_{\mathbf{q}(t)}^{n,n'} \cdot \mathbf{F} = i\langle u_{\mathbf{q}(t)}^n | \partial_t | u_{\mathbf{q}(t)}^{n'} \rangle$$

# Wilson lines and Wilczek-Zee connections

## Example (two-band system)

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{1,1} & -\zeta_{\mathbf{q}(t)}^{1,2} \\ -\zeta_{\mathbf{q}(t)}^{2,1} & E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

where

$$\zeta_{\mathbf{q}(t)}^{n,n'} = A_{\mathbf{q}(t)}^{n,n'} \cdot \mathbf{F} = i\langle u_{\mathbf{q}(t)}^n | \partial_t | u_{\mathbf{q}(t)}^{n'} \rangle$$

and thus

# Wilson lines and Wilczek-Zee connections

## Example (two-band system)

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{1,1} & -\zeta_{\mathbf{q}(t)}^{1,2} \\ -\zeta_{\mathbf{q}(t)}^{2,1} & E_{\mathbf{q}(t)}^1 - \zeta_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

where

$$\zeta_{\mathbf{q}(t)}^{n,n'} = A_{\mathbf{q}(t)}^{n,n'} \cdot \mathbf{F} = i\langle u_{\mathbf{q}(t)}^n | \partial_t | u_{\mathbf{q}(t)}^{n'} \rangle$$

and thus

$$A_{\mathbf{q}(t)}^{n,n'} = i\langle u_{\mathbf{q}}^n | \nabla_{\mathbf{q}} | u_{\mathbf{q}}^{n'} \rangle \Big|_{\mathbf{q}=\mathbf{q}(t)}$$

# Wilson line regime of *the* two-band model

# Wilson line regime of *the* two-band model

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} -\xi_{\mathbf{q}(t)}^{1,1} & -\xi_{\mathbf{q}(t)}^{1,2} \\ -\xi_{\mathbf{q}(t)}^{2,1} & -\xi_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

# Wilson line regime of *the* two-band model

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} -\xi_{\mathbf{q}(t)}^{1,1} & -\xi_{\mathbf{q}(t)}^{1,2} \\ -\xi_{\mathbf{q}(t)}^{2,1} & -\xi_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

- Defining  $\xi_{\mathbf{q}(t)}$  as the matrix with elements  $\xi_{\mathbf{q}(t)}^{n,n'}$



# Wilson line regime of *the* two-band model

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} -\tilde{\zeta}_{\mathbf{q}(t)}^{1,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{1,2} \\ -\tilde{\zeta}_{\mathbf{q}(t)}^{2,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

- Defining  $\tilde{\zeta}_{\mathbf{q}(t)}$  as the matrix with elements  $\tilde{\zeta}_{\mathbf{q}(t)}^{n,n'}$
- state evolution

# Wilson line regime of *the* two-band model

$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} -\tilde{\zeta}_{\mathbf{q}(t)}^{1,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{1,2} \\ -\tilde{\zeta}_{\mathbf{q}(t)}^{2,1} & -\tilde{\zeta}_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$

- Defining  $\tilde{\zeta}_{\mathbf{q}(t)}$  as the matrix with elements  $\tilde{\zeta}_{\mathbf{q}(t)}^{n,n'}$
- state evolution

$$|\psi(t)\rangle = \mathcal{T} \exp(i \int \tilde{\zeta}_{\mathbf{q}(t)} dt) |\psi(0)\rangle \equiv W |\psi(0)\rangle$$

# Wilson line regime of *the* two-band model


$$i\partial_t \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix} = \begin{pmatrix} -\xi_{\mathbf{q}(t)}^{1,1} & -\xi_{\mathbf{q}(t)}^{1,2} \\ -\xi_{\mathbf{q}(t)}^{2,1} & -\xi_{\mathbf{q}(t)}^{2,2} \end{pmatrix} \begin{pmatrix} \alpha^1(t) \\ \alpha^2(t) \end{pmatrix}$$


- Defining  $\xi_{\mathbf{q}(t)}$  as the matrix with elements  $\xi_{\mathbf{q}(t)}^{n,n'}$
- state evolution


$$|\psi(t)\rangle = \mathcal{T} \exp(i \int \xi_{\mathbf{q}(t)} dt) |\psi(0)\rangle \equiv W |\psi(0)\rangle$$

- thus  $W_{\mathbf{q}(0) \rightarrow \mathbf{q}(t)} = \mathcal{P} \exp \left( i \int_C A_{\mathbf{q}} d\mathbf{q} \right)$


# Measuring Wilson lines








- Matrix elements of Wilson line operator



- Matrix elements of Wilson line operator

$$W_{\mathbf{Q} \rightarrow \mathbf{q}}^{m,n} = \langle \Phi_{\mathbf{q}}^m | e^{i(\mathbf{q}-\mathbf{Q}) \cdot \hat{\mathbf{r}}} | \Phi_{\mathbf{Q}}^n \rangle = \langle u_{\mathbf{q}}^m | u_{\mathbf{Q}}^n \rangle$$



- Matrix elements of Wilson line operator

$$W_{\mathbf{Q} \rightarrow \mathbf{q}}^{m,n} = \langle \Phi_{\mathbf{q}}^m | e^{i(\mathbf{q}-\mathbf{Q}) \cdot \hat{\mathbf{r}}} | \Phi_{\mathbf{Q}}^n \rangle = \langle u_{\mathbf{q}}^m | u_{\mathbf{Q}}^n \rangle$$

- saturation value  $W_{\Gamma \rightarrow \mathbf{q}}^{11} = \langle u_{\mathbf{q}}^1 | u_{\Gamma}^1 \rangle$  of population after transport measures overlap between  $|u\rangle$



# Reconstructing band eigenstates

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin
- cell-periodic Bloch functions at a fixed reference quasimomentum  $Q$

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin
- cell-periodic Bloch functions at a fixed reference quasimomentum  $\mathbf{Q}$

$$|1\rangle = |u_{\mathbf{Q}}^1\rangle \quad |2\rangle = |u_{\mathbf{Q}}^2\rangle$$

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin
- cell-periodic Bloch functions at a fixed reference quasimomentum  $\mathbf{Q}$

$$|1\rangle = |u_{\mathbf{Q}}^1\rangle \quad |2\rangle = |u_{\mathbf{Q}}^2\rangle$$

- such that  $|u_{\mathbf{q}}^1\rangle = \cos(\frac{\theta_{\mathbf{q}}}{2})|1\rangle + \sin(\frac{\theta_{\mathbf{q}}}{2})e^{i\phi_{\mathbf{q}}}|2\rangle$

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin
- cell-periodic Bloch functions at a fixed reference quasimomentum  $\mathbf{Q}$

$$|1\rangle = |u_{\mathbf{Q}}^1\rangle \quad |2\rangle = |u_{\mathbf{Q}}^2\rangle$$

- such that  $|u_{\mathbf{q}}^1\rangle = \cos(\frac{\theta_{\mathbf{q}}}{2})|1\rangle + \sin(\frac{\theta_{\mathbf{q}}}{2})e^{i\phi_{\mathbf{q}}}|2\rangle$
- thus  $(\theta_{\mathbf{q}}, \phi_{\mathbf{q}})$  characterize a state

# Reconstructing band eigenstates

- cell-periodic Bloch state as pseudo-spin
- cell-periodic Bloch functions at a fixed reference quasimomentum  $\mathbf{Q}$


$$|1\rangle = |u_{\mathbf{Q}}^1\rangle \quad |2\rangle = |u_{\mathbf{Q}}^2\rangle$$

- such that  $|u_{\mathbf{q}}^1\rangle = \cos(\frac{\theta_{\mathbf{q}}}{2})|1\rangle + \sin(\frac{\theta_{\mathbf{q}}}{2})e^{i\phi_{\mathbf{q}}}|2\rangle$
- thus  $(\theta_{\mathbf{q}}, \phi_{\mathbf{q}})$  characterize a state
- Throughout this work basis states are chosen at reference point  $\mathbf{Q} = \Gamma$



# Measuring mixing angle $\theta_q$



# Measuring mixing angle $\theta_q$




# Measuring mixing angle $\theta_q$





# Measuring relative phase $\phi_q$


# Measuring relative phase $\phi_q$




# Measuring relative phase $\phi_q$




**b**



**b**




**b**



- AB-site degeneracy (blue) —  $\pi$  jump




**b**



- AB-site degeneracy (blue) —  $\pi$  jump
- AB-site offset (red) by elliptically-polarized lattice beam — continuous varying phase

**b**



- AB-site degeneracy (blue) —  $\pi$  jump
- AB-site offset (red) by elliptically-polarized lattice beam — continuous varying phase
- 3-fold symmetry of system (no matter whether offset)

# Determining Wilson line eigenvalue

# Determining Wilson line eigenvalue

## Wilson-Zak loop

$$W_{\mathbf{q} \rightarrow \mathbf{q} + \mathbf{G}}$$

# Determining Wilson line eigenvalue

## Wilson-Zak loop

$$W_{\mathbf{q} \rightarrow \mathbf{q} + \mathbf{G}}$$

- $\hat{A} \implies \hat{A}_{U(1)} + \hat{A}_{SU(2)}$

# Determining Wilson line eigenvalue

## Wilson-Zak loop

$$W_{\mathbf{q} \rightarrow \mathbf{q} + \mathbf{G}}$$

- $\hat{A} \implies \hat{A}_{U(1)} + \hat{A}_{SU(2)}$
- the  $U(2)$  Wilson line  $\implies U(1)$  global phase multiplied by a  $SU(2)$  matrix.

## Wilson-Zak loop


$$W_{q \rightarrow q+G}$$

- $\hat{A} \implies \hat{A}_{U(1)} + \hat{A}_{SU(2)}$
- the  $U(2)$  Wilson line  $\implies U(1)$  global phase multiplied by a  $SU(2)$  matrix.
- $SU(2)$  matrix eigenvalues  $e^{\pm i\zeta}$






# Mapping dispersion relation



# Mapping dispersion relation



by varying the reference quasimomentum  $Q$

- ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡

# Thank you !