

question

2 views

Daily Challenge 20.7

Use the Weierstrass substitution $t = \tan\left(\frac{x}{2}\right)$ to compute

$$\int \frac{dx}{a \sin(x) + b \cos(x)}.$$

You may like to refer to the slides or Wikipedia.

daily_challenge

Updated 4 months ago by Christian Ferko

the students' answer, where students collectively construct a single answer

We let $t = \tan\left(\frac{x}{2}\right)$; we then see that we can substitute out our sine and cosine,

$$\begin{aligned} \int \frac{dx}{a \sin(x) + b \cos(x)} &= \int \frac{dx}{a \frac{2t}{1+t^2} + b \frac{1-t^2}{1+t^2}} \\ &= \int \frac{(1+t^2) dx}{a2t + b(1-t^2)} \end{aligned}$$

And then the $dx = \frac{2}{1+t^2}$:

$$\begin{aligned} \int \frac{(1+t^2) dx}{a2t + b(1-t^2)} &= 2 \int \frac{dt}{a2t + b(1-t^2)} \\ &= 2 \int \frac{dt}{-bt^2 + a2t + b} \end{aligned}$$

Let me just start faffing about with certain chain rules of $\log(-bt^2 + a2t + b)$ and see what we have to divide out. Okay, that didn't work because of the presence of the t^2 , so we have to get creative. There is no way to apply partial fractions to this, as intuitively the polynomial cannot be factored. Let's try writing this in a slightly more obvious way;

$$2 \int \frac{dt}{-bt^2 + a2t + b} = 2 \int (-bt^2 + a2t + b)^{-1} dt$$

How about we try a u-sub where $u = t^2$ and thus $\sqrt{u} = t$, then $du = 2tdt$. This complicates what we have going on by even more, where we can rewrite our problem as

$$2 \int (-bt^2 + a2t + b)^{-1} dt = 2 \int ((2t) \cdot (-bu + a\sqrt{u} + b))^{-1} dt$$

This didn't help and I still don't have any better ideas or alternate u-sub. We cannot apply integration by parts here, as no two functions are multiplied. False, it just doesn't help. Let's go back to trying to factor the denominator. We shall apply the wikipedia article on completing the square. We then see that:

$$2 \int \frac{dt}{-bt^2 + a2t + b} = 2 \int \frac{dt}{-b(t - \frac{a}{b})^2 + b + 2\frac{a}{b}}$$

We can now factor a $b + 2\frac{a}{b}$ out of the denominator and thus totally out to see that

$$2 \int \frac{dt}{-b(t - \frac{a}{b})^2 + b + 2\frac{a}{b}} = \frac{2}{b + 2\frac{a}{b}} \int \frac{dt}{\frac{-b}{b + 2\frac{a}{b}}(t - \frac{a}{b})^2 + 1}$$

And apply the hinted formula for $\int \frac{dx}{1+x^2} = \arctan(x)$

To conclude that

$$\frac{2}{b + 2\frac{a}{b}} \int \frac{dt}{\frac{-b}{b + 2\frac{a}{b}}(t - \frac{a}{b})^2 + 1} = \frac{2}{\sqrt{-b^2 + 2a}} \cdot \arctan\left(\sqrt{\frac{-b}{b + 2\frac{a}{b}}} t\right) + C$$

where then $t = \tan \frac{x}{2}$ and we conclude that

$$\int \frac{dx}{a \sin(x) + b \cos(x)} = \frac{2}{\sqrt{-b^2 + 2a}} \cdot \arctan\left(\sqrt{\frac{-b}{b + 2\frac{a}{b}}} \tan \frac{x}{2}\right) + C$$

Updated 2 months ago by Logan Pachulski

the instructors' answer, where instructors collectively construct a single answer

Click to start off the wiki answer

followup discussions for lingering questions and comments

☒ Resolved ☐ Unresolved



Christian Ferko 2 months ago

Hint: complete the square in the denominator and use that $\int \frac{dx}{1+x^2} = \arctan(x)$.



Logan Pachulski 2 months ago :logwow: