28,6

(a)! Apply the limit comparison test's evaluate by comparison to in lim h sin (in) = lim sin (in)

Since this goer to oras we can apply L'Holpitals rules

willim cos(in):-1: in= = lim (coo(in)) = 1

Since the hour manic sension is divergent, then

Since the hour manic sension is divergent, then

Since the hour manic sension is divergent, then

Since the hour manic sension is divergent.

(b): We see that

at $\sum a_n = b_k - b_0$ Then by taking the limit of each side and applying $bk \rightarrow 0$, $\sum_{n=1}^{\infty} a_n = 0 - b_0 = -b_0$

Mon, solve for A and Bin

A B - 1

A Th

A

Then, we must compute $a_n = \frac{1}{h} - \frac{1}{4n\pi l}$, we must compute $a_n = b_n - b_n$

(c) We see that since (2) and (3) (anverge individually

(1) $\frac{\sum_{k=1}^{\infty} 2^{k+3}}{\sum_{k=1}^{\infty}} = \sum_{k=1}^{\infty} (\frac{2}{5})^{k} + \sum_{k=1}^{\infty} (\frac{2}{5})^{k}$ then apply the sum of the generalic series for $r^{\frac{1}{2}}$, to see that

 $\sum_{h=1}^{\infty} r^{h} = \frac{r}{1-r} + \frac{r}{1-r$

 $\frac{2}{3} + \frac{3}{7}$

 $=\frac{4}{8}+\frac{9}{6}$

- 13 D