

On the Category Theoretic Implications on Induced Representation of Finite Groups

Part 1: Categories

Atonu Roy Chowdhury

BRAC University

atonuroychowdhury@gmail.com

September 19, 2024


What even is Category Theory?

What even is Category Theory?

Is it a branch of math developoed by cats?


What even is Category Theory?

Is it a branch of math developoed by cats?



What even is Category Theory?

Is it a branch of math developoed by cats?




Sadly, no! :(

What do we do in math?

What do we do in math?

In Linear Algebra


we study vector spaces and linear transformations between them



What do we do in math?

In Group Theory


we study groups and group homomorphisms between them



What do we do in math?

In Topology

we study topological spaces and continuous functions between them



What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- We can compose them:

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- We can compose them:
 - Composition of group homomorphisms is a group homomorphism:

$$\begin{array}{ccccc} G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & G_3 \\ & \searrow & & \nearrow & \\ & & f_2 \circ f_1 & & \end{array}$$

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- We can compose them:
 - Composition of group homomorphisms is a group homomorphism:

$$\begin{array}{ccccc} G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & G_3 \\ & \searrow & & \nearrow & \\ & & f_2 \circ f_1 & & \end{array}$$

- Composition of linear maps is a linear map:

$$\begin{array}{ccccc} U & \xrightarrow{f_1} & V & \xrightarrow{f_2} & W \\ & \searrow & & \nearrow & \\ & & f_2 \circ f_1 & & \end{array}$$

- and so on ...

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- There is an identity of the composition!!

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- There is an identity of the composition!!
 - Every group has an identity group homomorphism: id_G

$$\text{id}_G \left(\text{loop} \right) G \xrightarrow{f} H \left(\text{loop} \right) \text{id}_H$$

$$f = f \circ \text{id}_G = \text{id}_H \circ f.$$

What do we do in math?

Essentially, we study structures (i.e. groups, topological spaces, vector spaces) and maps between them.

What properties do these maps have?

- There is an identity of the composition!!
 - Every group has an identity group homomorphism: id_G

$$\text{id}_G \left(\text{loop} \right) G \xrightarrow{f} H \left(\text{loop} \right) \text{id}_H$$

$$f = f \circ \text{id}_G = \text{id}_H \circ f.$$

- Every vector space has an identity transformation: $\mathbb{1}_V$

$$\mathbb{1}_V \left(\text{loop} \right) V \xrightarrow{f} W \left(\text{loop} \right) \mathbb{1}_W$$

$$f = f \circ \mathbb{1}_V = \mathbb{1}_W \circ f.$$

- and so on ...

Now we strip down all the details, and we have Category Theory!

Now we strip down all the details, and we have Category Theory!

Category theory is the bird's eye view of mathematics.

— Tom Leinster

Definition 1

A **category** \mathcal{C} consists of

- a collection of **objects**, often denoted as \mathcal{C}_0 ;
- a collection **arrows** from one object to another, often denoted as \mathcal{C}_1 ; if f is an arrow from A to B , we write $f : A \rightarrow B$ and call $A = \text{dom } f$ and $B = \text{cod } f$; $\text{Hom}_{\mathcal{C}}(A, B)$ is the collection of all arrows from the object A to the object B .

such that

- given any two arrows f, g with $\text{cod } f = \text{dom } g$, there is a composition $g \circ f : \text{dom } f \rightarrow \text{cod } g$;
- for every $A \in \mathcal{C}_0$, there is a unique identity arrow $1_A \in \text{Hom}_{\mathcal{C}}(A, A)$.

These data have the following properties:

- composition is associative: given $f : A \rightarrow B, g : B \rightarrow C, h : C \rightarrow D$, $h \circ (g \circ f) = (h \circ g) \circ f$.
- identity arrow is the identity of composition: for any $f : A \rightarrow B$, $1_B \circ f = f = f \circ 1_A$.

Caution!

Objects need not be sets, and arrows need not be functions!

Examples of Categories

Example 1

Any structured sets (groups, vector spaces, topological spaces, etc) and structure preserving maps between them. For instance,

- ① **Groups** is the category of all groups. **Groups**₀ is the collection of all groups, **Groups**₁ is the collection of all group homomorphisms.
- ② **Vect**_ℂ is the category of all vector spaces over the field ℂ. (**Vect**_ℂ)₀ is the collection of all vector spaces over ℂ, (**Vect**_ℂ)₁ is the collection of all linear maps between them.
- ③ **Sets** is the category of all sets. **Sets**₀ is the collection of all sets, **Sets**₁ is the collection of all functions between sets.
- ④ and so on . . .

Examples of Categories

Example 2

A group itself is also a category!!!

Examples of Categories

Example 2

A group itself is also a category!!!

Let G be a group. Then G can be considered a category, $\mathcal{C}(G)$.


Examples of Categories

Example 2

A group itself is also a category!!!

Let G be a group. Then G can be considered a category, $\mathcal{C}(G)$.

$\mathcal{C}(G)_0 = \{G\}$, $\mathcal{C}(G) = G$. In other words, there is only one object, G itself. The arrows are the elements of G .




Examples of Categories

Example 2

A group itself is also a category!!!

Let G be a group. Then G can be considered a category, $\mathcal{C}(G)$.

$\mathcal{C}(G)_0 = \{G\}$, $\mathcal{C}(G) = G$. In other words, there is only one object, G itself. The arrows are the elements of G .



The identity arrow $\mathbb{1}_G$ is the identity element e of G . The composition of arrows is given by the group operation: $g_1 \circ g_2 = g_1 g_2$.

Category of Categories

So, a group is a category. Then **Groups** is a category of some categories.
Is there a category of all categories? What are the arrows in that category?

Category of Categories

So, a group is a category. Then **Groups** is a category of some categories.

Is there a category of all categories? What are the arrows in that category?

Suppose \mathcal{C} and \mathcal{D} are two categories. Then an arrow $F : \mathcal{C} \rightarrow \mathcal{D}$ is supposed to preserve the “structure” of the categories.

Category of Categories

So, a group is a category. Then **Groups** is a category of some categories.

Is there a category of all categories? What are the arrows in that category?

Suppose \mathcal{C} and \mathcal{D} are two categories. Then an arrow $F : \mathcal{C} \rightarrow \mathcal{D}$ is supposed to preserve the “structure” of the categories.

The only structure on the categories we have are composition of arrows and the identity arrow.

Definition 2

A **functor** $F : \mathcal{C} \rightarrow \mathcal{D}$ between categories \mathcal{C} and \mathcal{D} is two mappings $F_0 : \mathcal{C}_0 \rightarrow \mathcal{D}_0$, and $F_1 : \mathcal{C}_1 \rightarrow \mathcal{D}_1$ such that

- if $f : A \rightarrow B$ is an arrow in \mathcal{C} , then $F_1(f) : F_0(A) \rightarrow F_0(B)$ in \mathcal{D} , i.e. F preserves domains and codomains;
- for every $A \in \mathcal{C}_0$, $F_1(1_A) = 1_{F_0(A)}$;
- given two composable arrows $f : A \rightarrow B$ and $g : B \rightarrow C$ in \mathcal{C} ,

$$F_1(g \circ f) = F_1(g) \circ F_1(f).$$

Category of Categories

Definition 2


A **functor** $F : \mathcal{C} \rightarrow \mathcal{D}$ between categories \mathcal{C} and \mathcal{D} is two mappings $F_0 : \mathcal{C}_0 \rightarrow \mathcal{D}_0$, and $F_1 : \mathcal{C}_1 \rightarrow \mathcal{D}_1$ such that

- if $f : A \rightarrow B$ is an arrow in \mathcal{C} , then $F_1(f) : F_0(A) \rightarrow F_0(B)$ in \mathcal{D} , i.e. F preserves domains and codomains;
- for every $A \in \mathcal{C}_0$, $F_1(1_A) = 1_{F_0(A)}$;
- given two composable arrows $f : A \rightarrow B$ and $g : B \rightarrow C$ in \mathcal{C} ,

$$F_1(g \circ f) = F_1(g) \circ F_1(f).$$

We shall often abuse the notation by writing $F(f) : F(A) \rightarrow F(B)$.

Category of Categories



Category of Categories

In order to make a category of categories, we need identity functors $\mathbb{1}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$. This is easy, we can just map everything to itself.

Category of Categories

In order to make a category of categories, we need identity functors $\mathbb{1}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$. This is easy, we can just map everything to itself.

What about composition?

$$\mathcal{C} \xrightarrow{F} \mathcal{D} \xrightarrow{G} \mathcal{E}$$

Given $A \in \mathcal{C}_0$, we can just define $(G \circ F)_0(A) = G_0(F_0(A))$. For an arrow $f : A \rightarrow B$ in \mathcal{C} , we can define $(G \circ F)_1(f) = G_1(F_1(f))$.

Category of Categories

In order to make a category of categories, we need identity functors $\mathbb{1}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$. This is easy, we can just map everything to itself.

What about composition?

$$\mathcal{C} \xrightarrow{F} \mathcal{D} \xrightarrow{G} \mathcal{E}$$


Given $A \in \mathcal{C}_0$, we can just define $(G \circ F)_0(A) = G_0(F_0(A))$. For an arrow $f : A \rightarrow B$ in \mathcal{C} , we can define $(G \circ F)_1(f) = G_1(F_1(f))$.

Then we have the category of all categories, **Cat**.

Making New categories From Old Ones

Example 3

The opposite category \mathcal{C}^{op} has the same objects as \mathcal{C} , but the arrows are reversed.



Often we use this interpretation that an arrow $f : X \rightarrow Y$ in \mathcal{C}^{op} is really an arrow $f : Y \rightarrow X$ in \mathcal{C} . This is an abuse of notation since we are dropping the superscript op from the arrows in \mathcal{C}^{op} .

Making New categories From Old Ones

Example 4

The product of 2 categories \mathcal{C} and \mathcal{D} , written as $\mathcal{C} \times \mathcal{D}$.

- the objects are (C, D) for $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,
- and the arrows are

$$(f, g) : (C, D) \rightarrow (C', D'),$$

where $f : C \rightarrow C'$ and $g : D \rightarrow D'$ are arrows in \mathcal{C} and \mathcal{D} , respectively.

Making New categories From Old Ones

Example 4

The product of 2 categories \mathcal{C} and \mathcal{D} , written as $\mathcal{C} \times \mathcal{D}$.

- the objects are (C, D) for $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,
- and the arrows are

$$(f, g) : (C, D) \rightarrow (C', D'),$$

where $f : C \rightarrow C'$ and $g : D \rightarrow D'$ are arrows in \mathcal{C} and \mathcal{D} , respectively.

$$\begin{array}{ccc} \mathbb{1}_{\mathcal{C}} \circlearrowleft C & \xrightarrow{f} C' \xrightarrow{f'} C'' & \mathbb{1}_{\mathcal{D}} \circlearrowleft D \xrightarrow{g} D' \xrightarrow{g'} D'' \\ & \searrow f' \circ f \nearrow & \searrow g' \circ g \nearrow \\ \mathbb{1}_{(C,D)} = (\mathbb{1}_C, \mathbb{1}_D) \circlearrowleft (C, D) & \xrightarrow{(f,g)} (C', D') \xrightarrow{(f',g')} (C'', D'') & \\ & \searrow (f',g') \circ (f,g) = (f' \circ f, g' \circ g) \nearrow & \end{array}$$

Making New categories From Old Ones

Example 5

We can also form a category with the arrows of a category \mathcal{C} . It is known as the **arrow category** of \mathcal{C} , and is denoted as $\text{Arr}(\mathcal{C})$. The objects are arrows of \mathcal{C} . What are the arrows of $\text{Arr}(\mathcal{C})$?

Making New categories From Old Ones

Example 5

We can also form a category with the arrows of a category \mathcal{C} . It is known as the **arrow category** of \mathcal{C} , and is denoted as $\text{Arr}(\mathcal{C})$. The objects are arrows of \mathcal{C} . What are the arrows of $\text{Arr}(\mathcal{C})$?

Suppose $f : A \rightarrow B, g : C \rightarrow D \in \text{Arr}(\mathcal{C})_0$. An arrow $x : f \rightarrow g$ in the arrow category is a pair of arrows $x_1 : A \rightarrow C$ and $x_2 : B \rightarrow D$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{x_1} & C \\ f \downarrow & & \downarrow g \\ B & \xrightarrow{x_2} & D \end{array} \quad , \text{ i.e. } x_2 \circ f = g \circ x_1.$$

Making New categories From Old Ones

Example 5

We can also form a category with the arrows of a category \mathcal{C} . It is known as the **arrow category** of \mathcal{C} , and is denoted as $\text{Arr}(\mathcal{C})$. The objects are arrows of \mathcal{C} . What are the arrows of $\text{Arr}(\mathcal{C})$?

Suppose $f : A \rightarrow B, g : C \rightarrow D \in \text{Arr}(\mathcal{C})_0$. An arrow $x : f \rightarrow g$ in the arrow category is a pair of arrows $x_1 : A \rightarrow C$ and $x_2 : B \rightarrow D$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{x_1} & C \\ f \downarrow & & \downarrow g \\ B & \xrightarrow{x_2} & D \end{array} \quad , \text{ i.e. } x_2 \circ f = g \circ x_1.$$

Identity and composition are defined in the obvious way.

Hom-sets

The categories we see in our everyday life are **locally small** categories, i.e. $\text{Hom}_{\mathcal{C}}(A, B)$ are sets. We shall not worry about it anymore and assume that all categories are locally small.

Hom-sets

For a locally small category \mathcal{C} , and $X \in \mathcal{C}_0$, we have a functor

$$\mathrm{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Sets},$$

called the Hom functor.

Hom-sets

For a locally small category \mathcal{C} , and $X \in \mathcal{C}_0$, we have a functor

$$\mathrm{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Sets},$$

called the Hom functor.

It takes objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(X, A)$.

Hom-sets

For a locally small category \mathcal{C} , and $X \in \mathcal{C}_0$, we have a functor

$$\mathrm{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Sets},$$

called the Hom functor.

It takes objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(X, A)$.

Given an arrow $f : A \rightarrow B$ in \mathcal{C} , the Hom functor is supposed to take it to a set function $f^* : \mathrm{Hom}_{\mathcal{C}}(X, A) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, B)$. How does f^* work?

Hom-sets

For a locally small category \mathcal{C} , and $X \in \mathcal{C}_0$, we have a functor

$$\mathrm{Hom}_{\mathcal{C}}(X, -) : \mathcal{C} \rightarrow \mathbf{Sets},$$

called the Hom functor.

It takes objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(X, A)$.

Given an arrow $f : A \rightarrow B$ in \mathcal{C} , the Hom functor is supposed to take it to a set function $f^* : \mathrm{Hom}_{\mathcal{C}}(X, A) \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, B)$. How does f^* work?

$$\begin{array}{ccc} X & & \\ g \downarrow & \searrow f^*(g)=f \circ g & \\ A & \xrightarrow{f} & B \end{array}$$

Similarly can we define a functor

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}?$$

Hom-sets

Similarly can we define a functor

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}?$$

It will take objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(A, X)$.

Hom-sets

Similarly can we define a functor

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}?$$

It will take objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(A, X)$.

Given an arrow $f : A \rightarrow B$ in \mathcal{C} , the this functor is supposed to take it to a set function $f_* : \mathrm{Hom}_{\mathcal{C}}(A, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(B, X)$. How should f_* work?

Hom-sets

Similarly can we define a functor

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}?$$

It will take objects $A \in \mathcal{C}_0$ to the set $\mathrm{Hom}_{\mathcal{C}}(A, X)$.

Given an arrow $f : A \rightarrow B$ in \mathcal{C} , the this functor is supposed to take it to a set function $f_* : \mathrm{Hom}_{\mathcal{C}}(A, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(B, X)$. How should f_* work?

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ g \downarrow & \swarrow f_*(g)=??? & \\ X & & \end{array}$$

Hom-sets

Given an arrow $f : A \rightarrow B$, seemingly there's an issue with defining $f_* : \text{Hom}_{\mathcal{C}}(A, X) \rightarrow \text{Hom}_{\mathcal{C}}(B, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ g \downarrow & \swarrow f_*(g)=??? & \\ X & & \end{array}$$

Hom-sets

Given an arrow $f : A \rightarrow B$, seemingly there's an issue with defining $f_* : \text{Hom}_{\mathcal{C}}(A, X) \rightarrow \text{Hom}_{\mathcal{C}}(B, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ g \downarrow & \swarrow f_*(g)=??? & \\ X & & \end{array}$$

But what we can do is, we can define $f_* : \text{Hom}_{\mathcal{C}}(B, X) \rightarrow \text{Hom}_{\mathcal{C}}(A, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ f_*(g)=g \circ f \downarrow & \swarrow g & \\ X & & \end{array}$$

Hom-sets

Given an arrow $f : A \rightarrow B$, seemingly there's an issue with defining $f_* : \text{Hom}_{\mathcal{C}}(A, X) \rightarrow \text{Hom}_{\mathcal{C}}(B, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ g \downarrow & \swarrow f_*(g)=??? & \\ X & & \end{array}$$

But what we can do is, we can define $f_* : \text{Hom}_{\mathcal{C}}(B, X) \rightarrow \text{Hom}_{\mathcal{C}}(A, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ f_*(g)=g \circ f \downarrow & \swarrow g & \\ X & & \end{array}$$

So $\text{Hom}_{\mathcal{C}}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}$ reverses the direction of arrows?

Hom-sets

The solution is tweaking the domain category to be the opposite category of \mathcal{C} .

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Sets}.$$

Hom-sets

The solution is tweaking the domain category to be the opposite category of \mathcal{C} .

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Sets}.$$

The objects of $\mathcal{C}^{\mathrm{op}}$ are the same as \mathcal{C} .

Hom-sets

The solution is tweaking the domain category to be the opposite category of \mathcal{C} .

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Sets}.$$

The objects of $\mathcal{C}^{\mathrm{op}}$ are the same as \mathcal{C} . The arrows of $\mathcal{C}^{\mathrm{op}}$ are $f^{\mathrm{op}} : B \rightarrow A$, corresponding to an arrow $f : A \rightarrow B$ in \mathcal{C} .

Hom-sets

The solution is tweaking the domain category to be the opposite category of \mathcal{C} .

$$\mathrm{Hom}_{\mathcal{C}}(-, X) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Sets}.$$

The objects of $\mathcal{C}^{\mathrm{op}}$ are the same as \mathcal{C} . The arrows of $\mathcal{C}^{\mathrm{op}}$ are $f^{\mathrm{op}} : B \rightarrow A$, corresponding to an arrow $f : A \rightarrow B$ in \mathcal{C} .

The functor $\mathrm{Hom}_{\mathcal{C}}(-, X)$ takes the arrow $f^{\mathrm{op}} : B \rightarrow A$ to a set function $f_* : \mathrm{Hom}_{\mathcal{C}}(B, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(A, X)$.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ f_*(g)=g \circ f \downarrow & \swarrow g & \\ X & & \end{array}$$

Hom-sets

Generalizing further, we define the functor

$$\mathrm{Hom}_{\mathcal{C}}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Sets}.$$

Hom-sets

Generalizing further, we define the functor

$$\mathrm{Hom}_{\mathcal{C}}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Sets}.$$

In the object level, it takes (A, B) to the set $\mathrm{Hom}_{\mathcal{C}}(A, B)$.

Hom-sets

Generalizing further, we define the functor

$$\mathrm{Hom}_{\mathcal{C}}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Sets}.$$

In the object level, it takes (A, B) to the set $\mathrm{Hom}_{\mathcal{C}}(A, B)$.

In the arrow level, let (f^{op}, g) be an arrow in $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$. Then $f : A \rightarrow B$ and $g : X \rightarrow Y$ are arrows in \mathcal{C} .

Hom-sets

Generalizing further, we define the functor

$$\mathrm{Hom}_{\mathcal{C}}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Sets}.$$

In the object level, it takes (A, B) to the set $\mathrm{Hom}_{\mathcal{C}}(A, B)$.

In the arrow level, let (f^{op}, g) be an arrow in $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$. Then $f : A \rightarrow B$ and $g : X \rightarrow Y$ are arrows in \mathcal{C} .

$\mathrm{Hom}_{\mathcal{C}}(f^{\mathrm{op}}, g)$ will be a set function $\mathrm{Hom}_{\mathcal{C}}(B, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(A, Y)$.

Hom-sets

Generalizing further, we define the functor

$$\mathrm{Hom}_{\mathcal{C}}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Sets}.$$

In the object level, it takes (A, B) to the set $\mathrm{Hom}_{\mathcal{C}}(A, B)$.

In the arrow level, let (f^{op}, g) be an arrow in $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$. Then $f : A \rightarrow B$ and $g : X \rightarrow Y$ are arrows in \mathcal{C} .

$\mathrm{Hom}_{\mathcal{C}}(f^{\mathrm{op}}, g)$ will be a set function $\mathrm{Hom}_{\mathcal{C}}(B, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(A, Y)$.

$$\begin{array}{ccc} A & & X \\ f \downarrow & \nearrow x & \downarrow g \\ B & & Y \end{array}$$

Given $x \in \mathrm{Hom}_{\mathcal{C}}(B, X)$, we define $\mathrm{Hom}_{\mathcal{C}}(f^{\mathrm{op}}, g) = g \circ x \circ f$.

Isomorphisms

We define

- isomorphism between groups,
- isomorphism between vector spaces,
- homeomorphism between topological spaces,
- diffeomorphisms between smooth manifolds,
- and so on . . .

Category theory captures this pattern of “sameness” as well.

Definition 3

An arrow $f : A \rightarrow B$ in a category \mathcal{C} is called an **isomorphism** if there exists another arrow $g : B \rightarrow A$ such that $f \circ g = \mathbb{1}_B$ and $g \circ f = \mathbb{1}_A$.
If there is an isomorphism from A to B , we call A and B **isomorphic objects**.

Definition 3

An arrow $f : A \rightarrow B$ in a category \mathcal{C} is called an **isomorphism** if there exists another arrow $g : B \rightarrow A$ such that $f \circ g = \mathbb{1}_B$ and $g \circ f = \mathbb{1}_A$.
If there is an isomorphism from A to B , we call A and B **isomorphic objects**.

This definition aligns with our definition of isomorphisms in other categories.

Isomorphism and Functors

Suppose A and B are isomorphic objects in a category \mathcal{C} . Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then are $F(A)$ and $F(B)$ isomorphic objects in the category \mathcal{D} ?

Isomorphism and Functors

Suppose A and B are isomorphic objects in a category \mathcal{C} . Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then are $F(A)$ and $F(B)$ isomorphic objects in the category \mathcal{D} ?

 \mathcal{C} \mathcal{D}

$$A \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} B$$

$$F(A) \begin{array}{c} \xrightarrow{F(f)} \\ \xleftarrow{F(g)} \end{array} F(B)$$

Isomorphism and Functors

Suppose A and B are isomorphic objects in a category \mathcal{C} . Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then are $F(A)$ and $F(B)$ isomorphic objects in the category \mathcal{D} ?

 \mathcal{C} \mathcal{D}

$$A \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} B$$

$$F(A) \begin{array}{c} \xrightarrow{F(f)} \\ \xleftarrow{F(g)} \end{array} F(B)$$

$$F(f) \circ F(g) = F(f \circ g) = F(1_A) = 1_{F(A)}.$$

$$F(g) \circ F(f) = F(g \circ f) = F(1_B) = 1_{F(B)}.$$

Isomorphism and Functors

Suppose A and B are isomorphic objects in a category \mathcal{C} . Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then are $F(A)$ and $F(B)$ isomorphic objects in the category \mathcal{D} ?

 \mathcal{C} \mathcal{D}

$$A \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} B$$

$$F(A) \begin{array}{c} \xrightarrow{F(f)} \\ \xleftarrow{F(g)} \end{array} F(B)$$

$$F(f) \circ F(g) = F(f \circ g) = F(1_A) = 1_{F(A)}.$$

$$F(g) \circ F(f) = F(g \circ f) = F(1_B) = 1_{F(B)}.$$

Therefore, functors preserve isomorphisms.

Isomorphism and Hom-sets

Suppose A and B are isomorphic objects in a category. Then for any other object X , since functors preserve isomorphisms, $\text{Hom}_{\mathcal{C}}(X, A)$ and $\text{Hom}_{\mathcal{C}}(X, B)$ are isomorphic objects in **Sets**.

Isomorphism and Hom-sets

Suppose A and B are isomorphic objects in a category. Then for any other object X , since functors preserve isomorphisms, $\text{Hom}_{\mathcal{C}}(X, A)$ and $\text{Hom}_{\mathcal{C}}(X, B)$ are isomorphic objects in **Sets**.

Similarly, $\text{Hom}_{\mathcal{C}}(A, X)$ and $\text{Hom}_{\mathcal{C}}(B, X)$ are also isomorphic objects in **Sets**.

Isomorphism and Hom-sets


Suppose A and B are isomorphic objects in a category. Then for any other object X , since functors preserve isomorphisms, $\text{Hom}_{\mathcal{C}}(X, A)$ and $\text{Hom}_{\mathcal{C}}(X, B)$ are isomorphic objects in **Sets**.

Similarly, $\text{Hom}_{\mathcal{C}}(A, X)$ and $\text{Hom}_{\mathcal{C}}(B, X)$ are also isomorphic objects in **Sets**.

Does the converse hold? If $\text{Hom}_{\mathcal{C}}(A, X)$ and $\text{Hom}_{\mathcal{C}}(B, X)$ are also isomorphic objects in **Sets**, then can we say that A and B are isomorphic objects in \mathcal{C} ?

Isomorphisms

When someone says, "a finite dimensional vector space V is isomorphic to its dual V^* "



Natural Transformation

Given two categories \mathcal{C} and \mathcal{D} , we can form the functor category $\text{Fun}(\mathcal{C}, \mathcal{D})$ whose objects are all the functors from \mathcal{C} to \mathcal{D} . What should be the arrows in this category?

This is similar to the construction of arrow category.

Natural Transformation

Definition 4

Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be two functors. Then a **natural transformation** $\eta : F \Rightarrow G$ is a family of arrows

$$\{\eta_X : F(X) \rightarrow G(X)\}_{X \in \mathcal{C}_0}$$

in \mathcal{D} such that for every arrow $f : X \rightarrow Y$ in \mathcal{C} , the following diagram commutes:


$$\begin{array}{ccc} F(X) & \xrightarrow{F(f)} & F(Y) \\ \eta_X \downarrow & & \downarrow \eta_Y \\ G(X) & \xrightarrow{G(f)} & G(Y) \end{array}$$

In other words, $\eta_Y \circ F(f) = G(f) \circ \eta_X$.

Given such a natural transformation $\eta : F \Rightarrow G$, the arrow η_X is called the component of η at X .

Natural Transformation

If $F, G : \mathcal{C} \rightarrow \mathcal{D}$ are functors, and $\eta : F \Rightarrow G$ is a natural transformation, it is denoted as follows



Natural Transformation

Why do we care about natural transformations?

Natural Transformation


Why do we care about natural transformations?

Because they take commutative diagrams to commutative diagrams!!!

Natural Transformation

Why do we care about natural transformations?

Because they take commutative diagrams to commutative diagrams!!!



Natural Transformation



If F and G are functors between the categories C and D , then a **natural transformation** η from F to G is a family of morphisms that satisfies two requirements.

1. The natural transformation must associate, to every object X in C , a morphism $\eta_X : F(X) \rightarrow G(X)$ between objects of D . The morphism η_X is called the **component** of η at X .
2. Components must be such that for every morphism $f : X \rightarrow Y$ in C we have:

$$\eta_Y \circ F(f) = G(f) \circ \eta_X$$

The last equation can conveniently be expressed by the **commutative diagram**

$$\begin{array}{ccccc}
 X & & F(X) & \xrightarrow{\eta_X} & G(X) \\
 f \downarrow & & F(f) \downarrow & & \downarrow G(f) \\
 Y & & F(Y) & \xrightarrow{\eta_Y} & G(Y)
 \end{array}$$

Natural Isomorphism

Definition 5

Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be functors. A natural transformation $\eta : F \Rightarrow G$ is called a **natural isomorphism** if all its components

$$\{\eta_X : F(X) \rightarrow G(X)\}_{X \in \mathcal{C}_0}$$

are isomorphisms in \mathcal{D} .

Natural Isomorphism

Definition 5

Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be functors. A natural transformation $\eta : F \Rightarrow G$ is called a **natural isomorphism** if all its components

$$\{\eta_X : F(X) \rightarrow G(X)\}_{X \in \mathcal{C}_0}$$

are isomorphisms in \mathcal{D} .

Since natural transformations are arrows in the functor category $\text{Fun}(\mathcal{C}, \mathcal{D})$, natural isomorphisms are just isomorphisms in $\text{Fun}(\mathcal{C}, \mathcal{D})$.

Natural Isomorphism

What do we mean when we say V and V^{**} are “naturally isomorphic” (when V is a finite dimensional vector space)? Where is the natural isomorphism here?

Natural Isomorphism

What do we mean when we say V and V^{**} are “naturally isomorphic” (when V is a finite dimensional vector space)? Where is the natural isomorphism here?

Let $\mathbf{Vect}_{\mathbb{K}}^{\text{fin}}$ be the category of all finite dimensional vector spaces over the field \mathbb{K} . Consider the functors $\mathbb{1}_{\mathbf{Vect}_{\mathbb{K}}^{\text{fin}}}, (-)^{**} : \mathbf{Vect}_{\mathbb{K}}^{\text{fin}} \rightarrow \mathbf{Vect}_{\mathbb{K}}^{\text{fin}}$.

Natural Isomorphism

What do we mean when we say V and V^{**} are “naturally isomorphic” (when V is a finite dimensional vector space)? Where is the natural isomorphism here?

Let $\mathbf{Vect}_{\mathbb{K}}^{\text{fin}}$ be the category of all finite dimensional vector spaces over the field \mathbb{K} . Consider the functors $\mathbb{1}_{\mathbf{Vect}_{\mathbb{K}}^{\text{fin}}}, (-)^{**} : \mathbf{Vect}_{\mathbb{K}}^{\text{fin}} \rightarrow \mathbf{Vect}_{\mathbb{K}}^{\text{fin}}$.

The second functor sends a vector space V to its double dual V^{**} , and a linear map $f : V \rightarrow W$ to $(f^T)^T : V^{**} \rightarrow W^{**}$.

Natural Isomorphism

Then we can define an isomorphism $\eta_V : V \rightarrow V^{**}$ such that

$$\eta_V(\mathbf{v})(\varphi) = \varphi(\mathbf{v}),$$

for $\mathbf{v} \in V$ and $\varphi \in V^*$.


Natural Isomorphism

Then we can define an isomorphism $\eta_V : V \rightarrow V^{**}$ such that

$$\eta_V(\mathbf{v})(\varphi) = \varphi(\mathbf{v}),$$

for $\mathbf{v} \in V$ and $\varphi \in V^*$.

Then $\{\eta_V : V \rightarrow V^{**}\}_{V \in (\mathbf{Vect}_{\mathbb{K}}^{\text{fin}})_0}$ is a natural isomorphism, because the following diagram commutes:



Adjoint

Isomorphism	Equivalence	Adjoint
$F : \mathcal{C} \rightarrow \mathcal{D},$ $G : \mathcal{D} \rightarrow \mathcal{C},$ s.t. $F \circ G = \mathbb{1}_{\mathcal{D}}$ and $G \circ F = \mathbb{1}_{\mathcal{C}}$	$F : \mathcal{C} \rightarrow \mathcal{D},$ $G : \mathcal{D} \rightarrow \mathcal{C},$ s.t. natural isomorphisms $F \circ G \Rightarrow \mathbb{1}_{\mathcal{D}}$ and $G \circ F \Rightarrow \mathbb{1}_{\mathcal{C}}$	$F : \mathcal{C} \rightarrow \mathcal{D},$ $G : \mathcal{D} \rightarrow \mathcal{C},$ s.t. natural transformations $F \circ G \Rightarrow \mathbb{1}_{\mathcal{D}}$ and $G \circ F \Leftarrow \mathbb{1}_{\mathcal{C}}$

Adjoint

Definition 6

Two functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are called **adjoints** to each other if

$$\mathrm{Hom}_{\mathcal{D}}(F(C), D) \cong \mathrm{Hom}_{\mathcal{C}}(C, G(D)) \text{ naturally.}$$

Adjoint

Definition 6

Two functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are called **adjoints** to each other if

$$\mathrm{Hom}_{\mathcal{D}}(F(C), D) \cong \mathrm{Hom}_{\mathcal{C}}(C, G(D)) \text{ naturally.}$$

In other words, there exists a natural isomorphism η :

$$\begin{array}{ccc} & \mathrm{Hom}_{\mathcal{D}}(F(-), -) & \\ \swarrow & \Downarrow \eta & \searrow \\ \mathcal{C}^{\mathrm{op}} \times \mathcal{D} & & \mathbf{Sets} \\ \nwarrow & \Downarrow & \nearrow \\ & \mathrm{Hom}_{\mathcal{C}}(-, G(-)) & \end{array}$$

Adjoint

Definition 6

Two functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are called **adjoints** to each other if

$$\mathrm{Hom}_{\mathcal{D}}(F(C), D) \cong \mathrm{Hom}_{\mathcal{C}}(C, G(D)) \text{ naturally.}$$

In other words, there exists a natural isomorphism η :

$$\begin{array}{ccc} & \mathrm{Hom}_{\mathcal{D}}(F(-), -) & \\ \swarrow & \Downarrow \eta & \searrow \\ \mathcal{C}^{\mathrm{op}} \times \mathcal{D} & & \mathbf{Sets} \\ \nwarrow & \Downarrow & \nearrow \\ & \mathrm{Hom}_{\mathcal{C}}(-, G(-)) & \end{array}$$

If this happens, we call F a **left adjoint** of G ; and we call G a **right adjoint** of F . We write this as $F \dashv G$.

Adjoint Example

Tensor product of vector spaces

The **tensor product** of two \mathbb{K} -vector spaces U and V is another \mathbb{K} -vector space $U \otimes V$ equipped with a bilinear map $\theta : U \times V \rightarrow U \otimes V$ that is *universal*:

Adjoint Example

Tensor product of vector spaces

The **tensor product** of two \mathbb{K} -vector spaces U and V is another \mathbb{K} -vector space $U \otimes V$ equipped with a bilinear map $\theta : U \times V \rightarrow U \otimes V$ that is *universal*: for any bilinear map $\beta : U \times V \rightarrow W$, there exists a unique linear map $\alpha : U \otimes V \rightarrow W$ such that the following diagram commutes:

$$\begin{array}{ccc} U \times V & \xrightarrow{\theta} & U \otimes V \\ \forall \text{ bilinear } \beta \downarrow & \swarrow \exists! \alpha & \\ W & & \end{array}$$

In other words, $\beta = \alpha \circ \theta$.

Adjoint Example

So we have a 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{bilinear maps } U \times V \rightarrow W\}.$$

Adjoint Example

So we have a 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{bilinear maps } U \times V \rightarrow W\}.$$

We also have another 1-1 correspondence

$$\{\text{bilinear maps } U \times V \rightarrow W\} \leftrightarrow \{\text{linear maps } U \rightarrow \text{Hom}(V, W)\};$$

Adjoint Example

So we have a 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{bilinear maps } U \times V \rightarrow W\}.$$

We also have another 1-1 correspondence

$$\begin{aligned} \{\text{bilinear maps } U \times V \rightarrow W\} &\leftrightarrow \{\text{linear maps } U \rightarrow \text{Hom}(V, W)\}; \\ (f : U \times V \rightarrow W) &\mapsto (\mathbf{u} \mapsto f(\mathbf{u}, -)), \end{aligned}$$

Adjoint Example

So we have a 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{bilinear maps } U \times V \rightarrow W\}.$$

We also have another 1-1 correspondence

$$\begin{aligned} \{\text{bilinear maps } U \times V \rightarrow W\} &\leftrightarrow \{\text{linear maps } U \rightarrow \text{Hom}(V, W)\}; \\ (f : U \times V \rightarrow W) &\mapsto (\mathbf{u} \mapsto f(\mathbf{u}, -)), \\ ((\mathbf{u}, \mathbf{v}) \mapsto g(\mathbf{u})(\mathbf{v})) &\mapsto (g : U \rightarrow \text{Hom}(V, W)). \end{aligned}$$

Adjoint Example

So we have a 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{bilinear maps } U \times V \rightarrow W\}.$$

We also have another 1-1 correspondence

$$\begin{aligned} \{\text{bilinear maps } U \times V \rightarrow W\} &\leftrightarrow \{\text{linear maps } U \rightarrow \text{Hom}(V, W)\}; \\ (f : U \times V \rightarrow W) &\mapsto (\mathbf{u} \mapsto f(\mathbf{u}, -)), \\ ((\mathbf{u}, \mathbf{v}) \mapsto g(\mathbf{u})(\mathbf{v})) &\mapsto (g : U \rightarrow \text{Hom}(V, W)). \end{aligned}$$

Therefore, we have the 1-1 correspondence

$$\{\text{linear maps } U \otimes V \rightarrow W\} \leftrightarrow \{\text{linear maps } U \rightarrow \text{Hom}(V, W)\}.$$

Adjoint Example

In other words,

$$\mathrm{Hom}(U \otimes V, W) \cong \mathrm{Hom}(U, \mathrm{Hom}(V, W)).$$

Does that mean $- \otimes V$ and $\mathrm{Hom}(V, -) : \mathbf{Vect}_{\mathbb{K}} \rightarrow \mathbf{Vect}_{\mathbb{K}}$ are adjoint functors?

Adjoint Example

In other words,

$$\mathrm{Hom}(U \otimes V, W) \cong \mathrm{Hom}(U, \mathrm{Hom}(V, W)).$$

Does that mean $- \otimes V$ and $\mathrm{Hom}(V, -) : \mathbf{Vect}_{\mathbb{K}} \rightarrow \mathbf{Vect}_{\mathbb{K}}$ are adjoint functors? Well, not yet. We need to show the naturality of this isomorphism.

$$\begin{array}{ccc} & \mathrm{Hom}_{\mathbf{Vect}_{\mathbb{K}}}(- \otimes V, -) & \\ \mathrm{Vect}_{\mathbb{K}}^{\mathrm{op}} \times \mathrm{Vect}_{\mathbb{K}} & \begin{array}{c} \downarrow \eta \\ \downarrow \end{array} & \mathbf{Sets} \\ & \mathrm{Hom}_{\mathbf{Vect}_{\mathbb{K}}}(-, \mathrm{Hom}(V, -)) & \end{array}$$

Adjoint Example

Given an arrow $(\alpha_1^{\text{op}}, \alpha_2) : (U, W) \rightarrow (U', W)$ in $\mathbf{Vect}_{\mathbb{K}}^{\text{op}} \times \mathbf{Vect}_{\mathbb{K}}$, we need to show the commutativity of the following diagram in the category **Sets**:

$$\begin{array}{ccc} \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U \otimes V, W) & \xrightarrow{\eta_{(U, W)}} & \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U, \text{Hom}(V, W)) \\ \downarrow F(\alpha_1^{\text{op}}, \alpha_2) & & \downarrow G(\alpha_1^{\text{op}}, \alpha_2) \\ \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U' \otimes V, W') & \xrightarrow{\eta_{(U', W')}} & \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U', \text{Hom}(V, W')) \end{array}$$

where $F = \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (- \otimes V, -)$ and $G = \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (-, \text{Hom}(V, -))$.

Adjoint Example

Given an arrow $(\alpha_1^{\text{op}}, \alpha_2) : (U, W) \rightarrow (U', W)$ in $\mathbf{Vect}_{\mathbb{K}}^{\text{op}} \times \mathbf{Vect}_{\mathbb{K}}$, we need to show the commutativity of the following diagram in the category **Sets**:

$$\begin{array}{ccc} \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U \otimes V, W) & \xrightarrow{\eta_{(U, W)}} & \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U, \text{Hom}(V, W)) \\ \downarrow F(\alpha_1^{\text{op}}, \alpha_2) & & \downarrow G(\alpha_1^{\text{op}}, \alpha_2) \\ \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U' \otimes V, W') & \xrightarrow{\eta_{(U', W')}} & \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (U', \text{Hom}(V, W')) \end{array}$$

where $F = \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (- \otimes V, -)$ and $G = \text{Hom}_{\mathbf{Vect}_{\mathbb{K}}} (-, \text{Hom}(V, -))$.

And this diagram indeed commutes! So $- \otimes V$ is the left adjoint of $\text{Hom}(V, -)$.

Adjoint

Are adjoints unique? Can a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ have two right adjoints $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{C}$?

Adjoint

Are adjoints unique? Can a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ have two right adjoints $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{C}$? If this happens, then for any $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,

Adjoint

Are adjoints unique? Can a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ have two right adjoints $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{C}$? If this happens, then for any $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,

$$\mathrm{Hom}_{\mathcal{C}}(C, G_1(D)) \cong \mathrm{Hom}_{\mathcal{D}}(F(C), D)$$

Adjoint

Are adjoints unique? Can a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ have two right adjoints $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{C}$? If this happens, then for any $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,

$$\begin{aligned}\mathrm{Hom}_{\mathcal{C}}(C, G_1(D)) &\cong \mathrm{Hom}_{\mathcal{D}}(F(C), D) \\ &\cong \mathrm{Hom}_{\mathcal{C}}(C, G_2(D)).\end{aligned}$$

Adjoint

Are adjoints unique? Can a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ have two right adjoints $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{C}$? If this happens, then for any $C \in \mathcal{C}_0$ and $D \in \mathcal{D}_0$,

$$\begin{aligned}\mathrm{Hom}_{\mathcal{C}}(C, G_1(D)) &\cong \mathrm{Hom}_{\mathcal{D}}(F(C), D) \\ &\cong \mathrm{Hom}_{\mathcal{C}}(C, G_2(D)).\end{aligned}$$

Therefore,

$$\mathrm{Hom}_{\mathcal{C}}(C, G_1(D)) \cong \mathrm{Hom}_{\mathcal{C}}(C, G_2(D)).$$

Does this mean G_1 and G_2 are isomorphic functors?

Yoneda Lemma



Yoneda Lemma

Theorem 6 (Yoneda Lemma)

For any functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}$ and any $X \in \mathcal{C}_0$, the natural transformations $\text{Hom}_{\mathcal{C}}(-, X) \Rightarrow F$ are in bijection with the elements of the set $F(X)$. In other words,

$$\text{Hom}_{\text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})}(\text{Hom}_{\mathcal{C}}(-, X), F) \cong F(X),$$

and this isomorphism is natural in both F and X .

Yoneda Lemma

Theorem 6 (Yoneda Lemma)

For any functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}$ and any $X \in \mathcal{C}_0$, the natural transformations $\text{Hom}_{\mathcal{C}}(-, X) \Rightarrow F$ are in bijection with the elements of the set $F(X)$. In other words,

$$\text{Hom}_{\text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})}(\text{Hom}_{\mathcal{C}}(-, X), F) \cong F(X),$$

and this isomorphism is natural in both F and X .

The last line means that we have a natural isomorphism

$$\begin{array}{ccc} (X, F) \mapsto \text{Hom}_{\text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})}(\text{Hom}_{\mathcal{C}}(-, X), F) & & \\ \text{C}^{\text{op}} \times \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets}) & \xrightarrow{\quad \eta \quad} & \mathbf{Sets} \\ (X, F) \mapsto F(X) & & \end{array}$$

Yoneda Lemma

What does it even mean?

Yoneda Lemma

What does it even mean?

Definition 7

A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *fully faithful* if the set-functions

$$F_{X,Y} : \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}}(F(X), F(Y))$$

are bijective for all $X, Y \in \mathcal{C}_0$.

Yoneda Lemma

What does it even mean?

Definition 7

A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *fully faithful* if the set-functions

$$F_{X,Y} : \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}}(F(X), F(Y))$$

are bijective for all $X, Y \in \mathcal{C}_0$.

Theorem 7

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a fully faithful functor. Then $X \cong Y$ in \mathcal{C} if and only if $F(X) \cong F(Y)$ in \mathcal{D} .

Corollary 8

The **Yoneda embedding**

$$\mathcal{Y} : \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})$$

$$X \mapsto \text{Hom}(-, X)$$

$$(f : X \rightarrow Y) \mapsto (f_* : \text{Hom}_{\mathcal{C}}(-, X) \rightarrow \text{Hom}_{\mathcal{C}}(-, Y))$$

is *fully faithful*.

Yoneda Lemma

Corollary 8

The **Yoneda embedding**

$$\mathcal{Y} : \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})$$

$$X \mapsto \text{Hom}(-, X)$$

$$(f : X \rightarrow Y) \mapsto (f_* : \text{Hom}_{\mathcal{C}}(-, X) \rightarrow \text{Hom}_{\mathcal{C}}(-, Y))$$

is *fully faithful*.

Proof.

Take $F = \text{Hom}_{\mathcal{C}}(-, Y)$ in Yoneda Lemma. This gives us

$$\text{Hom}_{\text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})}(\text{Hom}_{\mathcal{C}}(-, X), \text{Hom}_{\mathcal{C}}(-, Y)) \cong \text{Hom}_{\mathcal{C}}(X, Y).$$

Isomorphism in sets is bijection!!



Yoneda Lemma

Corollary 9

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(-, X)$ and $\text{Hom}_{\mathcal{C}}(-, Y)$ are naturally isomorphic functors.

Yoneda Lemma

Corollary 9

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(-, X)$ and $\text{Hom}_{\mathcal{C}}(-, Y)$ are naturally isomorphic functors.

Proof.

Because Yoneda embedding $X \mapsto \text{Hom}_{\mathcal{C}}(-, X)$ is fully faithful!!! ■

Yoneda Lemma

Corollary 9

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(-, X)$ and $\text{Hom}_{\mathcal{C}}(-, Y)$ are naturally isomorphic functors.

Proof.

Because Yoneda embedding $X \mapsto \text{Hom}_{\mathcal{C}}(-, X)$ is fully faithful!!! ■

We usually use this variant of Yoneda lemma to prove isomorphisms.

Yoneda Lemma Summarized

Interchanging \mathcal{C} and \mathcal{C}^{op} throughout, we get the following result:

Corollary 10

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(X, -)$ and $\text{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic functors.

Yoneda Lemma Summarized

Interchanging \mathcal{C} and \mathcal{C}^{op} throughout, we get the following result:

Corollary 10

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(X, -)$ and $\text{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic functors.

Proof.



Yoneda Lemma Summarized

Interchanging \mathcal{C} and \mathcal{C}^{op} throughout, we get the following result:

Corollary 10

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(X, -)$ and $\text{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic functors.

Proof.

$$X \cong Y \text{ in } \mathcal{C} \iff X \cong Y \text{ in } \mathcal{C}^{\text{op}}$$



Yoneda Lemma Summarized

Interchanging \mathcal{C} and \mathcal{C}^{op} throughout, we get the following result:

Corollary 10

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(X, -)$ and $\text{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic functors.

Proof.

$$\begin{aligned} X \cong Y \text{ in } \mathcal{C} &\iff X \cong Y \text{ in } \mathcal{C}^{\text{op}} \\ &\iff \text{Hom}_{\mathcal{C}^{\text{op}}}(-, X) \cong \text{Hom}_{\mathcal{C}^{\text{op}}}(-, Y) \end{aligned}$$



Yoneda Lemma Summarized

Interchanging \mathcal{C} and \mathcal{C}^{op} throughout, we get the following result:

Corollary 10

In a category \mathcal{C} , $X \cong Y$ if and only if $\text{Hom}_{\mathcal{C}}(X, -)$ and $\text{Hom}_{\mathcal{C}}(Y, -)$ are naturally isomorphic functors.

Proof.

$$\begin{aligned} X \cong Y \text{ in } \mathcal{C} &\iff X \cong Y \text{ in } \mathcal{C}^{\text{op}} \\ &\iff \text{Hom}_{\mathcal{C}^{\text{op}}}(X, -) \cong \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, -) \\ &\iff \text{Hom}_{\mathcal{C}}(-, X) \cong \text{Hom}_{\mathcal{C}}(-, Y) \\ &\iff \text{Hom}_{\mathcal{C}}(X, -) \cong \text{Hom}_{\mathcal{C}}(Y, -). \end{aligned}$$



Yoneda Lemma Summarized



Yoneda Lemma



Tell me who your
Hom-ies are and I'll
tell you who you are

References

- ① *Category Theory*, by Steve Awodey
- ② *Category Theory in Context*, by Emily Riehl
- ③ *Basic Category Theory*, by Tom Leinster
- ④ *Categories for the Working Mathematician*, by Saunders Mac Lane
- ⑤ Math3ma blog: <https://www.math3ma.com/blog/the-yoneda-lemma>
- ⑥ Ncatlab: <https://ncatlab.org/nlab/show/Yoneda+lemma>

Thank you for joining!

The slides are available in my webpage

https://atonurc.github.io/assets/catrep_talk_1.pdf

