

The Baker–Campbell–Hausdorff Formula and Lie's Third Theorem

Atonu Roy Chowdhury

BRAC University

atonuroychowdhury@gmail.com

5 January, 2026

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem

Definition 1

A **Lie group** is a group G , that also has a manifold structure;

Definition 1

A **Lie group** is a group G , that also has a manifold structure; and these two structures are compatible,

Definition 1

A **Lie group** is a group G , that also has a manifold structure; and these two structures are compatible, i.e. the multiplication map

$$\begin{aligned} m : G \times G &\rightarrow G \\ (g, h) &\mapsto gh \end{aligned}$$

and the inverse map

$$\begin{aligned} i : G &\rightarrow G \\ g &\mapsto g^{-1} \end{aligned}$$

are smooth.

Examples of Lie Groups

Example 1

- $(\mathbb{R}, +)$

Examples of Lie Groups

Example 1

- $(\mathbb{R}, +)$
- $(\mathbb{R}_{>0}, \times)$

Examples of Lie Groups

Example 1

- $(\mathbb{R}, +)$
- $(\mathbb{R}_{>0}, \times)$
- $GL(n, \mathbb{R})$ with respect to matrix multiplication

Examples of Lie Groups

Example 1

- $(\mathbb{R}, +)$
- $(\mathbb{R}_{>0}, \times)$
- $GL(n, \mathbb{R})$ with respect to matrix multiplication
- $SL(n, \mathbb{R})$ with respect to matrix multiplication

Definition 2 (Lie algebra)

A **Lie algebra** over a field \mathbb{K} is a \mathbb{K} -vector space V equipped with a bilinear operation $[-, -] : V \times V \rightarrow V$ (called the **Lie bracket**) such that

Definition 2 (Lie algebra)

A **Lie algebra** over a field \mathbb{K} is a \mathbb{K} -vector space V equipped with a bilinear operation $[-, -] : V \times V \rightarrow V$ (called the **Lie bracket**) such that

- ④ $[-, -]$ is antisymmetric, i.e. $[v_1, v_2] = -[v_2, v_1]$ for all $v_1, v_2 \in V$; and

Definition 2 (Lie algebra)

A **Lie algebra** over a field \mathbb{K} is a \mathbb{K} -vector space V equipped with a bilinear operation $[-, -] : V \times V \rightarrow V$ (called the **Lie bracket**) such that

- Ⓐ $[-, -]$ is antisymmetric, i.e. $[v_1, v_2] = -[v_2, v_1]$ for all $v_1, v_2 \in V$; and
- Ⓑ for any $u, v, w \in V$,

$$[[u, v], w] + [[w, u], v] + [[v, w], u] = 0. \quad (1)$$

Definition 2 (Lie algebra)

A **Lie algebra** over a field \mathbb{K} is a \mathbb{K} -vector space V equipped with a bilinear operation $[-, -] : V \times V \rightarrow V$ (called the **Lie bracket**) such that

- Ⓐ $[-, -]$ is antisymmetric, i.e. $[v_1, v_2] = -[v_2, v_1]$ for all $v_1, v_2 \in V$; and
- Ⓑ for any $u, v, w \in V$,

$$[[u, v], w] + [[w, u], v] + [[v, w], u] = 0. \quad (1)$$

(1) is also known as the Jacobi identity.

Examples of Lie Algebra

Example 2

If V is an associative \mathbb{K} -algebra with respect to the product \cdot , then we can define the Lie bracket as

$$[v_1, v_2] = v_1 \cdot v_2 - v_2 \cdot v_1. \quad (2)$$

Then V is a Lie algebra with respect to $[-, -]$.

Examples of Lie Algebra

Example 3

Let M be a manifold, and $\mathfrak{X}(M)$ be the space of smooth vector fields on M . Given two vector fields X and Y , XY is not a vector field, in general. However, $XY - YX$ is a vector field.

Examples of Lie Algebra

Example 3

Let M be a manifold, and $\mathfrak{X}(M)$ be the space of smooth vector fields on M . Given two vector fields X and Y , XY is not a vector field, in general. However, $XY - YX$ is a vector field. Now we define the Lie bracket on $\mathfrak{X}(M)$ as

$$[X, Y] = XY - YX, \quad (3)$$

and then $\mathfrak{X}(M)$ becomes a Lie algebra with respect to $[-, -]$.

Examples of Lie Algebra

Example 3

Let M be a manifold, and $\mathfrak{X}(M)$ be the space of smooth vector fields on M . Given two vector fields X and Y , XY is not a vector field, in general. However, $XY - YX$ is a vector field. Now we define the Lie bracket on $\mathfrak{X}(M)$ as

$$[X, Y] = XY - YX, \quad (3)$$

and then $\mathfrak{X}(M)$ becomes a Lie algebra with respect to $[-, -]$.

Note that $\mathfrak{X}(M)$ is, in general, an infinite dimensional Lie algebra. For example, if we take $M = \mathbb{R}$, then

$$\frac{d}{dx}, x \frac{d}{dx}, x^2 \frac{d}{dx}, x^3 \frac{d}{dx}, \dots$$

are all linearly independent vector fields on \mathbb{R} .

Examples of Lie Algebra

For a Lie group, we can get a finite dimensional subalgebra of $\mathfrak{X}(G)$. It's the space of all left-invariant vector fields on G .

Examples of Lie Algebra

For a Lie group, we can get a finite dimensional subalgebra of $\mathfrak{X}(G)$. It's the space of all left-invariant vector fields on G .

Definition 3

A vector field $X \in \mathfrak{X}(G)$ is called **left-invariant** if it is L_g -invariant for every $g \in G$, i.e. $(L_g)_*X = X$. Then $\mathfrak{L}(G)$ denotes the space of all left-invariant vector fields.

Left-invariant Vector Field

Theorem 4

$\mathfrak{L}(G)$ is a Lie algebra with respect to the Lie bracket $[X, Y] = XY - YX$.

Proof.

See any introductory Lie groups textbook. (Don't ask a toddler on the street please.) ■

Left-invariant Vector Field

Theorem 4

$\mathfrak{L}(G)$ is a Lie algebra with respect to the Lie bracket $[X, Y] = XY - YX$.

Proof.

See any introductory Lie groups textbook. (Don't ask a toddler on the street please.)

"Mathematics should be accessible to everyone."

Meanwhile, mathematicians:

Lemma 5.1. $\text{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/m\mathbb{Z})^\times$.

Proof. Ask a toddler on the street.

Left-invariant Vector Field

Theorem 5

$\mathfrak{L}(G)$ is finite dimensional. Moreover, $\mathfrak{L}(G) \cong T_e G$.

Proof.

Again, see any introductory Lie groups textbook. ■

Lie Algebra Associated to a Lie Group

If G is a Lie group, then $\mathfrak{g} = \mathfrak{L}(G)$ (or equivalently, $\mathfrak{g} = T_e G$) is called the Lie algebra associated to G .

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem

Exponential map

The exponential map is a way of going from the Lie algebra \mathfrak{g} to the Lie group G . So, it's a map

$$\exp : \mathfrak{g} \rightarrow G.$$

Exponential map

The exponential map is a way of going from the Lie algebra \mathfrak{g} to the Lie group G . So, it's a map

$$\exp : \mathfrak{g} \rightarrow G.$$

Given $X \in \mathfrak{g} = \mathfrak{L}(G) \cong T_e G$, let $\alpha : \mathbb{R} \rightarrow G$ be the integral curve of X starting at the identity, i.e. $\alpha(0) = e$. Then we define

$$\exp(X) = \alpha(1). \tag{4}$$

Examples of Exponential Map

Example 6

- For $G = \mathbb{R}_{>0}$, $\mathfrak{g} = T_e G = T_1 \mathbb{R}_{>0} \cong \mathbb{R}$. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is the usual exponential map

$$\begin{aligned}\exp : \mathbb{R} &\rightarrow \mathbb{R}_{>0} \\ t &\mapsto e^t.\end{aligned}\tag{5}$$

Examples of Exponential Map

Example 6

- For $G = \mathbb{R}_{>0}$, $\mathfrak{g} = T_e G = T_1 \mathbb{R}_{>0} \cong \mathbb{R}$. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is the usual exponential map

$$\begin{aligned}\exp : \mathbb{R} &\rightarrow \mathbb{R}_{>0} \\ t &\mapsto e^t.\end{aligned}\tag{5}$$

- For $G = \mathrm{GL}(n, \mathbb{R})$, $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{R}) = \mathrm{End}(\mathbb{R}^n) = \mathbb{R}^{n \times n}$. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is the matrix exponential

$$\begin{aligned}\exp : \mathbb{R}^{n \times n} &\rightarrow \mathrm{GL}(n, \mathbb{R}) \\ A &\mapsto e^A = \sum_{n=0}^{\infty} \frac{1}{n!} A^n.\end{aligned}\tag{6}$$

Examples of Exponential Map

Example 7

- For $G = S^1$, $\mathfrak{g} = T_1 S^1 \cong \mathbb{R}$. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is

$$\begin{aligned}\exp : \mathbb{R} &\rightarrow S^1 \\ t &\mapsto e^{it}.\end{aligned}\tag{7}$$

Examples of Exponential Map

Example 7

- For $G = S^1$, $\mathfrak{g} = T_1 S^1 \cong \mathbb{R}$. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is

$$\begin{aligned}\exp : \mathbb{R} &\rightarrow S^1 \\ t &\mapsto e^{it}.\end{aligned}\tag{7}$$

- For $G = \mathrm{SL}(n, \mathbb{R})$, $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{R}) =$ all traceless $n \times n$ matrices. Then the exponential map $\exp : \mathfrak{g} \rightarrow G$ is the matrix exponential

$$\begin{aligned}\exp : \mathfrak{sl}(n, \mathbb{R}) &\rightarrow \mathrm{SL}(n, \mathbb{R}) \\ A &\mapsto e^A = \sum_{n=0}^{\infty} \frac{1}{n!} A^n.\end{aligned}\tag{8}$$

Properties of Exponential Map

Proposition 8

If $f : G \rightarrow H$ is a Lie group homomorphism, then $(df)_e : T_e G \rightarrow T_e H$ is a Lie algebra homomorphism. Furthermore, the following diagram commutes:

$$\begin{array}{ccc} T_e G & \xrightarrow{(df)_e} & T_e H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (9)$$

Properties of Exponential Map

Proposition 9

$(d \exp)_0 : T_0(T_e G) \cong T_e G \rightarrow T_e G$ is the identity map.

Properties of Exponential Map

Proposition 9

$(d \exp)_0 : T_0(T_e G) \cong T_e G \rightarrow T_e G$ is the identity map.

Therefore, \exp is a local diffeomorphism near $\mathbf{0} \in T_e G$,

Properties of Exponential Map

Proposition 9

$(d \exp)_0 : T_0(T_e G) \cong T_e G \rightarrow T_e G$ is the identity map.

Therefore, \exp is a local diffeomorphism near $\mathbf{0} \in T_e G$, i.e. there is a neighborhood $U \ni e$ and a neighborhood $\mathfrak{u} \ni \mathbf{0}$ such that \exp is a diffeomorphism of \mathfrak{u} and U . Let's write $\log = \exp^{-1}$.

$$\begin{array}{ccc} & \xrightarrow{\exp} & \\ \mathfrak{u} \subseteq \mathfrak{g} & & U \subseteq G \\ & \xleftarrow{\log} & \end{array}$$

Properties of Exponential Map

Is $\exp : \mathfrak{g} \rightarrow G$ a group homomorphism? Does it hold that $\exp(X + Y) = \exp(X)\exp(Y)$?

Properties of Exponential Map

Is $\exp : \mathfrak{g} \rightarrow G$ a group homomorphism? Does it hold that $\exp(X + Y) = \exp(X)\exp(Y)$?

It sure does hold for $G = \mathbb{R}_{>0}, S^1$. In fact, one can show that it holds for every abelian Lie group (for a proof, please go through [7]).

Properties of Exponential Map

Is $\exp : \mathfrak{g} \rightarrow G$ a group homomorphism? Does it hold that $\exp(X + Y) = \exp(X)\exp(Y)$?

It sure does hold for $G = \mathbb{R}_{>0}, S^1$. In fact, one can show that it holds for every abelian Lie group (for a proof, please go through [7]).

But it's not true for non-abelian Lie groups, for example $GL(n, \mathbb{R})$ or $SL(n, \mathbb{R})$.

Properties of Exponential Map

Is $\exp : \mathfrak{g} \rightarrow G$ a group homomorphism? Does it hold that $\exp(X + Y) = \exp(X)\exp(Y)$?

It sure does hold for $G = \mathbb{R}_{>0}, S^1$. In fact, one can show that it holds for every abelian Lie group (for a proof, please go through [7]).

But it's not true for non-abelian Lie groups, for example $GL(n, \mathbb{R})$ or $SL(n, \mathbb{R})$.

To what extent does $\exp : \mathfrak{g} \rightarrow G$ fail to be a group homomorphism for nonabelian G ?

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula**
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem

Proposition 10

If $X, Y \in \mathfrak{g}$ with $[X, Y] = 0$,

$$\exp(X) \exp(Y) = \exp(X + Y).$$

BCH Formula

Theorem 11 (BCH Formula)

For “small” $X, Y \in \mathfrak{g}$,

$$\exp(X)\exp(Y) = \exp\left(X + Y + \frac{1}{2}[X, Y] + \frac{1}{12}([X, [X, Y]] + [Y, [Y, X]]) + \cdots\right)$$

The latter terms are nested commutators. Here, “small” $X, Y \in \mathfrak{g}$ means that there exists a neighborhood \mathfrak{u} of $\mathbf{0} \in \mathfrak{g}$ such that for $X, Y \in \mathfrak{u}$, the RHS converges, and the equation above holds.

BCH Formula

The exact coefficients, or the exact terms of the BCH formula have been computed. The following is taken from [3].

Dynkin's formula:

$$(1.7.3) \quad \mu(X, Y) = Y + X + \sum_{k=1}^{\infty} \frac{(-1)^k}{k+1} \sum_{\substack{l_1, \dots, l_k \geq 0, \\ m_1, \dots, m_k \geq 0, \\ l_j + m_j > 0}} \frac{1}{l_1 + \dots + l_k + 1} \\ \times \left(\frac{(\operatorname{ad} X)^{l_1}}{l_1!} \circ \frac{(\operatorname{ad} Y)^{m_1}}{m_1!} \circ \dots \circ \frac{(\operatorname{ad} X)^{l_k}}{l_k!} \circ \frac{(\operatorname{ad} Y)^{m_k}}{m_k!} \right) (X).$$

BCH Formula

The exact coefficients, or the exact terms of the BCH formula have been computed. The following is taken from [3].

Dynkin's formula:

$$(1.7.3) \quad \mu(X, Y) = Y + X + \sum_{k=1}^{\infty} \frac{(-1)^k}{k+1} \sum_{\substack{l_1, \dots, l_k \geq 0, \\ m_1, \dots, m_k \geq 0, \\ l_j + m_j > 0}} \frac{1}{l_1 + \dots + l_k + 1} \\ \times \left(\frac{(\operatorname{ad} X)^{l_1}}{l_1!} \circ \frac{(\operatorname{ad} Y)^{m_1}}{m_1!} \circ \dots \circ \frac{(\operatorname{ad} X)^{l_k}}{l_k!} \circ \frac{(\operatorname{ad} Y)^{m_k}}{m_k!} \right) (X).$$

But the exact formula is rarely useful. What's useful is the mere fact that this formula exists!

BCH Formula

The exact coefficients, or the exact terms of the BCH formula have been computed. The following is taken from [3].

Dynkin's formula:

$$(1.7.3) \quad \mu(X, Y) = Y + X + \sum_{k=1}^{\infty} \frac{(-1)^k}{k+1} \sum_{\substack{l_1, \dots, l_k \geq 0, \\ m_1, \dots, m_k \geq 0, \\ l_j + m_j > 0}} \frac{1}{l_1 + \dots + l_k + 1} \\ \times \left(\frac{(\operatorname{ad} X)^{l_1}}{l_1!} \circ \frac{(\operatorname{ad} Y)^{m_1}}{m_1!} \circ \dots \circ \frac{(\operatorname{ad} X)^{l_k}}{l_k!} \circ \frac{(\operatorname{ad} Y)^{m_k}}{m_k!} \right) (X).$$

But the exact formula is rarely useful. What's useful is the mere fact that this formula exists! Because it gives a way of multiplying Lie group elements (near the identity) just from the Lie bracket.

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?**
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem

Lie Algebra to Lie Group

We already know that if $f : G \rightarrow H$ is a Lie group homomorphism, then $(df)_e : T_e G \rightarrow T_e H$ is a Lie algebra homomorphism, and the following diagram commutes:

$$\begin{array}{ccc} T_e G & \xrightarrow{(df)_e} & T_e H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (10)$$

Lie Algebra to Lie Group

We already know that if $f : G \rightarrow H$ is a Lie group homomorphism, then $(df)_e : T_e G \rightarrow T_e H$ is a Lie algebra homomorphism, and the following diagram commutes:

$$\begin{array}{ccc} T_e G & \xrightarrow{(df)_e} & T_e H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (10)$$

Now we want to know whether the converse holds, i.e. given any Lie algebra homomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$, is there a corresponding Lie group homomorphism $f : G \rightarrow H$ such that $\Phi = (df)_e$?

Lie Algebra to Lie Group

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (11)$$

We want to define something like $f(\exp(X)) = \exp(\Phi(X))$. But it has several issues:

Lie Algebra to Lie Group

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (11)$$

We want to define something like $f(\exp(X)) = \exp(\Phi(X))$. But it has several issues:

- Can every element $g \in G$ be written as $\exp X$ for some $X \in \mathfrak{g}$?

Lie Algebra to Lie Group

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (11)$$

We want to define something like $f(\exp(X)) = \exp(\Phi(X))$. But it has several issues:

- Can every element $g \in G$ be written as $\exp X$ for some $X \in \mathfrak{g}$?
- If $g = \exp(X)$, is this unique? If not, why is $\exp(\Phi(X))$ independent of the choice of X ?

Lie Algebra to Lie Group

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (11)$$

We want to define something like $f(\exp(X)) = \exp(\Phi(X))$. But it has several issues:

- Can every element $g \in G$ be written as $\exp X$ for some $X \in \mathfrak{g}$?
- If $g = \exp(X)$, is this unique? If not, why is $\exp(\Phi(X))$ independent of the choice of X ?
- Is f a group homomorphism?

Lie Algebra to Lie Group

Consider $G = S^1$ and $H = \mathbb{R}$. Both of them have isomorphic Lie algebras, $\mathfrak{g} = \mathfrak{h} = \mathbb{R}$. So there is a Lie algebra isomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$. But there is no corresponding nontrivial Lie group homomorphism. Assume there is one, say ϕ .

$$\begin{array}{ccc} \mathfrak{g} = \mathbb{R} & \xrightarrow{\Phi} & \mathfrak{h} = \mathbb{R} \\ \exp_{S^1} \downarrow & & \downarrow \exp_{\mathbb{R}} \\ S^1 & \xrightarrow{\phi} & \mathbb{R} \end{array} \quad (12)$$

Lie Algebra to Lie Group

Consider $G = S^1$ and $H = \mathbb{R}$. Both of them have isomorphic Lie algebras, $\mathfrak{g} = \mathfrak{h} = \mathbb{R}$. So there is a Lie algebra isomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$. But there is no corresponding nontrivial Lie group homomorphism. Assume there is one, say ϕ .

$$\begin{array}{ccc} \mathfrak{g} = \mathbb{R} & \xrightarrow{\Phi} & \mathfrak{h} = \mathbb{R} \\ \exp_{S^1} \downarrow & & \downarrow \exp_{\mathbb{R}} \\ S^1 & \xrightarrow{\phi} & \mathbb{R} \end{array} \quad (12)$$

Suppose $\phi(e^{i\theta}) = x \neq 0$. Then

$$\phi(e^{in\theta}) = nx, \text{ for any } n \in \mathbb{Z}.$$

Lie Algebra to Lie Group

Consider $G = S^1$ and $H = \mathbb{R}$. Both of them have isomorphic Lie algebras, $\mathfrak{g} = \mathfrak{h} = \mathbb{R}$. So there is a Lie algebra isomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$. But there is no corresponding nontrivial Lie group homomorphism. Assume there is one, say ϕ .

$$\begin{array}{ccc} \mathfrak{g} = \mathbb{R} & \xrightarrow{\Phi} & \mathfrak{h} = \mathbb{R} \\ \exp_{S^1} \downarrow & & \downarrow \exp_{\mathbb{R}} \\ S^1 & \xrightarrow{\phi} & \mathbb{R} \end{array} \quad (12)$$

Suppose $\phi(e^{i\theta}) = x \neq 0$. Then

$$\phi(e^{in\theta}) = nx, \text{ for any } n \in \mathbb{Z}.$$

But since S^1 is compact, $\phi(S^1)$ needs to be compact, i.e. closed and bounded. But it's not bounded if $x \neq 0$. Therefore, the only Lie group homomorphism $\phi : S^1 \rightarrow \mathbb{R}$ is the trivial one. But its corresponding Lie algebra homomorphism is also trivial, NOT an isomorphism.

Lie Algebra to Lie Group

Therefore, the group homomorphism $f : G \rightarrow H$ cannot always be defined. We need some extra condition(s).

Lie Algebra to Lie Group

Therefore, the group homomorphism $f : G \rightarrow H$ cannot always be defined. We need some extra condition(s).

Notice that S^1 has a “hole”, and \mathbb{R} doesn't. More concretely, S^1 is **NOT** simply connected, since

$$\pi_1(S^1) = \mathbb{Z}. \quad (13)$$

Lie Algebra to Lie Group

Therefore, the group homomorphism $f : G \rightarrow H$ cannot always be defined. We need some extra condition(s).

Notice that S^1 has a “hole”, and \mathbb{R} doesn't. More concretely, S^1 is **NOT** simply connected, since

$$\pi_1(S^1) = \mathbb{Z}. \quad (13)$$

So we can guess that for simply connected Lie group G , such a group homomorphism $f : G \rightarrow H$ exists for every Lie group homomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$.

Lie Algebra to Lie Group

Therefore, the group homomorphism $f : G \rightarrow H$ cannot always be defined. We need some extra condition(s).

Notice that S^1 has a “hole”, and \mathbb{R} doesn't. More concretely, S^1 is **NOT** simply connected, since

$$\pi_1(S^1) = \mathbb{Z}. \quad (13)$$

So we can guess that for simply connected Lie group G , such a group homomorphism $f : G \rightarrow H$ exists for every Lie group homomorphism $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$.

Turns out, this is actually correct!!

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group**
- 6 Lie's Third Theorem

Simply Connected Lie Group

How does simply connectedness help us?

Simply Connected Lie Group

How does simply connectedness help us?

Essentially, what we do is:

Simply Connected Lie Group

How does simply connectedness help us?

Essentially, what we do is:

- 1 We define $f(\exp(X)) = \exp(\Phi(X))$ on a neighborhood of $e \in G$.

Simply Connected Lie Group

How does simply connectedness help us?

Essentially, what we do is:

- 1 We define $f(\exp(X)) = \exp(\Phi(X))$ on a neighborhood of $e \in G$.
- 2 We extend this definition by taking **ANY** path from $e \in G$ to $g \in G$, and then define $f(g)$.

Simply Connected Lie Group

How does simply connectedness help us?

Essentially, what we do is:

- 1 We define $f(\exp(X)) = \exp(\Phi(X))$ on a neighborhood of $e \in G$.
- 2 We extend this definition by taking **ANY** path from $e \in G$ to $g \in G$, and then define $f(g)$.
- 3 Then we show that this is independent of the path we choose, since any two paths are homotopic, because G is simply connected.

Step 1

We take a neighborhood $\mathfrak{u} \subseteq \mathfrak{g}$ and $\mathfrak{v} \subseteq \mathfrak{h}$ such that $\Phi(\mathfrak{u}) \subseteq \mathfrak{v}$, and BCH formula holds on \mathfrak{u} and \mathfrak{v} . Since \exp is a local diffeomorphism, it's a diffeomorphism of \mathfrak{u} and $\exp(\mathfrak{u}) =: U$.

Step 1

We take a neighborhood $\mathfrak{u} \subseteq \mathfrak{g}$ and $\mathfrak{v} \subseteq \mathfrak{h}$ such that $\Phi(\mathfrak{u}) \subseteq \mathfrak{v}$, and BCH formula holds on \mathfrak{u} and \mathfrak{v} . Since \exp is a local diffeomorphism, it's a diffeomorphism of \mathfrak{u} and $\exp(\mathfrak{u}) =: U$.

So, on $U \subseteq G$, we can uniquely define $f(\exp(X)) = \exp(\Phi(X))$. Then why is f a (locally) group homomorphism?

Step 1

We take a neighborhood $\mathfrak{u} \subseteq \mathfrak{g}$ and $\mathfrak{v} \subseteq \mathfrak{h}$ such that $\Phi(\mathfrak{u}) \subseteq \mathfrak{v}$, and BCH formula holds on \mathfrak{u} and \mathfrak{v} . Since \exp is a local diffeomorphism, it's a diffeomorphism of \mathfrak{u} and $\exp(\mathfrak{u}) =: U$.

So, on $U \subseteq G$, we can uniquely define $f(\exp(X)) = \exp(\Phi(X))$. Then why is f a (locally) group homomorphism?

The answer is BCH formula!!

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

$$f(\exp(X)\exp(Y)) = f\left(\exp\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right)$$

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

$$\begin{aligned} f(\exp(X)\exp(Y)) &= f\left(\exp\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \end{aligned}$$

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

$$\begin{aligned} f(\exp(X)\exp(Y)) &= f\left(\exp\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi(X) + \Phi(Y) + \frac{1}{2}[\Phi(X), \Phi(Y)] + \dots\right) \end{aligned}$$

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

$$\begin{aligned} f(\exp(X)\exp(Y)) &= f\left(\exp\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi(X) + \Phi(Y) + \frac{1}{2}[\Phi(X), \Phi(Y)] + \dots\right) \\ &= \exp(\Phi(X))\exp(\Phi(Y)) \end{aligned}$$

Step 1

We want to show that $f(\exp(X)\exp(Y)) = f(\exp X)f(\exp Y)$.

$$\begin{aligned} f(\exp(X)\exp(Y)) &= f\left(\exp\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi\left(X + Y + \frac{1}{2}[X, Y] + \dots\right)\right) \\ &= \exp\left(\Phi(X) + \Phi(Y) + \frac{1}{2}[\Phi(X), \Phi(Y)] + \dots\right) \\ &= \exp(\Phi(X))\exp(\Phi(Y)) \\ &= f(\exp X)f(\exp Y). \end{aligned}$$

So f acts like a group homomorphism on U .

Step 2: Define $f(g)$

Now take any $g \in G$. Consider any path $\gamma : [0, 1] \rightarrow G$ from e to g .

Step 2: Define $f(g)$

Now take any $g \in G$. Consider any path $\gamma : [0, 1] \rightarrow G$ from e to g .

Since $[0, 1]$ is compact, there is a partition $0 = t_0 < t_1 < t_2 < \cdots < t_m = 1$ such that for all $s, t \in [t_i, t_{i+1}]$ and for each i ,

$$\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U. \quad (14)$$

We call such a partition a “*good partition*” for γ .

Step 2: Define $f(g)$

Now take any $g \in G$. Consider any path $\gamma : [0, 1] \rightarrow G$ from e to g . Since $[0, 1]$ is compact, there is a partition $0 = t_0 < t_1 < t_2 < \cdots < t_m = 1$ such that for all $s, t \in [t_i, t_{i+1}]$ and for each i ,

$$\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U. \quad (14)$$

We call such a partition a “good partition” for γ . Then we write

$$g = (\gamma(t_m)\gamma(t_{m-1})^{-1}) (\gamma(t_{m-1})\gamma(t_{m-2})^{-1}) \cdots (\gamma(t_2)\gamma(t_1)^{-1}) (\gamma(t_1)\gamma(t_0)^{-1}).$$

Then we define

$$\begin{aligned} f(g) = & f(\gamma(t_m)\gamma(t_{m-1})^{-1}) f(\gamma(t_{m-1})\gamma(t_{m-2})^{-1}) \cdots \\ & \cdots f(\gamma(t_2)\gamma(t_1)^{-1}) f(\gamma(t_1)\gamma(t_0)^{-1}). \end{aligned} \quad (15)$$

Step 2: Independence of Partition

There can be another partition of the interval $[0, 1]$ such that for each i , for all $s, t \in [t'_i, t'_{i+1}]$, $\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U$. But $f(g)$ stays invariant!

Step 2: Independence of Partition

There can be another partition of the interval $[0, 1]$ such that for each i , for all $s, t \in [t'_i, t'_{i+1}]$, $\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U$. But $f(g)$ stays invariant! Let's say we add a point s in between t_i and t_{i+1} . Then the value of $f(g)$ with respect to this refined partition is

$$f(g) = \cdots f(\gamma(t_{i+1})\gamma(s)^{-1}) f(\gamma(s)\gamma(t_i)^{-1}) \cdots$$

Step 2: Independence of Partition

There can be another partition of the interval $[0, 1]$ such that for each i , for all $s, t \in [t'_i, t'_{i+1}]$, $\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U$. But $f(g)$ stays invariant! Let's say we add a point s in between t_i and t_{i+1} . Then the value of $f(g)$ with respect to this refined partition is

$$\begin{aligned} f(g) &= \cdots f(\gamma(t_{i+1})\gamma(s)^{-1}) f(\gamma(s)\gamma(t_i)^{-1}) \cdots \\ &= \cdots f(\gamma(t_{i+1})\gamma(t_i)^{-1}) \cdots \end{aligned}$$

which is the same as the value obtained from the previous partition! Therefore, $f(g)$ is invariant under refinement of partition.

Step 2: Independence of Partition

There can be another partition of the interval $[0, 1]$ such that for each i , for all $s, t \in [t'_i, t'_{i+1}]$, $\gamma(t)\gamma(s)^{-1} \in \exp(\mathfrak{u}) = U$. But $f(g)$ stays invariant! Let's say we add a point s in between t_i and t_{i+1} . Then the value of $f(g)$ with respect to this refined partition is

$$\begin{aligned} f(g) &= \cdots f(\gamma(t_{i+1})\gamma(s)^{-1}) f(\gamma(s)\gamma(t_i)^{-1}) \cdots \\ &= \cdots f(\gamma(t_{i+1})\gamma(t_i)^{-1}) \cdots \end{aligned}$$

which is the same as the value obtained from the previous partition! Therefore, $f(g)$ is invariant under refinement of partition. Given two partitions, one may take their union, which is refinement of both partitions, and thus we achieve the independence of partition!

Step 3: Independence of Path

Simply-connectedness comes into play here!

Step 3: Independence of Path

Simply-connectedness comes into play here! Suppose β, γ are two paths from e to g . Then there is a homotopy $H : [0, 1]^2 \rightarrow G$ such that

$$H(0, -) = \beta, \quad H(1, -) = \gamma; \quad (16)$$

$$H(-, 0) = e, \quad H(-, 1) = g. \quad (17)$$

Step 3: Independence of Path

Simply-connectedness comes into play here! Suppose β, γ are two paths from e to g . Then there is a homotopy $H : [0, 1]^2 \rightarrow G$ such that

$$H(0, -) = \beta, \quad H(1, -) = \gamma; \quad (16)$$

$$H(-, 0) = e, \quad H(-, 1) = g. \quad (17)$$


Since $[0, 1]^2$ is compact, there exists a positive integer N such that for every $(s, t), (s', t') \in [0, 1]^2$ with $|s - s'| < \frac{2}{N}, |t - t'| < \frac{2}{N}$,

$$H(s, t) H(s', t')^{-1} \in \exp(\mathfrak{u}) = U. \quad (18)$$

Step 3: Independence of Path

Now we define a sequence of paths $\alpha_{k,l}$ for $k = 0, 1, 2, \dots, N-1$ and $l = 0, 1, 2, \dots, N$:

$$\alpha_{k,l}(t) = \begin{cases} H\left(\frac{k+1}{N}, t\right) & \text{if } 0 \leq t \leq \frac{l-1}{N}, \\ H\left(\frac{k}{N}, t\right) & \text{if } \frac{l-1}{N} \leq t \leq 1, \\ H(\text{connecting diagonal}) & \text{if } \frac{l-1}{N} \leq t \leq \frac{l}{N}. \end{cases} \quad (19)$$



Step 3: Independence of Path

We will show that the value of $f(g)$ is the same across all $\alpha_{k,l}$'s. Then we're done, since $\alpha_{0,0} = \beta$ and $\alpha_{N,0} = \gamma$.

Step 3: Independence of Path

We will show that the value of $f(g)$ is the same across all $\alpha_{k,l}$'s. Then we're done, since $\alpha_{0,0} = \beta$ and $\alpha_{N,0} = \gamma$.

Note that $\alpha_{k,l}$ and $\alpha_{k,l+1}$ agree everywhere except on $[\frac{l-1}{N}, \frac{l+1}{N}]$. We consider the partition

$$0, \frac{1}{N}, \dots, \frac{l-1}{N}, \frac{l+1}{N}, \frac{l+2}{N}, \dots, 1.$$

This is a good partition for both $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

Step 3: Independence of Path

We will show that the value of $f(g)$ is the same across all $\alpha_{k,l}$'s. Then we're done, since $\alpha_{0,0} = \beta$ and $\alpha_{N,0} = \gamma$.

Note that $\alpha_{k,l}$ and $\alpha_{k,l+1}$ agree everywhere except on $[\frac{l-1}{N}, \frac{l+1}{N}]$. We consider the partition

$$0, \frac{1}{N}, \dots, \frac{l-1}{N}, \frac{l+1}{N}, \frac{l+2}{N}, \dots, 1.$$

This is a good partition for both $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

Since the value of $f(g)$ across a path depends only on the points of the partition, the value of $f(g)$ is the same across $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

Step 3: Independence of Path

We will show that the value of $f(g)$ is the same across all $\alpha_{k,l}$'s. Then we're done, since $\alpha_{0,0} = \beta$ and $\alpha_{N,0} = \gamma$.

Note that $\alpha_{k,l}$ and $\alpha_{k,l+1}$ agree everywhere except on $[\frac{l-1}{N}, \frac{l+1}{N}]$. We consider the partition

$$0, \frac{1}{N}, \dots, \frac{l-1}{N}, \frac{l+1}{N}, \frac{l+2}{N}, \dots, 1.$$

This is a good partition for both $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

Since the value of $f(g)$ across a path depends only on the points of the partition, the value of $f(g)$ is the same across $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

A similar argument shows that the value of $f(g)$ is the same across $\alpha_{k,N}$ and $\alpha_{k+1,0}$.

Step 3: Independence of Path

We will show that the value of $f(g)$ is the same across all $\alpha_{k,l}$'s. Then we're done, since $\alpha_{0,0} = \beta$ and $\alpha_{N,0} = \gamma$.

Note that $\alpha_{k,l}$ and $\alpha_{k,l+1}$ agree everywhere except on $[\frac{l-1}{N}, \frac{l+1}{N}]$. We consider the partition

$$0, \frac{1}{N}, \dots, \frac{l-1}{N}, \frac{l+1}{N}, \frac{l+2}{N}, \dots, 1.$$

This is a good partition for both $\alpha_{k,l}$ and $\alpha_{k,l+1}$.


Since the value of $f(g)$ across a path depends only on the points of the partition, the value of $f(g)$ is the same across $\alpha_{k,l}$ and $\alpha_{k,l+1}$.

A similar argument shows that the value of $f(g)$ is the same across $\alpha_{k,N}$ and $\alpha_{k+1,0}$.

Therefore, $f(g)$ is well-defined.

Step 4: f is a group homomorphism

Let β and γ be paths from e to g and h , respectively. Then from that, we can construct a path from e to gh in the following way:



Step 4: f is a group homomorphism

$$\alpha(t) = \begin{cases} \beta(2t) & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g \cdot \gamma(2t - 1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases} \quad (20)$$

We then combine the good partitions for γ and β to find a good partition for α .

Step 4: f is a group homomorphism

$$\alpha(t) = \begin{cases} \beta(2t) & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g \cdot \gamma(2t - 1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases} \quad (20)$$

We then combine the good partitions for γ and β to find a good partition for α . If $t_0 < \dots < t_m$ and $s_0 < \dots < s_n$ are the good partitions for γ and β , a good partition for α will be

$$\frac{t_0}{2} < \frac{t_1}{2} < \dots < \frac{t_m}{2} = \frac{s_0 + 1}{2} < \frac{s_1 + 1}{2} < \dots < \frac{s_n + 1}{2}.$$

Step 4: f is a group homomorphism

$$\alpha(t) = \begin{cases} \beta(2t) & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g \cdot \gamma(2t-1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases} \quad (20)$$

We then combine the good partitions for γ and β to find a good partition for α . If $t_0 < \dots < t_m$ and $s_0 < \dots < s_n$ are the good partitions for γ and β , a good partition for α will be

$$\frac{t_0}{2} < \frac{t_1}{2} < \dots < \frac{t_m}{2} = \frac{s_0+1}{2} < \frac{s_1+1}{2} < \dots < \frac{s_n+1}{2}.$$

$$f(gh) = \prod_{i=n}^1 f\left(\alpha\left(\frac{s_i+1}{2}\right) \alpha\left(\frac{s_{i-1}+1}{2}\right)^{-1}\right) \prod_{j=m}^1 f\left(\alpha\left(\frac{t_j}{2}\right) \alpha\left(\frac{t_{j-1}}{2}\right)^{-1}\right)$$

Step 4: f is a group homomorphism

$$\alpha(t) = \begin{cases} \beta(2t) & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g \cdot \gamma(2t-1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases} \quad (20)$$

We then combine the good partitions for γ and β to find a good partition for α . If $t_0 < \dots < t_m$ and $s_0 < \dots < s_n$ are the good partitions for γ and β , a good partition for α will be

$$\frac{t_0}{2} < \frac{t_1}{2} < \dots < \frac{t_m}{2} = \frac{s_0 + 1}{2} < \frac{s_1 + 1}{2} < \dots < \frac{s_n + 1}{2}.$$

$$\begin{aligned} f(gh) &= \prod_{i=n}^1 f \left(\alpha \left(\frac{s_i + 1}{2} \right) \alpha \left(\frac{s_{i-1} + 1}{2} \right)^{-1} \right) \prod_{j=m}^1 f \left(\alpha \left(\frac{t_j}{2} \right) \alpha \left(\frac{t_{j-1}}{2} \right)^{-1} \right) \\ &= \prod_{i=n}^1 f(g\gamma(s_i)\gamma(s_{i-1})^{-1}g^{-1}) \prod_{j=m}^1 f(\beta(t_j)\beta(t_{j-1})^{-1}) \end{aligned}$$

Step 4: f is a group homomorphism

$$\alpha(t) = \begin{cases} \beta(2t) & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g \cdot \gamma(2t-1) & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases} \quad (20)$$

We then combine the good partitions for γ and β to find a good partition for α . If $t_0 < \dots < t_m$ and $s_0 < \dots < s_n$ are the good partitions for γ and β , a good partition for α will be

$$\frac{t_0}{2} < \frac{t_1}{2} < \dots < \frac{t_m}{2} = \frac{s_0+1}{2} < \frac{s_1+1}{2} < \dots < \frac{s_n+1}{2}.$$

$$\begin{aligned} f(gh) &= \prod_{i=n}^1 f \left(\alpha \left(\frac{s_i+1}{2} \right) \alpha \left(\frac{s_{i-1}+1}{2} \right)^{-1} \right) \prod_{j=m}^1 f \left(\alpha \left(\frac{t_j}{2} \right) \alpha \left(\frac{t_{j-1}}{2} \right)^{-1} \right) \\ &= \prod_{i=n}^1 f(g\gamma(s_i)\gamma(s_{i-1})^{-1}g^{-1}) \prod_{j=m}^1 f(\beta(t_j)\beta(t_{j-1})^{-1}) \\ &= f(ghg^{-1})f(g). \end{aligned}$$

Step 4: f is a group homomorphism

So we have $f(gh) = f(ghg^{-1})f(g)$ for every $g, h \in G$.

Step 4: f is a group homomorphism

So we have $f(gh) = f(ghg^{-1})f(g)$ for every $g, h \in G$. Given any $a, b \in G$, choose $g = b$ and $h = b^{-1}ab$.

Step 4: f is a group homomorphism

So we have $f(gh) = f(ghg^{-1})f(g)$ for every $g, h \in G$. Given any $a, b \in G$, choose $g = b$ and $h = b^{-1}ab$.

$$f(gh) = f(b \cdot b^{-1}ab) = f(ab) \quad (21)$$

Step 4: f is a group homomorphism

So we have $f(gh) = f(ghg^{-1})f(g)$ for every $g, h \in G$. Given any $a, b \in G$, choose $g = b$ and $h = b^{-1}ab$.

$$f(gh) = f(b \cdot b^{-1}ab) = f(ab) \quad (21)$$

$$f(ghg^{-1})f(g) = f(b \cdot b^{-1}ab \cdot b^{-1})f(b) = f(a)f(b). \quad (22)$$

Step 4: f is a group homomorphism

So we have $f(gh) = f(ghg^{-1})f(g)$ for every $g, h \in G$. Given any $a, b \in G$, choose $g = b$ and $h = b^{-1}ab$.

$$f(gh) = f(b \cdot b^{-1}ab) = f(ab) \quad (21)$$

$$f(ghg^{-1})f(g) = f(b \cdot b^{-1}ab \cdot b^{-1})f(b) = f(a)f(b). \quad (22)$$

So f is a group homomorphism.

Step 5: $(df)_e = \Phi$

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array} \quad (23)$$

For $X_e \in T_e G$,

$$(df)_e X_e = \left. \frac{d}{dt} \right|_{t=0} f(\exp(tX_e))$$

.

Step 5: $(df)_e = \Phi$

$$\begin{array}{ccc}
 T_e G & \xrightarrow{\Phi} & T_{e'} H \\
 \exp_G \downarrow & & \downarrow \exp_H \\
 G & \xrightarrow{f} & H
 \end{array} \tag{23}$$

For $X_e \in T_e G$,

$$\begin{aligned}
 (df)_e X_e &= \left. \frac{d}{dt} \right|_{t=0} f(\exp(tX_e)) \\
 &= \left. \frac{d}{dt} \right|_{t=0} \exp(\Phi(tX_e))
 \end{aligned}$$

.

Step 5: $(df)_e = \Phi$

$$\begin{array}{ccc}
 T_e G & \xrightarrow{\Phi} & T_{e'} H \\
 \exp_G \downarrow & & \downarrow \exp_H \\
 G & \xrightarrow{f} & H
 \end{array} \tag{23}$$

For $X_e \in T_e G$,

$$\begin{aligned}
 (df)_e X_e &= \left. \frac{d}{dt} \right|_{t=0} f(\exp(tX_e)) \\
 &= \left. \frac{d}{dt} \right|_{t=0} \exp(\Phi(tX_e)) \\
 &= \left. \frac{d}{dt} \right|_{t=0} \exp(t\Phi(X_e)) \\
 & \quad .
 \end{aligned}$$

Step 5: $(df)_e = \Phi$

$$\begin{array}{ccc}
 T_e G & \xrightarrow{\Phi} & T_{e'} H \\
 \exp_G \downarrow & & \downarrow \exp_H \\
 G & \xrightarrow{f} & H
 \end{array} \tag{23}$$

For $X_e \in T_e G$,

$$\begin{aligned}
 (df)_e X_e &= \left. \frac{d}{dt} \right|_{t=0} f(\exp(tX_e)) \\
 &= \left. \frac{d}{dt} \right|_{t=0} \exp(\Phi(tX_e)) \\
 &= \left. \frac{d}{dt} \right|_{t=0} \exp(t\Phi(X_e)) \\
 &= \Phi(X_e).
 \end{aligned}$$

So $\Phi = (df)_e$.

Summary so far

So, we have proved the following:

Theorem 12 (Lie's second theorem)

If G is a simply connected Lie group, and $\Phi : \mathfrak{g} \rightarrow \mathfrak{h}$ is a Lie algebra homomorphism, there is a Lie group homomorphism $f : G \rightarrow H$ such that $\Phi = (df)_e$, and the following diagram commutes:

$$\begin{array}{ccc} T_e G & \xrightarrow{\Phi} & T_{e'} H \\ \exp_G \downarrow & & \downarrow \exp_H \\ G & \xrightarrow{f} & H \end{array}$$

(24)

Table of Contents

- 1 Lie Groups and Lie Algebras Basics
- 2 The Exponential Map
- 3 The Baker–Campbell–Hausdorff (BCH) Formula
- 4 How Well Does Lie Algebra Know Its Lie Group?
- 5 Simply Connected Lie Group
- 6 Lie's Third Theorem**

Question

Given any finite dimensional Lie algebra, is it true that it's the Lie algebra of some Lie group?

Question

Given any finite dimensional Lie algebra, is it true that it's the Lie algebra of some Lie group?

The answer is YES!

Question

Given any finite dimensional Lie algebra, is it true that it's the Lie algebra of some Lie group?

The answer is YES!

Furthermore, the Lie group is **unique** if we require it to be simply connected!

Lie's Third Theorem

Theorem 13 (Lie's Third Theorem)

For any finite dimensional Lie algebra \mathfrak{g} , there is a unique simply connected Lie group G whose Lie algebra is \mathfrak{g} .

Lie's Third Theorem

Theorem 13 (Lie's Third Theorem)

For any finite dimensional Lie algebra \mathfrak{g} , there is a unique simply connected Lie group G whose Lie algebra is \mathfrak{g} .

The proof uses Ado's theorem.

Lie's Third Theorem

Theorem 13 (Lie's Third Theorem)

For any finite dimensional Lie algebra \mathfrak{g} , there is a unique simply connected Lie group G whose Lie algebra is \mathfrak{g} .

The proof uses Ado's theorem.

Theorem 14 (Ado's theorem)

Every finite dimensional real Lie algebra is isomorphic to a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{R})$ for some n .

Proof

By Ado's theorem, \mathfrak{g} is a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{R})$.

Proof

By Ado's theorem, \mathfrak{g} is a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{R})$.

Theorem 15 (Chevalley's theorem, also known as Lie's first theorem)

Let G be a Lie group, and \mathfrak{g} its Lie algebra. Then there is a one-to-one correspondence

$$\{\text{connected Lie subgroups } H \subseteq G\} \leftrightarrow \{\text{Lie subalgebras } \mathfrak{h} \subseteq \mathfrak{g}\}.$$

Proof

By Ado's theorem, \mathfrak{g} is a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{R})$.

Theorem 15 (Chevalley's theorem, also known as Lie's first theorem)

Let G be a Lie group, and \mathfrak{g} its Lie algebra. Then there is a one-to-one correspondence

$$\{\text{connected Lie subgroups } H \subseteq G\} \leftrightarrow \{\text{Lie subalgebras } \mathfrak{h} \subseteq \mathfrak{g}\}.$$

Therefore, there is a connected Lie subgroup of $G' \subseteq \text{GL}(n, \mathbb{R})$ whose Lie algebra is \mathfrak{g} .

By Ado's theorem, \mathfrak{g} is a Lie subalgebra of $\mathfrak{gl}(n, \mathbb{R})$.

Theorem 15 (Chevalley's theorem, also known as Lie's first theorem)

Let G be a Lie group, and \mathfrak{g} its Lie algebra. Then there is a one-to-one correspondence

$$\{\text{connected Lie subgroups } H \subseteq G\} \leftrightarrow \{\text{Lie subalgebras } \mathfrak{h} \subseteq \mathfrak{g}\}.$$

Therefore, there is a connected Lie subgroup of $G' \subseteq \text{GL}(n, \mathbb{R})$ whose Lie algebra is \mathfrak{g} .

Take G to be the simply connected cover of G' . It has a group structure, and its Lie algebra is the same as the Lie algebra of G (details are in [7]).

Proof

Hence, given **any** finite dimensional Lie algebra \mathfrak{g} , there is a simply connected Lie group G whose Lie algebra is \mathfrak{g} . As for the uniqueness, we'll use the following:

Proof

Hence, given **any** finite dimensional Lie algebra \mathfrak{g} , there is a simply connected Lie group G whose Lie algebra is \mathfrak{g} . As for the uniqueness, we'll use the following:

Lemma 16

If G and H are simply connected Lie groups with $\mathfrak{g} \cong \mathfrak{h}$, then $G \cong H$.

Proof

Hence, given **any** finite dimensional Lie algebra \mathfrak{g} , there is a simply connected Lie group G whose Lie algebra is \mathfrak{g} . As for the uniqueness, we'll use the following:

Lemma 16

If G and H are simply connected Lie groups with $\mathfrak{g} \cong \mathfrak{h}$, then $G \cong H$.

Proof.

$$\begin{array}{ccccc} \mathfrak{g} & \xrightarrow{\Phi=(df)_e} & \mathfrak{h} & \xrightarrow{\Phi^{-1}=(df')_{e'}} & \mathfrak{g} \\ \exp_G \downarrow & & \exp_H \downarrow & & \downarrow \exp_G \\ G & \xrightarrow{f} & H & \xrightarrow{f'} & G \end{array}$$

Therefore, $(d(f' \circ f))_e = \mathbb{1}_{\mathfrak{g}}$.

Proof

Hence, given **any** finite dimensional Lie algebra \mathfrak{g} , there is a simply connected Lie group G whose Lie algebra is \mathfrak{g} . As for the uniqueness, we'll use the following:

Lemma 16

If G and H are simply connected Lie groups with $\mathfrak{g} \cong \mathfrak{h}$, then $G \cong H$.

Proof.

$$\begin{array}{ccccc} \mathfrak{g} & \xrightarrow{\Phi=(df)_e} & \mathfrak{h} & \xrightarrow{\Phi^{-1}=(df')_{e'}} & \mathfrak{g} \\ \exp_G \downarrow & & \exp_H \downarrow & & \downarrow \exp_G \\ G & \xrightarrow{f} & H & \xrightarrow{f'} & G \end{array}$$

Therefore, $(d(f' \circ f))_e = \mathbb{1}_{\mathfrak{g}}$. Since any connected Lie group is generated by any neighborhood of identity and \exp is a local diffeomorphism, we have $f' \circ f = \text{id}_G$. Similarly, $f \circ f' = \text{id}_H$. ■

Proof

Therefore, for any finite dimensional Lie algebra \mathfrak{g} , there is a unique simply connected Lie group G whose Lie algebra is \mathfrak{g} .

References

- [1] A. Kirillov Jr., *An Introduction to Lie Groups and Lie Algebras*, Cambridge Studies in Advanced Mathematics, vol. 113, Cambridge University Press, Cambridge, 2008.
- [2] B. C. Hall, *Lie Groups, Lie Algebras, and Representations*, 2nd ed., Graduate Texts in Mathematics, vol. 222, Springer, New York, 2015.
- [3] J. A. C. Kolk and J. J. Duistermaat, *Lie Groups*, Universitext, Springer, Berlin, 2000.
- [4] T. Bröcker and T. tom Dieck, *Representations of Compact Lie Groups*, Springer-Verlag, New York, 1985.
- [5] P. Etingof, *Lie Groups and Lie Algebras*, Lecture notes, MIT, 2020-2021; arXiv:2201.09397 [math.RT], 2022. URL: arxiv.org/pdf/2201.09397.pdf
- [6] C. Daly, *Baker-Campbell-Hausdorff Formula*, available at https://www.math.brown.edu/cdaly2/Notes/BCH_Formula.pdf.
- [7] Expository notes I've written: <https://atonurc.github.io/liegrp>.

Thank you for joining!

The slides are available in my webpage

https://atonurc.github.io/liegrp/BCH_Lie3_Talk.pdf

