MTH-ALG, Algebra Problem Sets Solutions

Atonu Roy Chowdhury, achowdhu@ictp.it

Contents

1 Problem Set 1 2

2 Problem Set 2 11

3 Problems on Groups from Lothar Göttsche's Notes 22

4 Problems on Rings from Lothar Göttsche's Notes 35

5 Problem Set 3 64

6 Problem Set 4 70

§1 Problem Set 1

Exercise 1.1. Let

$$G = \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/20\mathbb{Z}. \tag{1.1}$$

(a) How many elements of G have order 60?

(b) How many cyclic subgroups of G have order 30?

(c) How many injective homomorphisms

$$f : \mathbb{Z}/12\mathbb{Z} \longrightarrow G \tag{1.2}$$

are there?

Solution. (a) We shall prove something first:

> **Claim 1.1:** If $a \in G$ has order n, and $b \in H$ has order m, then $\operatorname{ord}((a,b)) = \operatorname{lcm}(n,m)$ in $G \times H$.
>
> *Proof.* Suppose $(a,b)^k = (e,e)$. Then $a^k = b^k = e$. This gives us that $n \mid k$ and $m \mid k$. Hence,
>
> $$\operatorname{lcm}(n,m) \mid k. \tag{1.3}$$
>
> Furthermore,
>
> $$(a,b)^{\operatorname{lcm}(n,m)} = \left(a^{\operatorname{lcm}(n,m)}, b^{\operatorname{lcm}(n,m)}\right) = (e,e)\,. \tag{1.4}$$
>
> Hence, $\operatorname{ord}((a,b)) = \operatorname{lcm}(n,m)$. □

Now, (a,b) will have order 60 if $\operatorname{lcm}(\operatorname{ord}(a), \operatorname{ord}(b)) = 60$. Since $\operatorname{ord}(a) \in \{1,2,3,6\}$ and $\operatorname{ord}(b) \in \{1,2,4,5,10,20\}$, $\operatorname{lcm}(\operatorname{ord}(a), \operatorname{ord}(b)) = 60$ can only happen in the following cases:

- $\operatorname{ord}(a) = 3, \operatorname{ord}(b) = 20$. Then the choices for b are all the numbers coprime to 20, so we have $\varphi(20) = 8$ choices. The choices for a are $2, 4$. Hence there are $8 \times 2 = 16$ possibilities.
- $\operatorname{ord}(a) = 3, \operatorname{ord}(b) = 20$. Then the choices for b are all the numbers coprime to 20, so we have $\varphi(20) = 8$ choices. The choices for a are $1, 5$. Hence there are $8 \times 2 = 16$ possibilities.

Hence, there are $16 + 16 = 32$ elements of order 60.

(b) Let's first calculate how many elements (a,b) are there of order 30. As before, the possibilities are:

- $\operatorname{ord}(a) = 3, \operatorname{ord}(b) = 10$. Then the choices for b are multiples of 2 by all the numbers coprime to 10, so we have $\varphi(10) = 4$ choices. The choices for a are $2, 4$. Hence there are $4 \times 2 = 8$ possibilities.

- $\bullet \operatorname{ord}(a) = 6, \operatorname{ord}(b) = 10$. Then the choices for b are multiples of 2 by all the numbers coprime to 10, so we have $\varphi(10) = 4$ choices. There are $\varphi(6) = 2$ choices for a. Hence there are $4 \times 2 = 16$ possibilities.

- $\bullet \operatorname{ord}(a) = 6, \operatorname{ord}(b) = 5$. Then the choices for b are multiples of 4 by all the numbers coprime to 5, so we have $\varphi(5) = 4$ choices. There are $\varphi(6) = 2$ choices for a. Hence there are $4 \times 2 = 16$ possibilities.

Hence, there are $8 + 8 + 8 = 24$ numbers of order 30. A subgroup of order 30 consists of $\varphi(30) = 8$ generators. Hence, the number of subgroups of order 30 is $24/8 = 3$.

(c) Note that f is completely fixed by $f\left(\overline{1}\right)$. For f to be injective, $f\left(\overline{1}\right)$ has to be an element of order 12. Let's now calculate how many elements (a, b) are there of order 12. The possibilities are:

- $\bullet \operatorname{ord}(a) = 3, \operatorname{ord}(b) = 4$. Then the choices for b are multiples of 5 by all the numbers coprime to 4, so we have $\varphi(4) = 2$ choices. The choices for a are $2, 4$. Hence there are $2 \times 2 = 4$ possibilities.

- $\bullet \operatorname{ord}(a) = 6, \operatorname{ord}(b) = 4$. Then the choices for b are multiples of 5 by all the numbers coprime to 4, so we have $\varphi(4) = 2$ choices. There are $\varphi(6) = 2$ choices for a. Hence there are $2 \times 2 = 4$ possibilities.

Hence, there are $4 + 4 = 8$ elements of order 12. As a result, there are 12 injective homomorphisms $f : \mathbb{Z}_{12} \to G$.

■

Exercise 1.2. Let $(G_i)_{i=1}^n$ be a family of normal subgroups of G such that

$$G = \left\langle \bigcup_{i=1}^{n} G_i \right\rangle. \tag{1.5}$$

Show that G is isomorphic to the direct product of the G_i if and only if one of the following conditions holds:

1. For every i,
$$G_i \cap \left\langle \bigcup_{j \neq i} G_j \right\rangle = \{e\}. \tag{1.6}$$

2. Every element $g \in G$ can be written uniquely as a product
$$g = g_1 g_2 \cdots g_n, \qquad g_i \in G_i. \tag{1.7}$$

Solution. We will show the equivalence of the following 3 statements:

1. The map $f : G_1 \times \cdots \times G_n \to G$ defined by

$$f(g_1, g_2, \ldots, g_n) = g_1 g_2 \cdots g_n \tag{1.8}$$

is an isomorphism.

2. Every element $g \in G$ can be written uniquely as a product

$$g = g_1 g_2 \cdots g_n, \qquad g_i \in G_i. \tag{1.9}$$

3. For every i,

$$G_i \cap \left\langle \bigcup_{j \neq i} G_j \right\rangle = \{e\}. \tag{1.10}$$

Claim 1.2: Assuming 2, if $g_i \in G_i$ and $g_j \in G_j$ for $i \neq j$, then $g_i g_j = g_j g_i$.

Proof. WLOG, assume $i < j$. Since G_i is normal, $g_j g_i g_j^{-1} \in G_i$, so let's call this element g_i'. Therefore,

$$g_j g_i = g_i' g_j. \tag{1.11}$$

Similarly, since G_j is normal, $g_i^{-1} g_j g_i \in G_j$, so let's call this element g_j'. Therefore,

$$g_j g_i = g_i g_j'. \tag{1.12}$$

But this gives two different representations of the element $g_j g_i$:

$$g_j g_i = e \cdots e \underbrace{g_i'}_{i\text{-th slot}} e \cdots e \underbrace{g_j}_{j\text{-th slot}} e \cdots e = e \cdots e \underbrace{g_i}_{i\text{-th slot}} e \cdots e \underbrace{g_j'}_{j\text{-th slot}} e \cdots e. \tag{1.13}$$

By the uniqueness, we have $g_i = g_i'$ and $g_j = g_j'$, proving that $g_j g_i = g_i g_j$. □

Claim 1.3: Assuming 3, if $g_i \in G_i$ and $g_j \in G_j$ for $i \neq j$, then $g_i g_j = g_j g_i$.

Proof. Consider the element $x := g_i g_j g_i^{-1} g_j^{-1}$. By the normality of G_j, $g_i g_j g_i^{-1} \in G_j$, so that $x \in G_j$. Similarly, by the normality of G_i, $g_j g_i^{-1} g_j^{-1} \in G_i$, so that $x \in G_i$. Since the intersection $G_i \cap G_j = \{e\}$, we can conclude that $x = e$, i.e. $g_i g_j = g_j g_i$. □

(1⇒2) Trivial.

(2⇒3) Let $x \in G_i \cap \left\langle \bigcup_{j \neq i} G_j \right\rangle$. Since $x \in G_i$, it can be uniquely represented as

$$x = e \cdots e \underbrace{x}_{i\text{-th slot}} e \cdots e. \tag{1.14}$$

Since $x \in \left\langle \bigcup_{j\neq i} G_j \right\rangle$, x can be written as a string of elements of G_j. Since elements from different G_j's commute, we can group the elements from the same G_j's and then form

$$x = g_1 g_2 \cdots g_{i-1} e g_{i+1} \cdots g_n. \tag{1.15}$$

by uniqueness, we must have that all g_j's must be e, and also $x = e$. Hence, $G_i \cap \left\langle \bigcup_{j\neq i} G_j \right\rangle = \{e\}$.

(3⇒1) First, we'll show that f is a homomorphism.

$$f(g_1, \ldots, g_n) f\left(g_1', \ldots, g_n'\right) = (g_1 \cdots g_n)\left(g_1' \cdots g_n'\right). \tag{1.16}$$

Since elements from different G_i's commute, this can be rewritten as

$$(g_1 \cdots g_n)\left(g_1' \cdots g_n'\right) = \left(g_1 g_1'\right)\left(g_2 g_2'\right) \cdots \left(g_n g n'\right) = f\left(g_1 g_1', \ldots, g_n g_n'\right), \tag{1.17}$$

i.e. f is a homomorphism. f is surjective indeed: since $G = \langle \bigcup_{i=1}^n G_i \rangle$, every element $g \in G$ can be written as a string of elements of G_i's. Since elements from different G_i's commute, we can group the elements from the same G_i's and then form

$$g = g_1 g_2 \cdots g_n, \quad \text{where } g_i \in G_i \text{ for each } i. \tag{1.18}$$

Now, suppose $(g_1, \ldots, g_n) \in \operatorname{Ker} f$. Then

$$e = g_1 g_2 \cdots g_n. \tag{1.19}$$

From this, we can write

$$\begin{aligned} g_i^{-1} &= g_i^{-1}\ g_1 g_2 \cdots g_n = g_1 g_2 \cdots g_{i-1}\left(g_i^{-1} g_i\right) g_{i+1} \cdots g_n \\ &= g_1 g_2 \cdots g_{i-1} g_{i+1} \cdots g_n \in \left\langle \bigcup_{j\neq i} G_j \right\rangle. \end{aligned} \tag{1.20}$$

(Here the reordering is possible because g_i^{-1} commutes with the other g_j's.) Since $G_i \cap \left\langle \bigcup_{j\neq i} G_j \right\rangle = \{e\}$, this forces that $g_i = e$. This is true for each i. Hence, $\operatorname{Ker} f$ is trivial, and hence it's injective. This completes the proof. ■

Exercise 1.3. Let G be a group, and define

$$\Delta = \{(x, x) \mid x \in G\}. \tag{1.21}$$

(a) Show that Δ is a subgroup of $G \times G$.

(b) Prove that Δ is a normal subgroup of $G \times G$ if and only if G is abelian. Moreover, in this case, show that the quotient group $(G \times G)/\Delta$ is isomorphic to G.

Solution. (a) Take $(x,x),(y,y) \in \Delta$. Then

$$(x,x) * (y,y)^{-1} = (x,x) * \left(y^{-1}, y^{-1}\right) = \left(xy^{-1}, xy^{-1}\right) \in \Delta. \tag{1.22}$$

Therefore, Δ is a subgroup of $G \times G$.

(b) All subgroups of an abelian group are normal. So if G is abelian, so is $G \times G$, and hence Δ is a normal subgroup.

Suppose Δ is a normal subgroup. Then given any $(g,h) \in G \times G$,

$$(g,h) * (x,x) * (g,h)^{-1} = (g,h) * (x,x) * \left(g^{-1}, h^{-1}\right) = \left(gxg^{-1}, hxh^{-1}\right). \tag{1.23}$$

This is in Δ, by hypothesis that Δ is normal. Therefore,

$$gxg^{-1} = hxh^{-1} \tag{1.24}$$

Since this holds for every $g, x, h \in G$, we choose $h = e$, so we have

$$gxg^{-1} = x \implies gx = xg, \tag{1.25}$$

proving that G is abelian. Suppose G is abelian, then we define

$$\begin{aligned} f : G \times G &\to G \\ (x,y) &\mapsto xy^{-1}. \end{aligned} \tag{1.26}$$

This is a homomorphism, because

$$f\left((g,h) * (x,y)\right) = f\left(gx, hy\right) = (gx)\left(hy\right)^{-1} = gxy^{-1}h^{-1} \tag{1.27}$$

$$= \left(gh^{-1}\right)\left(xy^{-1}\right) = f\left(g,h\right) f\left(x,y\right). \tag{1.28}$$

Therefore, f is a homomorphism. Now,

$$(x,y) \in \operatorname{Ker} f \iff xy^{-1} = e \iff x = y. \tag{1.29}$$

Therefore, $\operatorname{Ker} f = \Delta$. f is surjective, since any $g \in G$ is $f(g,e) = g$. Hence, by the first isomorphism theorem,

$$\operatorname{im} f \cong \frac{\operatorname{dom} f}{\operatorname{Ker} f} \implies G \cong \frac{G \times G}{\Delta}. \tag{1.30}$$

■

Exercise 1.4. A subgroup M of a group G is called *maximal* if $M \neq G$ and for every subgroup L of G,

$$M \subsetneq L \subseteq G \implies L = G. \tag{1.31}$$

(a) Let K, M be subgroups of a group G such that $K \trianglelefteq G$ and $K \subseteq M$. Prove that M is a maximal subgroup of G if and only if M/K is a maximal subgroup of G/K.

(b) Prove that in a finite group G, every proper subgroup $H \neq G$ is contained in some maximal subgroup of G.

(c) Prove that in a finite abelian group different from the trivial group, the maximal subgroups are precisely those of prime index.

(d) Is the previous point true if we remove the assumption that the group is abelian?

Solution. (a) Let $\pi : G \to G/K$ denote the projection map. Suppose M is a maximal subgroup of G. Assume for the sake of contradiction that M/K is **NOT** maximal. Then there exists another subgroup

$$\frac{M}{K} \subsetneq X \subsetneq \frac{G}{K}. \tag{1.32}$$

Now consider $\pi^{-1}(X)$.

Claim 1.4: If $f : G_1 \to G_2$ is a homomorphism, and $H_2 \subseteq G_2$ is a subgroup, then $f^{-1}(H_2) \subseteq G_1$ is a subgroup.

Proof. Take $x, y \in f^{-1}(H_2)$. Then $f(x), f(y) \in H_2$. Since $H_2 \subseteq G_2$ is a subgroup,

$$f\left(xy^{-1}\right) = f(x) f(y)^{-1} \in H_2, \tag{1.33}$$

proving that $xy^{-1} \in f^{-1}(H_2)$. So $f^{-1}(H_2) \subseteq G_1$ is a subgroup. □

Now, $\pi^{-1}(X)$ is a subgroup of G, that contains $\pi^{-1}(M/K) = M$. Hence, either $\pi^{-1}(X) = M$ or $\pi^{-1}(X) = G$.

$$\pi^{-1}(X) = \{g \in G \mid gK \in X\}. \tag{1.34}$$

If $\pi^{-1}(X) = G$, then $X = G/K$; and if $\pi^{-1}(X) = M$, then $X = M/K$; both of which contradict (1.32).

Conversely, suppose M/K is a maximal subgroup in G/K. Assume for the sake of contradiction that M is **NOT** maximal. Then there exists another subgroup

$$M \subsetneq H \subsetneq G. \tag{1.35}$$

Since image of a subgroup is also a subgroup, we have $\frac{M}{K} \subset \frac{H}{K} \subset \frac{G}{K}$. By the maximality of M/K, either $H/K = M/K$ or $H/K = G/K$.

Claim 1.5: For any subgroup $L \supseteq K$, $\pi^{-1}(L/K) = L$.

Proof. It's clear that $L \subseteq \pi^{-1}(L/K)$. Now take any $g \in \pi^{-1}(L/K)$. Then $gK \in L/K$.

$$gK = lK \text{ for some } l \in L. \tag{1.36}$$

Therefore, $l^{-1}g \in K \subseteq L$. Hence, $g = l\left(l^{-1}g\right) \in L$. As a result, $L \supseteq \pi^{-1}(L/K)$. □

If $H/K = M/K$, by taking π^{-1}, we get $H = K$; and if $H/K = G/K$, then $H = G$; both of which contradict (1.35).

(b) Let H be a proper subgroup of a finite group G. We construct a sequence of subgroups

$$H =: H_0 \subsetneq H_1 \subsetneq H_2 \subsetneq \cdots \tag{1.37}$$

If H_i is maximal, then we are done, since H is contained in a maximal subgroup! Otherwise, there exists some $g_{i+1} \in G - H_i$ such that

$$H_i \subsetneq H_{i+1} := \langle H_i, g_{i+1} \rangle \subsetneq G. \tag{1.38}$$

If there is no such g_{i+1}, then H_i is indeed maximal, because adding one element to H_i and making a group with it makes it the whole G.

Now, since G is a finite group, this sequence can't go on forever. Therefore, it has to stop somewhere, and thus we find a maximal subgroup that contains H.

(c) Let M be a subgroup of prime index, i.e. $[G : M] = p$. Let's pick a subgroup L with

$$M \subseteq L \subseteq G. \tag{1.39}$$

Then

$$[G : M] = [G : L]\,[L : M]\,. \tag{1.40}$$

(This follows from the fact that for finite groups $[G : H] = \frac{|G|}{|H|}$.) Since $[G : M]$ is a prime, it forces either $[G : L] = 1$, or $[L : M] = 1$; consequently $G = L$ or $L = M$. Therefore, M is maximal.

Now suppose M is a maximal subgroup in a finite abelian group G. Since G is abelian, M is normal, so we can form the quotient group G/M. By part (a), since M is maximal, we can conclude that $\{\overline{e}\} = M/M$ is a maximal subgroup of G/M.

Claim 1.6: If $\{e\}$ is a maximal subgroup of $G \neq \{e\}$, then $G \cong \mathbb{Z}_p$ for some prime p.

Proof. Adding one single non-identity element makes it the whole group, so for every $a \in G$ with $a \neq e$,

$$\langle a \rangle = G. \tag{1.41}$$

Hence, G is cyclic. If $|G|$ was infinite, then it would have subgroups other than the trivial one. But as $\{e\}$ is a maximal subgroup, it's the only subgroup. Therefore, $|G|$ can't be infinite.

Now, if $|G|$ was composite, say $|G| = mn$, then $a^{mn} = e$, proving a^m has order n. This gives rise to a subgroup of size n, namely $\langle a^m \rangle$, which is a contradiction! Therefore, $|G|$ is prime. □

Now, $\{\overline{e}\} = M/M$ is a maximal subgroup of G/M, proves that $|G/M| = p$. Therefore,

$$M \subsetneq G \text{ is maximal } \iff [G : M] \text{ is prime.} \tag{1.42}$$

(d) No. We can consider the group $G = A_4$ of alternating group of degree 4, which contains all even permutations of $\{1, 2, 3, 4\}$. The subgroup

$$H = \langle (1\ 2\ 3) \rangle \tag{1.43}$$

has size 3, so that $[G : H] = 2^2$ is not a prime. However, H is maximal. Indeed, if there exists another subgroup

$$H \subsetneq K \subsetneq G, \tag{1.44}$$

then $|K|$ divides $|G| = 12$, and $|H| = 3$ divides $|K|$, forces $|K| = 6$. If K contains a product of two 2-cycles, e.g. $(1\ 2)\,(3\ 4)$, then

$$(1\ 2\ 3)\,(1\ 2)\,(3\ 4) = (1\ 3\ 4) \in K \tag{1.45}$$

Since K is a group, it contains the inverse $(1\ 4\ 3)$ as well. These are all the 6 elements of K. But then

$$(1\ 2)\,(3\ 4)\,(1\ 2\ 3) = (2\ 4\ 3) \tag{1.46}$$

that gives one more element in K, which is a contradiction!

Therefore, K doesn't contain any product of two 2-cycles, i.e. the only non-identity elements in K are 3-cycles. But every 3-cycle comes with its inverse as well, and no 3-cycle is its own inverse. Therefore, $|K|$ can't be even! Thus we arrive at a contradiction. Hence, H is maximal.

■

Exercise 1.5. Let p be a prime number.

(a) Determine all elements of order dividing p in S_n. Denote by a_n the number of such elements and derive a formula for a_n.

(b) Prove the identity

$$\sum_{n=0}^{\infty} \frac{a_n}{n!} x^n = e^{x + \frac{x^p}{p}}. \tag{1.47}$$

Solution. (a) Every element $\sigma \in S_n$ can be decomposed into disjoint cycles. The disjoint cycles commute. Now, if $\operatorname{ord} \sigma \mid p$, then we must have $\sigma^p = e$. This will happen if and only if $c_i^p = e$ for each disjoint cycle c_i in σ.

Now, for a cycle c_i, $c_i^p = e$ happens if the length of the cycle is either 1 or p. Therefore, a_n will be the number of ways we can decompose $\{1, 2, \ldots, n\}$ into cycles of length 1 or p.

Now, suppose $\sigma \in S_n$ consists of m disjoint cycles of length p, and the other $n-mp$ elements are fixed points (i.e. cycles of length 1). Here, m ranges from 0 to $\left\lfloor \frac{n}{p} \right\rfloor$. Therefore,

$$a_n = \sum_{m=0}^{\left\lfloor \frac{n}{p} \right\rfloor} \text{number of } \sigma\text{'s with } m \text{ disjoint cycles of length } p. \tag{1.48}$$

First, we can choose the numbers involved in the p-cycles in $\binom{n}{mp}$ ways. Now, we can partition them into m sets of size p, which can be done in

$$\binom{mp}{p} \cdot \binom{mp-p}{p} \cdot \cdots \cdot \binom{p}{p} = \frac{(mp)!}{(p!)^m} \tag{1.49}$$

many ways. Now, each cycles can be permuted in $(p-1)!$ different ways. Finally, the cycles themselves can permute in $m!$ ways among themselves. So we need to divide by that. Therefore, the number of σ's with m disjoint cycles of length p is

$$\binom{n}{mp} \cdot \frac{(mp)!}{(p!)^m} \cdot ((p-1)!)^m \cdot \frac{1}{m!} = \frac{n!}{p^m m!\,(n-mp)!}. \tag{1.50}$$

Hence,

$$a_n = \sum_{m=0}^{\left\lfloor \frac{n}{p} \right\rfloor} \frac{n!}{p^m m!\,(n-mp)!}. \tag{1.51}$$

(b) Using (1.51), we have

$$f(x) := \sum_{n=0}^{\infty} \frac{a_n}{n!} x^n = \sum_{n=0}^{\infty} \sum_{m=0}^{\left\lfloor \frac{n}{p} \right\rfloor} \frac{x^n}{p^m m!\,(n-mp)!}. \tag{1.52}$$

We can swap the sum into

$$f(x) = \sum_{m=0}^{\infty} \sum_{n=mp}^{\infty} \frac{x^n}{p^m m!\,(n-mp)!}. \tag{1.53}$$

Suppose $k = n - mp$, so that $n = k + mp$. Then the sum can be rewritten as

$$f(x) = \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{x^{k+mp}}{p^m m! k!} = \sum_{m=0}^{\infty} \frac{1}{m!} \left(\frac{x^p}{p} \right)^m \sum_{k=0}^{\infty} \frac{x^k}{k!} = e^{\frac{x^p}{p}} e^x = e^{\frac{x^p}{p} + x}. \tag{1.54}$$

■

§2 Problem Set 2

Exercise 2.1. Find all p-Sylow subgroups of the following groups:

(a) $\mathbb{Z}/n\mathbb{Z}$;

(b) $\mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$;

(c) the dihedral group D_n (the group of symmetries of a regular n-polygon).

Solution. Before proceeding, we claim that for abelian groups, the p-Sylow subgroup is unique! Indeed, if H_1 and H_2 are two p-Sylow subgroups of G, then by Sylow's Second Theorem, they are related by a conjugation, i.e.

$$H_2 = gH_1g^{-1} \text{ for some } g \in G. \tag{2.1}$$

But if G is abelian, we have

$$H_2 = gH_1g^{-1} = H_1. \tag{2.2}$$

Furthermore, notice that if G is not necessarily abelian, but a p-Sylow subgroup happens to be normal, then also the p-Sylow subgroup is unique. Because if H_1 is a p-Sylow subgroup which happens to be normal, and H_2 is another p-Sylow subgroup, then

$$H_2 = gH_1g^{-1} = H_1. \tag{2.3}$$

We shall use these two properties below.

(a) Let $n = p^k m$, where $p \nmid m$. Since the group is abelian, there is exactly one p-Sylow subgroup, i.e. subgroup of order p^m. That's precisely the subgroup generated by m, since it has order p^k, i.e.

$$H = \langle m \rangle = \left\{0, m, 2m, 3m, \ldots, \left(p^k - 1\right) m\right\}. \tag{2.4}$$

(b) $|G| = 6 \cdot 12 = 2^3 \cdot 3^2$. Since the group is abelian, there is exactly one p-Sylow subgroup. For $p = 2$, we can choose a copy of $\mathbb{Z}_2$ inside $\mathbb{Z}_6$, and a copy of $\mathbb{Z}_4$ inside $\mathbb{Z}_{12}$, and take their direct product, and that will be the subgroup of order 2^3.

$$H = \{0, 3\} \times \{0, 3, 6, 9\} \subset \mathbb{Z}_6 \times \mathbb{Z}_{12}. \tag{2.5}$$

For $p = 3$, we can choose a copy of $\mathbb{Z}_3$ inside $\mathbb{Z}_6$, and a copy of $\mathbb{Z}_3$ inside $\mathbb{Z}_{12}$, and take their direct product, and that will be the subgroup of order 3^2.

$$H' = \{0, 2, 4\} \times \{0, 4, 8\} \subset \mathbb{Z}_6 \times \mathbb{Z}_{12}. \tag{2.6}$$

(c) $|D_n| = 2n$. Suppose $p \neq 2$. Then $p \mid 2n$ if and only if $p \mid n$. We write

$$2n = 2 \cdot p^k \cdot m, \tag{2.7}$$

where $p \nmid m$. Let a be rotation by $\frac{2\pi}{n}$. Then $\langle a \rangle$ is an order-n subgroup of D_n. We take

$$H = \langle a^m \rangle = \left\{ e, a^m, a^{2m}, \ldots, a^{\left(p^k-1\right)m} \right\}, \tag{2.8}$$

which is a subgroup of order p^k, i.e. a p-Sylow subgroup. Now, we claim that H is normal in D_n. Indeed, all the rotations commute with all the elements of H. If b is a reflection, then we have

$$ba^i b^{-1} = a^{-i}, \tag{2.9}$$

(this can be proven by induction on i), and hence

$$\left(a^j b\right)\ a^i\ \left(a^j b\right)^{-1} = a^j b a^i b a^{-j} = a^j a^{-i} a^{-j} = a^{-i}. \tag{2.10}$$

Therefore, H is normal, and hence there is only one p-Sylow subgroup for $p \neq 2$.

Now suppose $p = 2$. If $2 \nmid n$, then

$$n_2 \mid n \implies n_2 \leq n. \tag{2.11}$$

We can then exhibit n subgroups of order 2, namely the ones containing one reflection and the identity:

$$H_i = \left\{ e, a^i b \right\}, \quad \text{for } i = 0, 1, 2, \ldots, n-1. \tag{2.12}$$

Now consider the case $2 \mid n$. Suppose $n = 2^k m$, where m is odd. Then

$$|D_n| = 2n = 2^{k+1} m. \tag{2.13}$$

Then

$$n_2 \mid m \implies n_2 \leq m. \tag{2.14}$$

We now exhibit m subgroups of order 2^{k+1}. These will be isomorphic to D_{2^k}. We define

$$H_i = \left\langle a^m, a^i b \right\rangle, \quad \text{for } i = 0, 1, 2, \ldots, m-1 \tag{2.15}$$

Here, $a' := a^m$ is a rotation of order 2^k (since $a^{2^k m} = a^n = e$), and $b' := a^i b$ is a reflection of degree 2, and

$$b'a'b' = \left(a^i b\right)\left(a^m\right)\left(a^i b\right) = a^i \left(b a^{i+m} b\right) = a^i a^{-i-m} = a^{-m} = \left(a'\right)^{-1}, \tag{2.16}$$

and therefore,

$$H_i = \left\langle a', b' \mid \left(a'\right)^{2^k} = \left(b'\right)^2 = e,\ b'a'b' = \left(a'\right)^{-1} \right\rangle \cong D_{2^k}, \tag{2.17}$$

so each H_i is a subgroup of size 2^{k+1}, i.e. a 2-Sylow subgroup. Since $n_2 \leq m$, and we have exhibited m subgroups, these are the all of them!

■

Exercise 2.2. Find all 2-Sylow subgroups of the following groups:

(a) the symmetric group S_4;

(b) the symmetric group S_5.

Solution. (a) $|S_4| = 24 = 2^3 \cdot 3$. So we need to find all subgroups of size 8. By Sylow's theorems, if n_2 is the number of 2-Sylow subgroups,

$$n_2 \mid 3 \implies n_2 \in \{1, 3\}\,. \tag{2.18}$$

We claim that there are 3 of them, by finding 3 subgroups of order 8. First, note that there's a subgroup isomorphic to D_4.




Figure 1: Rotation in D_4, which can be realized as the element (1 2 3 4). To see the resemblence, in the left figure, 1 is in the top-right position, which is occupied by 2 in the right figure, so $\sigma(1) = 2$, etc.




Figure 2: Reflection in D_4, which can be realized as the element (2 4). To see the resemblence, in the left figure, 2 is in the top-left position, which is occupied by 4 in the right figure, so $\sigma(2) = 4$, etc.

These two generate D_4, so we can take the corresponding subgroup in S_4

$$G_1 = \langle (1\ 2\ 3\ 4)\,, (2\ 4) \rangle\,, \tag{2.19}$$

which is isomorphic to D_4, and hence a subgroup of order 8, i.e. a 2-Sylow subgroup. To find the others, it's sufficient to change the labels in the vertices of the square. There are

3-possible ways we can cyclically rename the vertices, which gives the other two subgroups:

$$G_2 = \langle (1\ 3\ 2\ 4), (1\ 2) \rangle, \tag{2.20}$$

$$G_3 = \langle (1\ 2\ 4\ 3), (1\ 4) \rangle. \tag{2.21}$$

These are all the 2-Sylow subgroups of size 8.

(b) $|S_5| = 120 = 2^3 \cdot 15$. So we need to find all subgroups of size 8. By Sylow's theorems, if n_2 is the number of 2-Sylow subgroups,

$$n_2 \mid 15 \implies n_2 \in \{1, 3, 5, 15\}. \tag{2.22}$$

Note that, given any 4-element subset of $\{1, 2, 3, 4, 5\}$, we can find 3 subgroups of size 8, as per part (a). The total number of 4-element subsets of $\{1, 2, 3, 4, 5\}$ is $\binom{5}{4} = 5$. Hence, this already gives us a recipe to build $5 \times 3 = 15$ subgroups of order 8. Since $n_2 \leq 15$, these are all of them!

■

Exercise 2.3. (a) Show that there are no simple groups of order 30, 42 or 105.

(b) More generally, show that there are no simple groups of order pqr, where p, q, r are distinct prime numbers.

Solution. We'll just solve (b). Assume for the sake of contradiction that ther exists a simple group G with $|G| = pqr$. WLOG, $p < q < r$.

Note that, we must have $n_r > 1$, because if $n_r = 1$, then the r-Sylow subgroup would've been normal, contradicting the simplicity of G. Also, $n_r \equiv 1 \pmod{r}$ forces that $n_r > r > q > p$. Now, since

$$n_r \mid pq, \tag{2.23}$$

the only possibility is $n_r = pq$. In other words, there are pq subgroups of order r, which gives

$$(\text{number of elements of order } r) = n_r (r-1) = pq (r-1) = |G| - pq. \tag{2.24}$$

Similarly, we must have $n_q > 1$, because if $n_q = 1$, then the q-Sylow subgroup would've been normal, contradicting the simplicity of G. Also, $n_q \equiv 1 \pmod{q}$ forces that $n_q > q > p$. Therefore,

$$(\text{number of elements of order } q) = n_q (q-1) > pq - p. \tag{2.25}$$

Therefore,

$$(\text{number of elements of order } p) = n_p (p-1) < p. \tag{2.26}$$

But this forces $n_p = 1$, which implies the p-Sylow subgroup is normal, contradicting the simplicity of G. Therefore, there is no simple group of order pqr. ■

Exercise 2.4. Consider the action of a finite group G on a finite set X, with N orbits. For every $g \in G$, consider the set

$$\mathrm{Fix}(g) := \{x \in X \mid g \cdot x = x\}. \tag{2.27}$$

(a) Derive Burnside's formula:

$$N = \frac{1}{|G|} \sum_{g \in G} |\mathrm{Fix}(g)| \,. \tag{2.28}$$

(b) Apply the previous formula to solve the following problem: count the numbers of bracelets made of 3 red beads and 7 blue beads. *Hint:* Two bracelets are indistinguishable if one is obtained by the other with a rotation or a reflection.

Solution. (a) Consider the set

$$S = \{(g, x) \mid g \cdot x = x\} \,. \tag{2.29}$$

We can count $|S|$ in two different ways. For every $g \in G$, we count the number of $x \in S$ fixed by g. Then for every $x \in X$, we count the number of $g \in G$ that fixes x. Therefore,

$$|S| = \sum_{g \in G} |\mathrm{Fix}(g)| = \sum_{x \in X} |G_x| \,, \tag{2.30}$$

where G_x is the stabilizer of x. Since X is the disjoint union of G-orbits,

$$\sum_{x \in X} |G_x| = \sum_{\text{orbits } Gx} \sum_{y \in Gx} |G_y| \,. \tag{2.31}$$

Now, for $y \in Gx$, $Gx = Gy$, so their stabilizers have the same size, i.e. $|G_x| = |G_y|$.

> **Claim 2.1:** For $y \in Gx$, $|G_x| = |G_y|$
>
> *Proof.* Since $y \in Gx$, $y = g \cdot x$ for some $g \in G$. If $g_0 \in G_x$ is in the stabilizer of x, then $g_0 \cdot x = x$. Now,
>
> $$\left(gg_0g^{-1}\right) \cdot y = \left(gg_0g^{-1}\right) \cdot (g \cdot x) = \left(gg_0g^{-1}g\right) \cdot x = g \cdot (g_0 \cdot x) = g \cdot x = y. \tag{2.32}$$
>
> Therefore, $gg_0g^{-1} \in G_y$, i.e. $G_y = gG_xg^{-1}$. Since conjugation preserves size, $|G_x| = |G_y|$. □

With this, we rewrite (2.31),

$$\sum_{x \in X} |G_x| = \sum_{\text{orbits } Gx} |Gx| \, |G_x| \tag{2.33}$$

By Orbit-Stabilizer theorem, $|G| = |Gx| \, |G_x|$. Hence,

$$\sum_{x \in X} |G_x| = \sum_{\text{orbits } Gx} |Gx| \, |G_x| = N \, |G| \,, \tag{2.34}$$

where N is the number of orbits. Hence,

$$N = \frac{1}{|G|} \sum_{x \in X} |G_x| = \frac{1}{G} \sum_{g \in G} |\mathrm{Fix}(g)| \,. \tag{2.35}$$

(b) Let $G = D_{10}$, the dihedral group on a regular 10-gon. Then $|G| = 20$. Let X be the set of all bracelets made of 3 red beads and 7 blue beads (without accounting for indistinguishability). Then

$$|X| = \binom{10}{3} = \frac{10 \cdot 9 \cdot 8}{3!} = 120. \tag{2.36}$$

Consider D_{10}'s action on X. Then the number of indistinguishable bracelets will be the number of orbits in this action. By Burnside's lemma, if N is the number of orbits,

$$N = \frac{1}{|G|} \sum_{g \in G} |\mathrm{Fix}(g)| \,. \tag{2.37}$$

For each $g \in G$, let's now compute the size of $\mathrm{Fix}(g)$.

- If $g = e$, $|\mathrm{Fix}(e)| = |X| = 120$.
- If g is a rotation by $\frac{2\pi}{10}k$ for $k = 1, \ldots, 9$, then we claim that $\mathrm{Fix}(g) = \varnothing$.

  If we consider the positions (in modulo 10), then a rotation is just adding a constant with the positions. If r_1, r_2, r_3 are the position of the red beads in a configuration, if g fixes that configuration, then we must have

  $$\{r_1, r_2, r_3\} = \{r_1 + a, r_2 + a, r_3 + a\} \pmod{10}. \tag{2.38}$$

  After adding, we have

  $$r_1 + r_2 + r_3 \equiv r_1 + r_2 + r_3 + 3a \pmod{10}. \tag{2.39}$$

  In other words, $10 \mid 3a$. But since 10 and 3 are coprime, this means $a = 0 \pmod 1 0$, i.e. the rotation is the trivial one. Therefore, for nontrivial rotations, $|\mathrm{Fix}(g)| = 0$.
- If g is reflection in a line ℓ through the midpoints of opposite edges, then there are the same number of beads on both sides of ℓ, and we can't arrange the beads of the same color evenly, because there are an odd number of red beads and an odd number of blue beads. So in this case, $|\mathrm{Fix}(g)| = 0$. There are 5 such reflections.
- If g is reflection in a line ℓ through two opposite vertices, then we can choose one of them to be red, and we can choose the other to be blue. This can be done in $\binom{2}{1} = 2$ ways.

  Then if we color one side of ℓ, the other side becomes fixed. On each side of ℓ, we have 4 points, and we can choose one of them to be red, which can be done in $\binom{4}{1} = 4$ ways. Therefore, $|\mathrm{Fix}(g)| = 2 \times 4 = 8$. There are 5 such reflections.

Now,

$$N = \frac{1}{20}\left(120 + 0 \times 9 + 0 \times 5 + 8 \times 5\right) = \frac{160}{20} = 8. \tag{2.40}$$

■

Exercise 2.5 (Frattini's Lemma)**.** Let $H \trianglelefteq G$ be a normal subgroup of a finite group. Let S be a p-Sylow of H. Show that the conjugates of S in G are in H as well and deduce the following:

$$G = HN_G(S), \tag{2.41}$$

where $N_G(S)$ denotes the normalizer of S in G.

Solution. For every $g \in G$, since $S \leq H$,

$$gSg^{-1} \subseteq gHg^{-1} = H, \tag{2.42}$$

so conjugates of S in G are in H as well! Now we shall prove that any $g \in G$ can be written as $g = hn$ for some $h \in H, n \in N_G(S)$.

Note that since $gSg^{-1} \subseteq H$, gSg^{-1} is also a p-Sylow subgroup of H, because conjugation preserves cardinality. Since both S and gSg^{-1} are p-Sylow subgroups of H, by the Second Sylow Theorem, they are related by a conjugation in H, i.e.

$$gSg^{-1} = hSh^{-1} \tag{2.43}$$

for some $h \in H$. This gives

$$h^{-1}gSg^{-1}h = S. \tag{2.44}$$

Since $N_G(S) = \left\{g \mid gSg^{-1} = S\right\}$, by (2.44), $h^{-1}g \in N_G(S)$.

$$h^{-1}g = n \text{ for some } N_G(S), \text{ for some } h \in H \tag{2.45}$$

$$\implies g = hn \text{ for some } N_G(S), \text{ for some } h \in H. \tag{2.46}$$

Therefore, $G = HN_G(S)$. ■

Exercise 2.6 (Short exact sequences)**.** Given a sequence of morphisms

$$N \xrightarrow{a} G \xrightarrow{b} K \tag{2.47}$$

we say that it is exact in G if

$$\operatorname{Ker} b = \operatorname{im} a. \tag{2.48}$$

A sequence of the form

$$e \longrightarrow N \xrightarrow{a} G \xrightarrow{b} K \longrightarrow e \tag{2.49}$$

which is exact in N, G, K is called a short exact sequence (where we denote with e the group with one element).

(a) Show that in the short exact sequence above a is injective and b surjective.

(b) Show that $a(N) \trianglelefteq G$ and

$$K \simeq G/a(N).$$

(c) Let N and K be groups. Show that you can define a short exact sequence:

$$e \longrightarrow N \xrightarrow{\ i\ } N \times K \xrightarrow{\ p\ } K \longrightarrow e \tag{2.50}$$

Solution. (a) Exactness at N gives

$$\operatorname{Ker} a = \operatorname{im} (e \to N) = \{e\}, \tag{2.51}$$

so a is injective. Exactness at K gives

$$\operatorname{im} b = \operatorname{Ker} (K \to e) = K, \tag{2.52}$$

so b is surjective.

(b) Exactness at G gives $a(N) = \operatorname{im} a = \operatorname{Ker} b$. Kernel is always normal, because for $h \in \operatorname{Ker} b$ and $g \in G$,

$$b\left(ghg^{-1}\right) = b(g)\, b(h)\, b(g)^{-1} = b(g)\, b(g)^{-1} = e, \tag{2.53}$$

so $ghg^{-1} \in \operatorname{Ker} b$, proving $\operatorname{Ker} b$ is normal. Now, since b is surjective,

$$\operatorname{im} b \cong \frac{\operatorname{dom} b}{\operatorname{Ker} b} \implies K \cong \frac{G}{\operatorname{Ker} b} = \frac{G}{a(N)}. \tag{2.54}$$

(c) We define

$$i(n) = (n, e_K) \text{ and } p(n, k) = k. \tag{2.55}$$

Then clearly i is injective and p is surjective. Also,

$$\operatorname{im} i = N \times \{e_K\} = \operatorname{Ker} p. \tag{2.56}$$

Hence, the sequence $e \to N \to N \times K \to K \to e$ is exact.

■

Exercise 2.7 (Semidirect products)**.** Let N and K be groups, and let

$$\varphi : K \longrightarrow \operatorname{Aut}(N) \tag{2.57}$$

be a group homomorphism. For each $k \in K$ and $n \in N$, we denote by $\varphi(k)(n)$ the image of n under the automorphism of N corresponding to k via φ.

(a) Define a binary operation on the set-theoretic product $N \times K$ by

$$(n,k) * \left(n',k'\right) = \left(n\ \varphi(k)\left(n'\right), kk'\right). \tag{2.58}$$

Show that this operation defines a group. We denote this group by $N \rtimes_\varphi K$ and call it the semidirect product of N by K with respect to φ.

(b) Show that there is a short exact sequence

$$e \longrightarrow N \xrightarrow{\ i\ } N \rtimes_\varphi K \xrightarrow{\ p\ } K \longrightarrow e \tag{2.59}$$

where

$$i(n) = (n,e) \qquad \text{and} \qquad p(n,k) = k. \tag{2.60}$$

(c) Show that if

$$\varphi : K \longrightarrow \mathrm{Aut}(N) \tag{2.61}$$

is the trivial homomorphism, then $N \rtimes_\varphi K$ is the usual direct product of groups $N \times K$.

(d) Conversely, let G be a group with subgroups $N, K \leq G$ such that

$$N \trianglelefteq G, \qquad N \cap K = \{e\}, \qquad G = NK. \tag{2.62}$$

Show that G is isomorphic to the semidirect product $N \rtimes_\varphi K$, where

$$\varphi(k)(n) = knk^{-1} \tag{2.63}$$

for $n \in N$ and $k \in K$.

Solution. (a) For simplicity of notations, I'll write $\varphi_k = \varphi(k)$. Since φ is a homomorphism, this will give us

$$\varphi_{kk'} = \varphi\left(kk'\right) = \varphi\left(k\right) \circ \varphi\left(k'\right) = \varphi_k \circ \varphi k'. \tag{2.64}$$

Associativity:

$$\begin{aligned}
\left[(n,k) * \left(n',k'\right)\right] * \left(n'',k''\right) &= \left(n\ \varphi_k\left(n'\right), kk'\right) * \left(n'',k''\right) \\
&= \left(n\ \varphi_k\left(n'\right)\ \ \varphi_{kk'}\left(n''\right), (kk')k''\right) \\
&= \left(n\ \varphi_k\left(n'\right)\ \ \left(\varphi_k\varphi_{k'}\left(n''\right)\right), (kk')k''\right) \\
&= \left(n\ \varphi_k\left(n'\varphi_{k'}\left(n''\right)\right), k\left(k'k''\right)\right). \qquad (2.65) \\
(n,k) * \left[\left(n',k'\right) * \left(n'',k''\right)\right] &= (n,k) * \left(n'\varphi_{k'}\left(n''\right), k'k''\right) \\
&= \left(n\ \varphi_k\left(n'\varphi_{k'}\left(n''\right)\right), k\left(k'k''\right)\right). \qquad (2.66)
\end{aligned}$$

We claim that (e_N, e_K) is the identity.

$$(e_N, e_K) * (n, k) = (e_N \varphi_{e_K}(n), e_K k) \tag{2.67}$$
$$(n, k) * (e_N, e_K) = (n\varphi_k(e_N), k e_K) = (n, k). \tag{2.68}$$

We claim that $(n,k)^{-1} = \left(\varphi_{k^{-1}}\left(n^{-1}\right), k^{-1}\right)$.

$$\begin{aligned}(n,k) * \left(\varphi_{k^{-1}}\left(n^{-1}\right), k^{-1}\right) &= \left(n\varphi_k\left(\varphi_{k^{-1}}\left(n^{-1}\right)\right), kk^{-1}\right) \\ &= \left(nn^{-1}, kk^{-1}\right) = (e_N, e_K) \qquad (2.69)\\ \left(\varphi_{k^{-1}}\left(n^{-1}\right), k^{-1}\right) * (n,k) &= \left(\varphi_{k^{-1}}\left(n^{-1}\right)\varphi_{k^{-1}}(n), k^{-1}k\right) \\ &= \left(\varphi_{k^{-1}}\left(n^{-1}n\right), k^{-1}k\right) = (e_N, e_K). \qquad (2.70)\end{aligned}$$

Therefore, $N \rtimes_\varphi K$ is a group.

(b) Let's first verify that i and p are group homomorphisms:

$$\begin{aligned}i(n_1) * i(n_2) = (n_1, e_K) * (n_2, e_K) &= (n_1 \varphi_{e_K}(n_2), e_K e_K) \qquad (2.71)\\ &= (n_1 n_2, e_K) = i(n_1 n_2). \qquad (2.72)\\ p\left((n,k) * \left(n', k'\right)\right) = p\left(n\ \varphi_k\left(n'\right), kk'\right) &= kk' = p(n,k)\, p\left(n', k'\right). \qquad (2.73)\end{aligned}$$

Clearly i is injective and p is surjective. Also,

$$\operatorname{im} i = N \times \{e_K\} = \operatorname{Ker} p. \tag{2.74}$$

Hence, the sequence $e \to N \to N \rtimes_\varphi K \to K \to e$ is exact.

(c) If φ is the trivial homomorphism, then $\varphi_k = \mathrm{id}_N$ for every k. Then

$$(n,k) * \left(n', k'\right) = \left(n\ \varphi_k\left(n'\right), kk'\right) = \left(n\ \mathrm{id}_N\left(n'\right), kk'\right) = \left(nn', kk'\right), \tag{2.75}$$

which is exactly the same as the group operation on the direct product $N \times K$.

(d) We define the map

$$\begin{aligned}f : N \rtimes_\varphi K &\to G \\ (n,k) &\mapsto nk.\end{aligned} \tag{2.76}$$

Let's first check that it's a homomorphism:

$$\begin{aligned}f\left((n,k) * \left(n', k'\right)\right) = f\left(\left(n\ \varphi_k\left(n'\right), kk'\right)\right) &= f\left(n\ kn'k^{-1}, kk'\right) \qquad (2.77)\\ &= nkn'k^{-1}\ kk' = nk\ n'k' \qquad (2.78)\\ &= f(n,k)\, f\left(n', k'\right). \qquad (2.79)\end{aligned}$$

It's injective because if $f(n,k) = e$, then

$$e = f(n,k) = nk \implies n = k^{-1} \in N \cap K = \{e\}, \tag{2.80}$$

so $\operatorname{Ker} f = \{(e,e)\}$. Also, f is surjective since $G = NK$, so every element $g \in G$ can be written as $g = nk$ for some $n \in N$ and $k \in K$. Therefore, f is an isomorphism.

■

§3 Problems on Groups from Lothar Göttsche's Notes

Exercise 3.1. Find a subgroup H of S_3 of order 3, i.e. with $|H| = 3$.

Solution. Consider

$$H = \langle (1\ 2\ 3) \rangle = \{e, (1\ 2\ 3), (1\ 3\ 2)\}. \tag{3.1}$$

Then $|H| = 3$, and H is a subgroup of S_3. In fact, H is isomorphic to $\mathbb{Z}_3$. ■

Exercise 3.2. Let G be a group. The center $C(G)$ of G is

$$C(G) := \{g \in G \mid ga = ag,\ \forall a \in G\}. \tag{3.2}$$

(a) Show $C(G)$ is a subgroup of G.

(b) Show $C(G)$ is commutative.

(c) What is the center of S_3?

Solution. (a) First, $C(G) \neq \varnothing$, since $e \in C(G)$. Now, given any $x, y \in C(G)$, we'll show that $xy \in C(G)$ and $x^{-1} \in C(G)$. First notice that

$$(xy)\, a = x\,(ya) = x\,(ay) = (xa)\, y = (ax)\, y = a\,(xy), \tag{3.3}$$

for every $a \in G$; and hence, $xy \in C(G)$. Now,

$$xa = ax \implies x^{-1}(xa)\, x^{-1} = x^{-1}(ax)\, x^{-1} \implies ax^{-1} = x^{-1}a, \tag{3.4}$$

for every $a \in G$; and hence, $x^{-1} \in C(G)$. Therefore, $C(G) \leq G$.

(b) The elements of $C(G)$ commute with **every** element of G. So, in particular, they commute among themselves. Therefore, $C(G)$ is abelian.

(c) We claim that $C(S_3) = \{e\}$.

$$(1\ 2)(1\ 2\ 3) = (2\ 3), \quad (1\ 2\ 3)(1\ 2) = (1\ 3). \tag{3.5}$$

Hence, neither (1 2), nor (1 2 3) belong in the center. Since center is a subgroup, $(1\ 2\ 3)^{-1} = (1\ 3\ 2)$ is also not in the center. We can rule out (2 3) and (1 3) similarly:

$$(2\ 3)(1\ 2\ 3) = (1\ 3), \quad (1\ 2\ 3)(2\ 3) = (1\ 2); \tag{3.6}$$

$$(1\ 3)(1\ 2\ 3) = (1\ 2), \quad (1\ 2\ 3)(1\ 3) = (2\ 3). \tag{3.7}$$

Therefore, $C(S_3) = \{e\}$.

■

Exercise 3.3. Prove that every subgroup of a cyclic group is cyclic.

Solution. Let $G = \langle a \rangle$ be a cyclic group and $H \leq G$. Then every element of H is of the form a^i for some $i \in \mathbb{Z}$. Let

$$m = \min \left\{ i \in \mathbb{Z}_{>0} \mid a^i \in H \right\}. \tag{3.8}$$

We claim that $H = \langle a^m \rangle$. Since $a^m \in H$, it's clear that $\langle a^m \rangle \subset H$. Assume for the sake of contradiction that there exists some $b \in H$ and $b \notin \langle a^m \rangle$. Since $b \in G = \langle a \rangle$,

$$b = a^n, \quad \text{for some } n \in \mathbb{Z}. \tag{3.9}$$

Now, by performing division algorithm on n, we can write

$$n = mq + r, \quad \text{for } 0 \leq r < m. \tag{3.10}$$

Since $b \notin \langle a^m \rangle$, $r > 0$. Now, since $b = a^{mq+r}, a^m \in H$, $a^r \in H$. But m is the minimal positive integer such that $a^m \in H$, and now we have found an even smaller positive integer. Contradiction! ■

Exercise 3.4. Let G be a finite group. Let $K \subset H$ and $H \subset G$ be subgroups. Show that

$$[G : K] = [G : H][H : K]. \tag{3.11}$$

Solution. We shall use the fact that for finite groups, $[G : H] = \frac{|G|}{|H|}$.

$$[G : H][H : K] = \frac{|G|}{|H|} \cdot \frac{|H|}{|K|} = \frac{|G|}{|K|} = [G : K]. \tag{3.12}$$

■

Exercise 3.5. Let G be a group and H, K normal subgroups. Then $H \cap K$ is a normal subgroup of G.

Solution. It's easy to see that $N := H \cap K$ is a subgroup of G. Indeed, $N \neq \varnothing$ as $e \in N$. For $x, y \in N$,

$$x, y \in H \text{ implies } xy^{-1} \in H; \text{ and } x, y \in K \text{ implies } xy^{-1} \in K. \tag{3.13}$$

Therefore, $xy^{-1} \in H \cap K = N$. Hence, $N \leq G$. Now, for any $g \in G$,

$$N \subseteq H \implies gNg^{-1} \subseteq gHg^{-1} = H, \quad \text{and } N \subseteq K \implies gNg^{-1} \subseteq gKg^{-1} = K. \tag{3.14}$$

Hence, $gNg^{-1} \subseteq H \cap K = N$, proving that $N \trianglelefteq G$. ■

Exercise 3.6. Let G be a group and H a subgroup of index 2. Show H is a normal subgroup of G.

Solution. We shall prove a more general statement:

Suppose G is a finite group, and let p be the smallest prime divisor of $|G|$. If H is a subgroup with $[G : H] = p$, then $H \trianglelefteq G$.

Consider the action of G on the left coset space of H, by left multiplication:

$$g \cdot (aH) := (ga) H. \tag{3.15}$$

This gives a group homomorphism

$$f : G \to \mathrm{Sym}\,(G/H) \cong S_p. \tag{3.16}$$

Then $\mathrm{Ker}\, f \trianglelefteq G$. Furthermore, if $k \in \mathrm{Ker}\, f$, then for all $aH \in G/H$,

$$k \cdot (aH) := (ka) H = aH. \tag{3.17}$$

In particular, choosing $a = e$ gives us $kH = H$, i.e. $k \in H$. Hence,

$$\mathrm{Ker}\, f \leq H \leq G. \tag{3.18}$$

By the first isomorphism theorem, $\mathrm{im}\, f \cong \frac{G}{\mathrm{Ker}\, f}$, and hence $|\mathrm{im}\, f|$ divides $|G|$. But since $\mathrm{im}\, f$ is (isomorphic to) a subgroup of S_p, it divides $|S_p| = p!$ as well. Therefore,

$$|\mathrm{im}\, f| \,\Big|\, \gcd\left(|G|, p!\right) = p, \tag{3.19}$$

since p is the smallest prime factor of $|G|$. Therefore, $|\mathrm{im}\, f| = |G/\mathrm{Ker}\, f| = [G : \mathrm{Ker}\, f]$ divides p. From (3.18), we have

$$[G : \mathrm{Ker}\, f] = [G : H]\,[H : \mathrm{Ker}\, f] = p\,[H : \mathrm{Ker}\, f]\,, \tag{3.20}$$

which divides p. Hence, we must have $[H : \mathrm{Ker}\, f] = 1$, i.e. $H = \mathrm{Ker}\, f$. Since kernels are always normal, $H \trianglelefteq G$. ■

Exercise 3.7. Let G be a group. Let G' be the subgroup of G generated by

$$U = \{xyx^{-1}y^{-1} \mid x, y \in G\}. \tag{3.21}$$

(a) Show G' is a normal subgroup of G.

(b) Show G/G' is abelian.

Solution. (a) We denote $[x, y] := xyx^{-1}y^{-1}$. If $f : G \to G$ is a homomorphism, then

$$f\left([x, y]\right) = f\left(xyx^{-1}y^{-1}\right) = f(x)\, f(y)\, f(x)^{-1}\, f(y)^{-1} = [f(x), f(y)]\,. \tag{3.22}$$

Therefore, $f(U) \subseteq U$. As a result,

$$f\left(G'\right) = f\left(\langle U\rangle\right) \subseteq \langle U\rangle = G'. \tag{3.23}$$

For any $g \in G$, $x \mapsto gxg^{-1}$ is a homomorphism, so $gG'g^{-1} \subseteq G'$ for any $g \in G$. Hence, $G' \trianglelefteq G$.

(b)

$$\begin{aligned} \left(xG'\right)\left(yG'\right) = \left(yG'\right)\left(xG'\right) &\iff (xy)\,G' = (yx)\,G' \\ &\iff (yx)^{-1}\,(xy) \in G' \\ &\iff x^{-1}y^{-1}xy \in G', \end{aligned} \tag{3.24}$$

which is true! Hence, G/G' is abelian.

■

Exercise 3.8. Let G be a group. Show that $\mathrm{Inn}(G)$ is a normal subgroup of $\mathrm{Aut}(G)$.

Solution. First, we'll show that $\mathrm{Inn}(G) \leq \mathrm{Aut}(G)$. Let $\tau_g, \tau_h \in \mathrm{Inn}(G)$, where

$$\tau_g\,(x) = gxg^{-1}, \quad \tau_h\,(x) = hxh^{-1}. \tag{3.25}$$

Then

$$(\tau_g \circ \tau_h)\,(x) = \tau_g\left(hxh^{-1}\right) = g\left(hxh^{-1}\right)g^{-1} = (gh)\,x\,(gh)^{-1} = \tau_{gh}(x). \tag{3.26}$$

Hence, $\tau_g \circ \tau_h = \tau_{gh} \in \mathrm{Inn}(G)$. Furthermore, we claim that $\tau_g^{-1} = \tau_{g^{-1}}$.

$$\tau_g \circ \tau_{g^{-1}} = \tau_{gg^{-1}} = \tau_e = \mathrm{id}_G, \quad \tau_{g^{-1}} \circ \tau_g = \tau_{g^{-1}g} = \tau_e = \mathrm{id}_G\,. \tag{3.27}$$

Hence, $\tau_g^{-1} = \tau_{g^{-1}} \in \mathrm{Inn}(G)$.

Now we show $\mathrm{Inn}(G) \trianglelefteq \mathrm{Aut}(G)$. Let $\varphi \in \mathrm{Aut}(G)$, and $\tau_g \in \mathrm{Inn}(G)$.

$$\left(\varphi \circ \tau_g \varphi^{-1}\right)(x) = \varphi\left(g \cdot \varphi^{-1}(x) \cdot g^{-1}\right) = \varphi(g) \cdot x \cdot \varphi(g)^{-1} = \tau_{\varphi(g)}(x). \tag{3.28}$$

Hence, $\varphi \circ \tau_g \varphi^{-1} = \tau_{\varphi(g)} \in \mathrm{Inn}(G)$, proving that $\mathrm{Inn}(G) \trianglelefteq \mathrm{Aut}(G)$. ■

Exercise 3.9. Give an example of a group G and subgroups $K \subset H \subset G$, such that K is a normal subgroup of H and H is a normal subgroup of G, but K is not a normal subgroup of G.

Solution. Consider

$$G = D_4 = \left\{e, r, r^2, r^3, s, sr, sr^2, sr^3\right\}, \tag{3.29}$$

$$H = \left\{e, r^2, s, sr^2\right\}, \tag{3.30}$$

$$K = \{e, s\}\,. \tag{3.31}$$

Then

$$[G : H] = [H : K] = 2, \tag{3.32}$$

so $K \trianglelefteq H \trianglelefteq G$. But $K \not\trianglelefteq G$, since

$$rsr^{-1} = r\left(sr^{-1}\right) = r\left(rs\right) = r^2 s \notin K. \tag{3.33}$$

Therefore, K is not a normal subgroup of G. ■

Exercise 3.10. Let H, K be normal subgroups of a group G with $H \cap K = \{1\}$. Show that

$$hk = kh \quad \text{for all } h \in H,\ k \in K. \tag{3.34}$$

Solution. Consider the element $hkh^{-1}k^{-1}$.

$$hkh^{-1}k^{-1} = \left(hkh^{-1}\right)k^{-1} \in K, \tag{3.35}$$

by the normality of K. Again,

$$hkh^{-1}k^{-1} = h\left(kh^{-1}k^{-1}\right) \in H, \tag{3.36}$$

by the normality of H. Therefore, $hkh^{-1}k^{-1} \in H \cap K = \{e\}$, so

$$hkh^{-1}k^{-1} = e \implies hk = kh. \tag{3.37}$$

■

Exercise 3.11. Let

$$\sigma := \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 2 & 3 & 4 & 5 & 1 & 6 & 7 & 9 & 8 \end{pmatrix} \in S_9. \tag{3.38}$$

Find the cycle decomposition of σ, find the cycle type of σ, and write an element of S_9 which is conjugated to σ.

Solution. One can check that

$$1 \to 2 \to 3 \to 4 \to 5 \to 1, \quad 8 \to 9 \to 8, \tag{3.39}$$

and $6, 7$ are fixed points. Therefore,

$$\sigma = (1\ 2\ 3\ 4\ 5)\,(6)\,(7)\,(8\ 9)\,. \tag{3.40}$$

So its cycle-type is $(5, 2, 1, 1)$. Any element of this cycle-type will be conjugate to σ, i.e. any

$$\tau = (a_1\ a_2\ a_3\ a_4\ a_5)\,(a_6)\,(a_7)\,(a_8\ a_9) \tag{3.41}$$

is conjugate to σ. ■

Exercise 3.12. In the following, verify if the maps $\varphi : G \to G$ are homomorphisms of groups. If they are, determine their kernel.

(a) $G = (\mathbb{R} \setminus \{0\},\ \cdot\ )$, the nonzero real numbers with multiplication, $\varphi(x) = x^2$.

(b) $G = (\mathbb{R}, +)$, the real numbers with addition. $\varphi(x) = x + 1$.

(c) G is an abelian group, $\varphi(x) = x^3$.

Solution. (a) Yes, $\varphi : \mathbb{R}^\times \to \mathbb{R}^\times$, $x \mapsto x^2$ is a homomorphism, since

$$\varphi(xy) = (xy)^2 = x^2y^2 = \varphi(x)\,\varphi(y). \tag{3.42}$$

Its kernel is

$$\operatorname{Ker}\varphi = \left\{x \in \mathbb{R}^\times \mid \varphi(x) = 1\right\} = \left\{x \in \mathbb{R}^\times \mid x^2 = 1\right\} = \{1, -1\}. \tag{3.43}$$

(b) $\varphi : (\mathbb{R}, +) \to (\mathbb{R}, +)$, $x \mapsto x + 1$ is not a homomorphism, since $\varphi(0) \neq 0$.

(c) If G is abelian, $\varphi : G \to G$, $x \mapsto x^3$ is a homomorphism, since

$$\varphi(xy) = (xy)^3 = (xy)(xy)(xy) = (xxx)(yyy) = x^3y^3 = \varphi(x)\,\varphi(y). \tag{3.44}$$

Its kernel is

$$\operatorname{Ker}\varphi = \left\{x \in G \mid x^3 = e\right\} = \{x \in G \mid \operatorname{ord}(x) = 1 \text{ or } 3\}. \tag{3.45}$$

■

Exercise 3.13. Let G be a group, let $\varphi : G \to G$ be an automorphism, and let $Z(G)$ be the center of G. Show

$$\varphi(Z(G)) \subseteq Z(G). \tag{3.46}$$

Solution. For $z \in Z(G)$, z commutes with every $x \in G$. Hence,

$$zx = xz \implies \varphi(zx) = \varphi(xz)\varphi(z)\varphi(x) = \varphi(x)\varphi(z). \tag{3.47}$$

Since φ is an automorphism, given any $g \in G$, we can find $x \in G$ such that $\varphi(x) = g$. Hence, for any $g \in G$,

$$\varphi(z)\ g = g\ \varphi(z), \tag{3.48}$$

i.e. $\varphi(z) \in Z(G)$, proving that $\varphi(Z(G)) \subseteq Z(G)$. ■

In fact, $\varphi(Z(G)) = Z(G)$. The other containment can be shown using the fact that φ^{-1} is also an automorphism, so $\varphi^{-1}(Z(G)) \subseteq Z(G)$. As a result,

$$Z(G) = \varphi\left(\varphi^{-1}(Z(G))\right) \subseteq \varphi(Z(G)). \tag{3.49}$$

Exercise 3.14. Let G be a cyclic group of prime order. Determine $\operatorname{Aut}(G)$.

Solution. For $G = \mathbb{Z}_p$, we claim that $\mathrm{Aut}(G) \cong \mathbb{Z}_p^\times$. Define

$$\begin{aligned} \varphi : \mathrm{Aut}\left(\mathbb{Z}_p\right) &\to \mathbb{Z}_p^\times \\ f &\mapsto f(1). \end{aligned} \tag{3.50}$$

This is a homomorphism since

$$\varphi\left(f \circ g\right) = \left(f \circ g\right)(1) = f\left(g(1)\right) = f\left(g(1) \cdot 1\right) = g(1)f(1) = f(1)g(1) = \varphi(f)\varphi(g). \tag{3.51}$$

It's injective because if $f \in \operatorname{Ker}\varphi$, then $f\left(1\right) = 1$, and then $f\left(a\right) = a$ for each $a \in \mathbb{Z}_p$, i.e. $f = \mathrm{id}$.

Furthermore, φ is surjective, because given any $b \in \mathbb{Z}_p^\times$, we can define an automorphism $f\left(a\right) = ab$, for which $f\left(1\right) = b$. Hence, φ is an isomorphism. ■

Exercise 3.15. If $G/Z(G)$ is cyclic, then G is abelian.

Solution. For simplicity of notations, we'll denote $N := Z(G)$. Since G/N is cyclic,

$$G/N = \langle xN \rangle = \left\{N, xN, x^2N, \ldots, x^nN\right\}. \tag{3.52}$$

Let's take any two element $g, h \in G$. Since the cosets in G/N partition G, we must have that $g = x^i z_1$ and $h = x^j z_2$ for some $z_1, z_2 \in N = Z(G)$. Then

$$\begin{aligned} gh &= \left(x^i z_1\right)\left(x^j z_2\right) = x^i\left(z_1 x^j\right) z_2 && (3.53)\\ &= x^i\left(x^j z_1\right) z_2 = x^j x^i\left(z_1 z_2\right) && (3.54)\\ &= x^j\left[\left(x^i z_1\right) z_2\right] = x^j\left[z_2\left(x^i z_1\right)\right] && (3.55)\\ &= \left(x^j z_2\right)\left(x^i z_1\right) = hg. && (3.56) \end{aligned}$$

Hence, G is abelian. ■

Exercise 3.16. An r-cycle is even if and only if r is odd.

Solution.

$$\left(a_1\ a_2\ \cdots a_r\right) = \left(a_1\ a_2\right)\left(a_2\ a_3\right)\cdots\left(a_{r-1}\ a_r\right), \tag{3.57}$$

so a r-cycle is a product of $r-1$ transitions. So it is even if and only if $r-1$ is even, i.e. r is odd. ■

Exercise 3.17. Let G be a group of order 30.

(a) Show a 3-Sylow or a 5-Sylow subgroup is normal in G.

(b) Show every 3-Sylow and every 5-Sylow subgroup is normal in G.

(c) Show G has a normal subgroup of order 15.

Solution. We prove some preliminary result first.

Claim 3.1: If $N \trianglelefteq G$, and $H \trianglelefteq N$ is a p-Sylow subgroup of N, then $H \trianglelefteq G$.

Proof. Since $H \trianglelefteq N$, $n_p(N) = 1$. Then for any $g \in G$,

$$gHg^{-1} \subseteq gNg^{-1} = N. \tag{3.58}$$

Hence, gHg^{-1} is a p-Sylow subgroup of N, since $\left|gHg^{-1}\right| = |H|$. But since $n_p(N) = 1$, we must have that $gHg^{-1} = H$, for any $g \in G$. As a result, $H \trianglelefteq G$. □

(a) Assume for the sake of contradiction that neither a 3-Sylow subgroup nor a 5-Sylow subgroup is normal. Then $n_3, n_5 > 1$.

Since $n_5 \mid 6$ and $n_5 \equiv 1 \pmod 5$, then we must have $n_5 = 6$. Therefore, there are $6 \times 4 = 24$ elements of order 5 (since the 5-Sylow subgroups intersect trivially, because the non-identity elements generate the whole subgroup since it's a cyclic group). Hence,

$$2n_3 = (\text{number of elements of order } 3) \leq 30 - 24 - 1 = 5, \tag{3.59}$$

implying that $n_3 \leq 2$. Since $n_3 \equiv 1 \pmod 3$, we must have $n_3 = 1$.

(b) We have shown earlier that $n_3 = 1$. Now we'll show that $n_5 = 1$. Let H_3 be the 3-Sylow subgroup and H_5 be a 5-Sylow subgroup. Consider $H = H_3H_5 \leq G$. Then

$$|H| = \frac{|H_3|\,|H_5|}{|H_3 \cap H_5|} = 15, \tag{3.60}$$

and from the proof of Exercise 3.24, we conclude that H is cyclic. Since $[G : H] = 2$, H is normal in G. Furthermore, since H is cyclic, it has a unique 5-Sylow subgroup, which is H_5. Therefore, by Claim 3.1, $H_5 \trianglelefteq G$, i.e. $n_5(G) = 1$.

(c) As before, we consider the subgroup $H = H_3H_5 \leq G$, which has order 15, i.e. its index is 2. Therefore, it's normal.

■

Exercise 3.18. If G is a group of order 385, show that its 11-Sylow subgroup is normal and its 7-Sylow subgroup is in the center of G.

Solution. $|G| = 385 = 5 \cdot 7 \cdot 11$. $n_{11} \mid 35$, so

$$n_{11} \in \{1, 5, 7, 35\}, \tag{3.61}$$

and $n_{11} \equiv 1 \pmod{11}$ forces that we must have $n_{11} = 1$. Hence, the 11-Sylow subgroup is normal. Now, $n_7 \mid 55$ gives

$$n_7 \in \{1, 5, 11, 55\}, \tag{3.62}$$

and $n_7 \equiv 1 \pmod 7$ forces that we must have $n_7 = 1$. Hence, the 7-Sylow subgroup H is normal.

Claim 3.2: If $H \trianglelefteq G$, then $C_G(H) \trianglelefteq G$ and $G/C_G(H)$ is isomorphic to a subgroup of $\mathrm{Aut}(H)$.

Proof. Consider the action of G on H by conjugation. This gives rise to a group homomorphism

$$\begin{aligned} \Psi : G &\to \mathrm{Aut}(H) \\ g &\mapsto \left(h \mapsto ghg^{-1}\right). \end{aligned} \tag{3.63}$$

Then

$$\begin{aligned} \mathrm{Ker}\,\Psi &= \left\{g \in G \mid ghg^{-1} = h \text{ for all } h \in H\right\} && (3.64) \\ &= \{g \in G \mid gh = hg \text{ for all } h \in H\} = C_G(H). && (3.65) \end{aligned}$$

Hence, $C_G(H) \trianglelefteq G$ and by the First isomorphism theorem, $G/C_G(H)$ is isomorphic to $\mathrm{im}\,\Psi \leq \mathrm{Aut}(H)$. □

Now, by Claim 3.2, $G/C_G(H)$ is isomorphic to a subgroup of $\mathrm{Aut}(H)$. Since $H \cong \mathbb{Z}_7$, by Exercise 3.14, $|\mathrm{Aut}(H)| = 6$. Therefore,

$$|G/C_G(H)| \,\Big|\, |\mathrm{Aut}(H)| = 6. \tag{3.66}$$

But $|G|$ is coprime with 6, and hence so is $|G/C_G(H)| = \frac{|G|}{|C_G(H)|}$. Therefore,

$$|G/C_G(H)| = 1 \implies G = C_G(H). \tag{3.67}$$

So every element of G commutes with every element of H, proving that $H \leq Z(G)$. ■

Alternate Solution. We can also finish without using Claim 3.2. As before, we show $n_7 = 1$, and hence, the 7-Sylow subgroup H is normal.

Claim 3.3: The size of conjugacy class divides $|G|$.

Proof. Let G act on itself by conjugation. Then $\mathrm{Orb}(g)$ is the conjugacy class $\mathrm{Conj}_G(g)$ of g. Then by Orbit-Stabilizer theorem,

$$|G| = |\mathrm{Orb}(g)|\,|\mathrm{Stab}(g)| = |\mathrm{Conj}_G(g)|\,|\mathrm{Stab}(g)|\,. \tag{3.68}$$

□

Since normal subgroups consist of conjugacy classes (i.e. if one element of a conjugacy class is inside the normal subgroup, then so is the entire conjugacy class), the 7-Sylow subgroup H

consists of conjugacy classes. The possible size of the conjugacy classes are the divisors of 385, so the possible sizes are

$$1, 5, 7, 11, \ldots. \tag{3.69}$$

If H contains a nontrivial conjugacy class (i.e. conjugacy class of size > 1), then it must be of size 5. But then there is a non-identity element $g \in H$ which has trivial conjugacy class, i.e. it's in the center.

Since $H \cong \mathbb{Z}_7$, every non-identity element generates H, so $g \in Z(G)$ implies $H = \langle g \rangle \subseteq Z(G)$. ■

Exercise 3.19. Let G be a group of order 231. Show the 11-Sylow subgroup is in the center of G.

Solution. $|G| = 231 = 3 \cdot 7 \cdot 11$. Now, $n_{11} \mid 21$ gives

$$n_7 \in \{1, 3, 7, 21\}, \tag{3.70}$$

and $n_{11} \equiv 1 \pmod{11}$ forces that we must have $n_{11} = 1$. Hence, the 11-Sylow subgroup H is normal. Now, by Claim 3.2, $G/C_G(H)$ is isomorphic to a subgroup of $\mathrm{Aut}(H)$. Since $H \cong \mathbb{Z}_{11}$, by Exercise 3.14, $|\mathrm{Aut}(H)| = 10$. Therefore,

$$|G/C_G(H)| \,\Big|\, |\mathrm{Aut}(H)| = 10. \tag{3.71}$$

But $|G|$ is coprime with 10, and hence so is $|G/C_G(H)| = \frac{|G|}{|C_G(H)|}$. Therefore,

$$|G/C_G(H)| = 1 \implies G = C_G(H). \tag{3.72}$$

So every element of G commutes with every element of H, proving that $H \leq Z(G)$. ■

Alternate Solution. As before, we show $n_{11} = 1$, and hence, the 11-Sylow subgroup H is normal.

> **Claim 3.4:** If $H \trianglelefteq G$ is a cyclic normal subgroup and $|H| = p$ is prime, then for any two non-identity elements $x, y \in H$, $|\mathrm{Conj}_G(x)| = |\mathrm{Conj}_G(y)|$.
>
> *Proof.* If x, y belong to the same conjugacy class, then we are done. Else, we define a function $f : \mathrm{Conj}_G(x) \to \mathrm{Conj}_G(y)$ as follows: since $H = \langle x \rangle$, $y = x^k$ for some $0 < k < p$. So we define $f(z) = z^k$ for every $z \in \mathrm{Conj}_G(x)$. This is well-defined since
>
> $$f\left(gxg^{-1}\right) = \left(gxg^{-1}\right)^k = gx^k g^{-1} = gyg^{-1} \in \mathrm{Conj}_G(y). \tag{3.73}$$
>
> This is surjective, since any $gyg^{-1} \in \mathrm{Conj}_G(y)$ is just the image of gxg^{-1}. Therefore, $|\mathrm{Conj}_G(x)| \geq |\mathrm{Conj}_G(y)|$. Switching x and y in the above argument, we obtain $|\mathrm{Conj}_G(y)| \geq |\mathrm{Conj}_G(x)|$. □

The possible size of the conjugacy classes are the divisors of 231, so the possible sizes are

$$1, 3, 7, 11, \ldots. \tag{3.74}$$

The conjugacy classes of the non-identity elements are all the same. Hence, the size of their conjugacy class must divide $|H| - 1 = 10$. The only common element of divisors of 10 and divisors of $|G| = 231$ is 1. Hence, all non-identity elements are in a singleton conjugacy class, i.e. they are in the center! ■

Exercise 3.20. If G is a group of order p^2q with p and q primes, show that G has a nontrivial normal subgroup.

Solution. We'll first consider the case $p^2q = 12$. Assume the contrary. Note that $12 = 2^2 \times 3$. Since $n_3 \mid 4$ and $n_3 > 1$ (since we assumed the group is simple) and $n_3 \equiv 1 \pmod 3$, we must have that $n_3 = 4$. This gives

$$(\text{number of elements of order } 3) = n_3(3-1) = 4 \cdot 2 = 8. \tag{3.75}$$

The 2-Sylow subgroups can't intersect any of the 3-Sylow subgroups. The identity element together with the other 3 elements form one 2-Sylow subgroup. Therefore, we don't have any room for any other 2-Sylow subgroups, implying $n_2 = 1$.

Now, we consider the general case: Assume for the sake of contradiction that ther exists a simple group G with $|G| = p^2q$. Note that

$$p + 1 \leq n_p \leq q. \tag{3.76}$$

The first bound follows from the fact that $n_p \neq 1$, and the second bound is given by $n_p \mid q$. Now, note that $n_q \mid p^2$ and $n_q > q > p$, so we must have $n_q = p^2$. This gives us

$$p^2 \equiv 1 \pmod q \implies q \mid (p-1)(p+1). \tag{3.77}$$

Since $q > p$, the scenario $q \mid p-1$ is impossible, so we must have $q \mid p+1$. But that way $q \leq p+1$. But we already know $q \geq p+1$. Therefore, $q = p+1$. This can only happen if $p = 2$, $q = 3$, i.e. $G = 12$, which has already been done! ■

Exercise 3.21. How many elements of order 5 are contained in a group of order 20?

Solution. Since $n_5 \mid 4$ and $n_5 \equiv 1 \pmod 5$, we must have $n_5 = 1$. Therefore, there exists a unique 5-Sylow subgroup. Hence, the number of elements of order 5 is 4. ■

Exercise 3.22. Prove: No group of order pq, with p and q prime numbers, is simple.

Solution. Assume for the sake of contradiction that ther exists a simple group G with $|G| = pq$. WLOG, $p < q$. Then $n_q \mid p$. But $n_q = p$ is not possible since $n_q > 1$ along with

$$n_q \equiv 1 \pmod{q} \tag{3.78}$$

would imply

$$n_q \geq q + 1 > p. \tag{3.79}$$

Hence, $n_q = 1$, and hence the q-Sylow subgroup is normal. ■

Exercise 3.23. Let G be a group of order $p^m r$. Show G contains a subgroup of order p^e for every $e \leq m$.

Solution. We proceed by induction on e. The base case $e = 0$ is trivial. Suppose H is a subgroup of order $p^e < p^m$, we shall construct a subgroup of order p^{e+1}.

Claim 3.5 (p-group lemma)**:** If a p-group (group of order p^n) acts on a set S, then if N is number of fixed points (i.e. singleton orbits) of this action,

$$N \equiv |S| \pmod{p}. \tag{3.80}$$

Proof. By Orbit-Stabilizer theorem, the possible orbit sizes are $1, p, p^2, \ldots, p^n$. Then

$$|S| = \sum_{\text{orbits}} |\mathrm{Orb}(x)| \equiv \sum_{\text{singleton orbits}} |\mathrm{Orb}(x)| = N \pmod{p}. \tag{3.81}$$

□

Claim 3.6 (Normalizer lemma)**:** If H is a p-subgroup of G, then

$$[N_G(H) : H] \equiv [G : H] \pmod{p}. \tag{3.82}$$

Proof. Let H act on the left cosets G/H by left multiplication

$$h \cdot (gH) = (hg)\, H. \tag{3.83}$$

gH is a fixed-point of this action if and only if

$$\begin{aligned} h \cdot (gH) = gH \text{ for all } h \in H &\iff \left(g^{-1}hg\right) H = H \text{ for all } h \in H && (3.84)\\ &\iff g^{-1}hg \in H \text{ for all } h \in H && (3.85)\\ &\iff g \in N_G(H). && (3.86) \end{aligned}$$

Therefore, the fixed points are precisely the cosets in $N_G(H)/H$. As a result, the number of fixed points is $[N_G(H) : H]$, and hence, by p-group lemma, $[N_G(H) : H] \equiv [G : H] \pmod{p}$. □

Now, since $H \trianglelefteq N_G(H)$, consider the quotient group $N_G(H)/H$. Since $p \mid [G:H] = p^{m-e}r$, by Normalizer lemma, $p \mid |N_G(H)/H|$. Therefore, by Cauchy's theorem, we can find an element of order p in $N_G(H)/H$.

Say $aH \in N_G(H)/H$ is an element of order p. Then $a^p \in H$, since it's of order p. Now, define

$$H' = \bigcup_{i=0}^{p-1} a^i H. \tag{3.87}$$

Then $H' \leq G$, because if we take $a^i h, a^j h_\prime \in H'$ for $h, h_\prime \in H$,

$$\left(a^i h\right)\left(a^j h_\prime\right)^{-1} = a^i h\ h_\prime^{-1} a^{-j} = a^i\ a^{-j}\ h' \in H' \tag{3.88}$$

for some $h' \in H$, since $a^{-j} \in N_G(H)$. So $H' \leq G$. Finally,

$$\left|H'\right| = \sum_{i=0}^{p-1} \left|a^i H\right| = \sum_{i=0}^{p-1} |H| = p\,|H| = p^{e+1}. \tag{3.89}$$

■

Exercise 3.24. Classify the groups of order 33.

Solution. We shall prove a general statement:

If G is a group of order pq and $p < q$ primes with $q \not\equiv 1 \pmod{p}$, then $G \cong \mathbb{Z}_{pq}$.

By the proof of Exercise 3.22, $n_q = 1$. $n_p \mid q$ and $q \not\equiv 1 \pmod{p}$ implies $n_p = 1$.

Therefore, $H_1 = \langle a \rangle, H_2 = \langle b \rangle$, the p-Sylow subgroup and the q-Sylow subgroup, respectively, are normal in G. Their intersection is trivial, since their orders are coprime. Therefore, a and b commute, since

$$aba^{-1}b^{-1} = \left(aba^{-1}\right)b^{-1} = a\left(ba^{-1}b^{-1}\right) \in H_1 \cap H_2 = \{e\}. \tag{3.90}$$

Now, a and b commute, their orders are coprime, so $\operatorname{ord}(ab) = pq$. Indeed,

$$(ab)^n = a^n b^n = e \iff a^n = b^{-n} \in H_1 \cap H_2 = \{e\} \tag{3.91}$$

$$\iff a^n = b^n = e \tag{3.92}$$

$$\iff p \mid n \text{ and } q \mid n. \tag{3.93}$$

So $n = pq$ is the smallest value to have $(ab)^n = e$. Now since $\operatorname{ord}(ab) = pq$, G is cyclic!

Now, for $|G| = 33 = 3 \times 11$, and $11 \not\equiv 1 \pmod{3}$, so any group of order 33 is cyclic. ■

§4 Problems on Rings from Lothar Göttsche's Notes

Exercise 4.1. Let R be a ring with 1, and let $a \in R$ be a unit. Show that there is a *unique* $b \in R$ with $ab = ba = 1$.

Solution. If a is a unit, there exists $b \in R$ such that $ab = ba = 1$ (by definition). To show uniqueness, suppose another $b' \in R$ exists, with $ab' = b'a = 1$. Then

$$b = b \cdot 1 = b\left(ab'\right) = (ba)\, b' = 1 \cdot b' = b'. \tag{4.1}$$

So the b is unique! ■

Exercise 4.2. Show that the units in $\mathbb{Z}[i]$ are $\{1, -1, i, -i\}$.

Solution. Let $a + ib \in \mathbb{Z}[i]$ be a unit. Then

$$(a + ib)(c + id) = 1 \tag{4.2}$$

for some $c + id \in \mathbb{Z}[i]$. Taking modulus squared on both sides, we get

$$\left(a^2 + b^2\right)\left(c^2 + d^2\right) = 1. \tag{4.3}$$

$a^2 + b^2 \in \mathbb{Z}_{\geq 0}$ that divides 1. Therefore, it has to be equal to 1. Now,

$$a^2 + b^2 = 1 \implies (a, b) = (\pm 1, 0), (0, \pm 1), \tag{4.4}$$

which corresponds to $1, -1, i, -i$. It's easy to check that they are units indeed. ■

Exercise 4.3. Show: The odd integers with usual addition and multiplication are not a ring.

Solution. The additive identity is not in the set, so it's not an abelian group with respect to addition. ■

Exercise 4.4. Let R be a ring and $a, b \in R$. Show $(a + b)^2 = a^2 + ab + ba + b^2$ (using only the definitions).

Solution.

$$(a + b)^2 = (a + b)(a + b) = (a + b)\, a + (a + b)\, b = a^2 + ba + ab + b^2. \tag{4.5}$$

■

Exercise 4.5. An element a in a ring R is called *nilpotent* if there is an $n \in \mathbb{Z}_{>0}$ with $a^n = 0$. Show: if a is nilpotent, then $1 + a$ is a unit.

Solution. We can assume WLOG that n is odd, because if n is even, $a^{n+1} = a^n \cdot a = 0 \cdot a = 0$, and we can just replace n by $n+1$.

$$1 = 1 + a^n = (1+a)\left(1 - a + a^2 - a^3 + \cdots + a^{n-1}\right). \tag{4.6}$$

■

Exercise 4.6. Show: In a commutative ring R, $a+b$ is nilpotent if a and b are nilpotent. Show that this may be false if R is not commutative.

Solution. Suppose $a^n = b^m = 0$ for $n, m \in \mathbb{Z}_{>0}$. Since the ring is commutative, we can apply Binomial formula:

$$(a+b)^{m+n} = \sum_{i=0}^{m+n} \binom{m+n}{i} a^i b^{m+n-i}. \tag{4.7}$$

If $i < m$, $m+n-i \geq n$, so the term $b^{m+n-i} = 0$. If $i \geq m$, then $a^i = 0$. Thus all the summands are 0, and hence $(a+b)^{m+n} = 0$.

If R is not commutative, say $R = \mathrm{Mat}_{2\times 2}$, then this assertion may not be true. For example, take

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}. \tag{4.8}$$

Then $A^2 = B^2 = 0$. But

$$A + B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \tag{4.9}$$

is an invertible element, so it can't be nilpotent. ■

Exercise 4.7. Show that a commutative ring R is an integral domain if and only if for all $a, b, c \in R$ with $a \neq 0$, the relation $ab = ac$ implies $b = c$.

Solution. Suppose R is an ID. Let $a, b, c \in R$ with $a \neq 0$. Then

$$ab = ac \implies a(b-c) = ab - ac = 0 \implies b - c = 0 \implies b = c. \tag{4.10}$$

Conversely, suppose for all $a, b, c \in R$ with $a \neq 0$, the relation $ab = ac$ implies $b = c$. Suppose $xy = 0$ for $x, y \in R$. We need to show that this implies $x = 0$ or $y = 0$. If $x \neq 0$, then

$$xy = 0 = x \cdot 0 \implies y = 0. \tag{4.11}$$

Therefore, R is an ID. ■

Exercise 4.8. Let U be a set, and let R be the set of subsets of U. Addition and multiplication on R are defined by

$$A + B := A \cup B, \qquad A \cdot B := A \cap B. \tag{4.12}$$

Prove or disprove: R a ring.

Solution. No. Let's first examine what's the additive identity.

$$A = A + 0 = A \cup 0. \tag{4.13}$$

for all $A \subseteq U$, so we have that $0 \subseteq A$ for all A. Then we must have $0 = \varnothing$.

Given any set A, we need to have its additive inverse, i.e. another set B such that $A + B = 0$. This means

$$A \cup B = \varnothing, \tag{4.14}$$

which only happens if $A = B = \varnothing$. Therefore, this does **NOT** form a ring since the nonempty elements don't have additive inverses. ■

Exercise 4.9. Let U be a set, and let R be the set of subsets of U. Addition and multiplication on R are defined by

$$A + B := (A \setminus B) \cup (B \setminus A), \qquad A \cdot B := A \cap B. \tag{4.15}$$

Prove or disprove: R a ring.

Solution. Yes, R is a ring.

- Addition is associative: since $A \triangle B$ denotes the elements that belong to exactly one of the sets A or B. Therefore, $(A \triangle B) \triangle C$ denotes the elements that belong to exactly one of the sets $A \triangle B$ or C.

  If an element of $(A \triangle B) \triangle C$ belongs to C, it's not in $A \triangle B$, so it's in both A, B or neither of A, B. On the other hand, if an element is not in C, then it's in $(A \triangle B)$.

  Therefore, $(A \triangle B) \triangle C$ consists of elements that belong to exactly one of A, B, C or all three. Similarly, $A \triangle (B \triangle C)$ also consists of elements that belong to exactly one of A, B, C or all three. Hence, they are the same!

- Additiion is clearly commutative.

- $\varnothing$ is the additive identity, since

$$A + \varnothing = (A \setminus \varnothing) \cup (\varnothing \setminus A) = A. \tag{4.16}$$

- Every set is its own additive inverse, since

$$A + A := (A \setminus A) \cup (A \setminus A) = \varnothing. \tag{4.17}$$

- Multiplication is clearly associative and commutative.

- U is the multiplicative identity, since

$$A \cdot U = A \cap U = A. \tag{4.18}$$

- We have to verify the distributive law:

$$A \cap (B \triangle C) = (A \cap B) \triangle (A \triangle C). \tag{4.19}$$

  This is clearly true, since $A \cap (B \triangle C)$ consists of elements that are in A and exactly one of B or C. That means they are in $A \cap B$ or $A \cap C$ (not both), and those elements are precisely the ones that make up $(A \cap B) \triangle (A \triangle C)$.

Therefore, R is a commutative unital ring. ■

Exercise 4.10. Let $\mathrm{GL}(2, \mathbb{R}) := \{A \in \mathrm{Mat}_{2\times 2}(\mathbb{R}) \mid \det(A) \neq 0\}$. Is $\mathrm{GL}(2, \mathbb{R})$ a subring of $M_{2\times 2}(\mathbb{R})$?

Solution. No, because $\mathrm{GL}(2, \mathbb{R})$ is not closed under addition, so it's not an additive subgroup of $\mathrm{Mat}_{2\times 2}(\mathbb{R})$. Indeed,

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \tag{4.20}$$

where both the summands are in $\mathrm{GL}(2, \mathbb{R})$, but the sum is not. ■

Exercise 4.11. Let R be a ring with $a^2 = a$ for all $a \in R$. Show: R is commutative.

Solution. Since $(a+b)^2 = a + b$, we have

$$a + b = (a+b)^2 = a^2 + b^2 + ab + ba = a + b + ab + ba \implies ab + ba = 0. \tag{4.21}$$

Then

$$2a = (2a)^2 = 4a^2 = 4a \implies a + a = 0 \implies a = -a. \tag{4.22}$$

Plugging it into $ab + ba = 0$, we get

$$ab = -ba = -b(-a) = ba. \tag{4.23}$$

■

Exercise 4.12. Determine the group of units in $\mathbb{Z}_{12}$.

Solution. If $a \in \mathbb{Z}_{12}$ is a unit, then there exists some $b \in \mathbb{Z}_{12}$ such that $ab = 1$, i.e.

$$ab = 1 + 12k \text{ for some } k \in \mathbb{Z}. \tag{4.24}$$

If $d = \gcd(a, 12)$, then it divides both ab and $12k$, and hence it divides 1. So $d = 1$. And, if $d = 1$, by Bezout's identity, there exists $b, k \in \mathbb{Z}$ such that

$$ab - 12k = 1. \tag{4.25}$$

Hence, $ab = 1$ in $\mathbb{Z}_{12}$. Therefore, $\mathbb{Z}_{12}^{\times} = \{1, 5, 7, 11\}$, all the numbers smaller than 12 that are coprime to 12. ■

Exercise 4.13. Let R be a ring and let $Z(R) := \{x \in R \mid xy = yx \text{ for all } y \in R\}$. Show that $Z(R)$ is a subring of R. $Z(R)$ is usually called the center of R.

Solution. Clearly $0 \in Z(R)$, since $0 \cdot y = y \cdot 0 = 0$ for all $y \in R$. We need to show that given $a, b \in Z(R)$, $a - b, ab \in Z(R)$. For any $y \in R$, since $a, b \in Z(R)$,

$$(a - b)\, y = ay - by = ya - yb = y\,(a - b)\,, \tag{4.26}$$

so $a - b \in Z(R)$.

$$(ab)\, y = a\,(by) = a\,(yb) = (ay)\, b = (ya)\, b = y\,(ab)\,, \tag{4.27}$$

so $ab \in Z(R)$. Hence, $Z(R)$ is a subring. ■

Exercise 4.14. If R is a division ring, show that $Z(R)$ is a field.

Solution. As before, $Z(R)$ is a subring of R. $Z(R)$ is also a division ring, for nonzero $a \in Z(R)$,

$$ay = ya \implies a^{-1}\,(ay)\, a^{-1} = a^{-1}\,(ya)\, a^{-1} \implies ya^{-1} = a^{-1}y, \tag{4.28}$$

for every $y \in R$, so $a^{-1} \in Z(R)$. Therefore, $Z(R)$ is a division ring. Now, we just need to show that the multiplication is commutative. It's trivial since the elements of $Z(R)$ commute with every element of R, so in particular, they commute with elements from $Z(R)$. ■

Exercise 4.15. Let R be an integral domain and $a, b \in R$. Assume $a^n = b^n$ and $a^m = b^m$ for two relatively prime integers n, m. Show that $a = b$.

Solution. Since $\gcd(m, n) = 1$, by Bezout's identity, there exist $x, y \in \mathbb{Z}$ such that

$$mx + ny = 1. \tag{4.29}$$

WLOG $x > 0$ and $y < 0$. Then

$$a^m = b^m \implies (a^m)^x = (b^m)^x \implies a^{mx} = b^{mx}. \tag{4.30}$$

Similarly, $a^{-ny} = b^{-ny}$. Therefore,

$$a \cdot a^{-ny} = a^{1-ny} = a^{mx} = b^{mx} = b^{1-ny} = b \cdot b^{-ny}. \tag{4.31}$$

If $a^{-ny} = b^{-ny} = 0$, then $a = b = 0$ since we're in an integral domain. If $a^{-ny} = b^{-ny} \neq 0$, we may cancel it out to get $a = b$. ■

Exercise 4.16. Let m be an integer which is not divisible by the square of an integer > 1.

(a) Show that $\mathbb{Z}[\sqrt{m}] := \{a + b\sqrt{m} \mid a, b \in \mathbb{Z}\}$ is an integral domain.

(b) Find the units in $\mathbb{Z}[\sqrt{m}]$ for all $m < 0$.

Solution. (a) $\mathbb{Z}[\sqrt{m}]$ is a subring of $\mathbb{C}$, which is a field and hence it has no nonzero zero divisors. Therefore, $\mathbb{Z}[\sqrt{m}]$ also can't have nonzero zero divisors. Hence, it's an integral domain.

(b) We have already shown that for $m = -1$,

$$\mathbb{Z}\left[\sqrt{-1}\right]^\times = \mathbb{Z}[i]^\times = \{1, -1, i, -i\}. \tag{4.32}$$

Now let's assume $m = -n$ for $n > 1$. Suppose $a + b\sqrt{m} = a + ib\sqrt{n}$ is a unit. Then there exists $c, d \in \mathbb{Z}$ such that

$$\left(a + ib\sqrt{n}\right)\left(c + id\sqrt{n}\right) = 1. \tag{4.33}$$

Taking norm-squared on both sides, we get

$$\left(a^2 + nb^2\right)\left(c^2 + nd^2\right) = 1. \tag{4.34}$$

$a^2 + nb^2$ is a non-negative integer that divides 1, hence it is 1. Now, if $b \neq 0$, then

$$a^2 + nb^2 \geq nb^2 \geq n > 1. \tag{4.35}$$

Therefore, $b = 0$. But that just means $a^2 = 1$, i.e. $a = \pm 1$. Therefore, for $m = -n$ with $n > 1$,

$$\mathbb{Z}\left[\sqrt{m}\right]^\times = \{1, -1\}. \tag{4.36}$$

■

Exercise 4.17. Show: A finite ring with 1 (not necessarily commutative) is a division ring.

Solution. This is false! Consider

$$\mathbb{Z}_4 = \{0, 1, 2, 3\}. \tag{4.37}$$

This is a finite ring with 1 (and also commutative). But this is **NOT** a division ring since 2 doesn't have any multiplicative inverse. I think the statement should be

A finite ring with 1, which doesn't have any nonzero zero-divisors, is a division ring.

Take any $a \in R$ with $a \neq 0$. Consider the set

$$\left\{a, a^2, a^3, \ldots\right\}. \tag{4.38}$$

Since our ring is finite, there must be a repitition, i.e. $a^m = a^n$ for $m < n$. After a bit of rearrangement, we get

$$0 = a^n - a^m = a^m\left(a^{n-m} - 1\right). \tag{4.39}$$

Since $a \neq 0$, $a^m \neq 0$ (this holds because the ring doesn't have any nonzero zero-divisors). Therefore,

$$a^{n-m} - 1 = 0 \implies a^{n-m} = 1 \implies a \cdot a^{n-m-1} = a^{n-m-1} \cdot a = 1, \tag{4.40}$$

implying that $a^{n-m-1} = a^{-1}$. ■

Exercise 4.18. Let R be a commutative ring with 1. Determine the units in $R[x]$.

Solution. We claim that $f(x) = \sum_{i=0}^{n} a_i x^i \in R[x]$ is a unit if and only if a_0 is a unit and a_i is nilpotent for all $1 \leq i \leq n$.

Let's prove the "if" direction first. Since $R \leq R[x]$, nilpotents in R are also nilpotents in $R[x]$. Since nilpotent elements form an ideal in a commutative ring (Exercise 4.33), $\sum_{i=1}^{n} a_i x^i$ is nilpotent in $R[x]$. If n is a nilpotent and u is a unit, then $u + n$ is a unit, since

$$u + n = u\left(1 + u^{-1}n\right), \tag{4.41}$$

and $u^{-1}n$ is a nilpotent (since nilpotents form an ideal, Exercise 4.33), so $1 + u^{-1}n$ is a unit by Exercise 4.5. Since units form a group under multiplication, $u\left(1 + u^{-1}n\right)$ is a unit. Now, a_0 is a unit, and $\sum_{i=1}^{n} a_i x^i$ is nilpotent, so their sum $f(x) = \sum_{i=0}^{n} a_i x^i \in R[x]$ is a unit.

Conversely, suppose $f(x) = \sum_{i=0}^{n} a_i x^i \in R[x]$ is a unit. Let $g(x) = \sum_{j=1}^{m} b_j x^j$ is its inverse. Then

$$1 = f(x)g(x) = \left(\sum_{i=0}^{n} a_i x^i\right)\left(\sum_{j=1}^{m} b_j x^j\right) = \sum_{k=0}^{m+n}\left(\sum_{i+j=k} a_i b_j\right) x^k \tag{4.42}$$

Equating the coefficients of x^0 on both sides, we get $a_0 b_0 = 1$, i.e. a_0 is a unit (and so is b_0). Equating the coefficients of x^{m+n} on both sides, we get

$$a_n b_m = 0. \tag{4.43}$$

We claim that $a_n^{l+1} b_{m-l} = 0$ for each $0 \leq l \leq m$. We prove it by induction on l. The base case $l = 0$ is shown above. Suppose it holds for $0, 1, \ldots, l-1$. Looking at the coefficient of x^{m+n-l}, we get

$$a_n b_{m-l} + a_{n-1} b_{m-l+1} + a_{n-2} b_{m-l+2} + \cdots + a_{n-l} b_m = 0. \tag{4.44}$$

Multiplying by a_n^l, we have

$$0 = a_n^{l+1} b_{m-l} + a_{n-1} a_n^l b_{m-l+1} + a_{n-2} a_n^l b_{m-l+2} + \cdots + a_{n-l} a_n^l b_m. \tag{4.45}$$

By inductive hypothesis, all $a_n^l b_{m-(l-j)} = 0$. Therefore, $a_n^{l+1} b_{m-l} = 0$. So we have proved that $a_n^{l+1} b_{m-l} = 0$ for each $0 \leq l \leq m$. Picking $l = m$, we get

$$a_n^{m+1} b_0 = 0. \tag{4.46}$$

Since b_0 is a unit, this proves $a_n^{m+1} = 0$, i.e. the leading coefficient of $f(x)$ is a nilpotent.

Since $f(x)$ is a unit, and $a_n x^n$ is a nilpotent in $R[x]$, $f(x) - a_n x^n$ is a unit (earlier we have shown that if n is a nilpotent and u is a unit, then $u + n$ is a unit). Then repeating the same steps on $f(x) - a_n x^n$, we get that a_{n-1} is a nilpotent. Hence, repeating, all a_i's are nilpotents. ■

Exercise 4.19. Let R be a ring and let I and J be ideals in R. Which of the following are ideals of R

$$\begin{aligned} &I + J := \{u + v \mid u \in I,\, v \in J\}, \\ &I \cap J, \\ &I \cup J? \end{aligned} \tag{4.47}$$

Give a proof or a counterexample.

Solution. Yes, $I + J$ is an ideal of R. First, let's show that $I + J$ is an additive subgroup. Clearly, it contains 0, since $0 = 0 + 0$ and 0 is in both I and J. Given $u_1 + v_1, u_2 + v_2 \in I + J$, where $u_i \in I$ and $v_i \in J$,

$$(u_1 + v_1) - (u_2 + v_2) = \underbrace{(u_1 - u_2)}_{\in I} + \underbrace{(v_1 - v_2)}_{\in J} \in I + J. \tag{4.48}$$

So $I + J$ is an additive subgroup of R. Now take any $r \in R$, and $u + v \in I + J$ for $u \in I$, $v \in J$.

$$r(u + v) = \underbrace{ru}_{\in I} + \underbrace{rv}_{\in J} \in I + J; \tag{4.49}$$

$$(u + v)r = \underbrace{ur}_{\in I} + \underbrace{vr}_{\in J} \in I + J. \tag{4.50}$$

So $I + J$ is an ideal of R.

$I \cap J$ is an ideal. Indeed, intersections of subgroups is a subgroup, so $I \cap J$ is an additive subgroup. Furthermore, given any $r \in R$ and $a \in I \cap J$,

$$ra \in I \text{ and } ra \in J, \tag{4.51}$$

since I, J are ideals. Also

$$ar \in I \text{ and } ar \in J, \tag{4.52}$$

since I, J are ideals. Therefore, $ar, ra \in I \cap J$, so $I \cap J$ is an ideal.

$I \cup J$ is not, in general, an ideal of R. For instance, if

$$R = \mathbb{Z}, \quad I = 2\mathbb{Z}, \quad J = 3\mathbb{Z}, \tag{4.53}$$

$I \cup J$ contains $2, 3$, but it doesn't contain $2 + 3 = 5$. So it's not an ideal. ■

Exercise 4.20. Let R be a noncommutative ring and $a \in R$. Show by example that $aR := \{ar \mid r \in R\}$ is not always an ideal.

Solution. Let $R = \mathrm{Mat}_{2\times 2}(\mathbb{R})$, and take the matrix $a = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. Then

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} = \begin{bmatrix} x & y \\ 0 & 0 \end{bmatrix} \tag{4.54}$$

So the ideal aR is

$$aR = \left\{ \begin{bmatrix} x & y \\ 0 & 0 \end{bmatrix} \mid x, y \in \mathbb{R} \right\}. \tag{4.55}$$

We'll show that this is not an ideal by picking an element $b \in aR$ and $s \in R$ such that $sb \notin aR$. That will prove that aR is not an ideal. We pick $b = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$, $s = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$,

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \notin aR. \tag{4.56}$$

■

Exercise 4.21. Let R be a ring, $I \subseteq R$ an ideal.

(a) Show that $r(I) := \{r \in R \mid ra = 0 \text{ for all } a \in I\}$ is an ideal in R.

(b) Show that $[R : I] := \{x \in R \mid xr \in I \text{ for all } r \in R\}$ is an ideal in R.

Solution. (a) Clearly $0 \in r(I)$. To show that it's an additive subgroup, let $r, r' \in r(I)$,

$$\left(r - r'\right) a = ra - r'a = 0 - 0 = 0 \tag{4.57}$$

for all $a \in I$. Now, let $r \in r(I)$ and $s \in R$.

$$(rs)\, a = r\,(sa) = 0, \quad \text{forall } a \in I \tag{4.58}$$

since $sa \in I$ for I is an ideal. Furthermore,

$$(sr)\, a = s\,(ra) = s \cdot 0 = 0. \tag{4.59}$$

Therefore, $sr, rs \in r(I)$, proving $r(I)$ is an ideal.

(b) Clearly $0 \in [R : I]$, since $0 \cdot r = 0 \in I$ for all $r \in R$. To show that it's an additive subgroup, let $x, x' \in [R : I]$,

$$\left(x - x'\right) r = xr - x'r \in I, \quad \forall\, r \in R; \tag{4.60}$$

since $xr, xr' \in I$. Therefore, $x - x' \in [R : I]$. Now, let $s \in R$ and $x \in [R : I]$.

$$(sx)\, r = s\,(xr) \in I, \tag{4.61}$$

since $xr \in I$ and I is an ideal, so $s(xr) \in I$. Furthermore,

$$(xs)\, r = x\,(sr) \in I, \tag{4.62}$$

since $x \in [R : I]$. Therefore, $sx, xs \in [R : I]$, , proving $[R : I]$ is an ideal.

■

Exercise 4.22. Prove that a subring of an integral domain is an integral domain.

Solution. If there's no nonzero zero-divisor in the larger ring, there can't be any nonzero zero-divisor in the subring. ■

Exercise 4.23. Is there an integral domain with precisely 10 elements?

Solution. No. Finite integral domains are fields (Exercise 4.17), and hence, their cardinalities have to be prime powers. ■

Exercise 4.24. Let R be a ring such that $x^3 = x$ for every $x \in R$. Show that R is a commutative ring.

Solution. We proceed by a sequence of steps.

1. If $x^2 = 0$, then $x = 0$. Indeed, if $x^2 = 0$,

$$x = x^3 = x^2 \cdot x = 0 \cdot x = 0. \tag{4.63}$$

2. If $xy = 0$, then $yx = 0$.

$$yx = (yx)^3 = yxyxyx = y\,(xy)\,xyx = 0. \tag{4.64}$$

3. $xy = x^2y^2xy$ for all $x, y \in R$.

$$0 = yx - yx = yx - y^3x = y\left(x - y^2x\right) = \left(x - y^2x\right)y \tag{4.65}$$

by 2. Multiplying by x^2 to the left, we have

$$0 = x^2\left(x - y^2x\right)y = x^3y - x^2y^2xy = xy - x^2y^2xy. \tag{4.66}$$

Therefore, $xy = x^2y^2xy$.

4. $(xy - yx)yx = 0$ for all $x, y \in R$. By 3,

$$0 = x^2y^2xy - xy = x^2y^2xy - (xy)^3 = xxyyxy - xyxyxy = x\,(xy - yx)\,yxy. \tag{4.67}$$

By 2,

$$0 = (xy - yx)\,yxyx = (xy - yx)\,yxyxyx = (xy - yx)\,(yx)^3 = (xy - yx)\,yx. \tag{4.68}$$

5. $xy = yx$ for all $x, y \in R$. For this, we'll show that $(xy - yx)^2 = 0$, then we'll be done by 1.

$$\begin{aligned}(xy - yx)^2 &= (xy - yx)\,xy - (xy - yx)\,yx &(4.69)\\ &= -(yx - xy)\,xy - (xy - yx)\,yx &(4.70)\\ &= -0 - 0 = 0 &(4.71)\end{aligned}$$

by 4. Hence, we are done!

■

Exercise 4.25. Let R be a ring with 1 and let $\varphi : R \to S$ be a surjective homomorphism of rings. Show that $\varphi(1)$ is the unit element of S.

Solution. Since φ is surjective, any $s \in S$ is of the form $s = \varphi(r)$ for some $r \in R$. Then

$$\begin{aligned}\varphi(1) \cdot s = \varphi(1)\varphi(r) = \varphi(1 \cdot r) = \varphi(r) = s, &\tag{4.72}\\ s \cdot \varphi(1) = \varphi(r)\varphi(1) = \varphi(r \cdot 1) = \varphi(r) = s. &\tag{4.73}\end{aligned}$$

Hence, $\varphi(1)$ is the unit element of S. ■

Exercise 4.26. Let k be a field. Find all automorphisms φ of $k[x]$ with the property that $\varphi(a) = a$ for all $a \in k$.

Solution. Every homomorphism $f : k[x] \to k[x]$ is uniquely determined by where it sends x. Since f is an automorphism, let $g = f^{-1}$. Suppose

$$f(x) = P(x) \in k[x] \text{ and } g(x) = Q(x). \tag{4.74}$$

Then since $f \circ g = \mathrm{id}$,

$$x = f(g(x)) = P(Q(x)). \tag{4.75}$$

The degree of $P(Q(x))$ is $\deg P \cdot \deg Q$, which has to be equal to $\deg x = 1$. Hence, both $\deg P = \deg Q = 1$. Therefore,

$$P(x) = ax + b \tag{4.76}$$

for $a, b \in k$ with $a \neq 0$. Then

$$f(x) = ax + b \tag{4.77}$$

is, indeed, invertible since its inverse is given by

$$g(x) = a^{-1}x - a^{-1}b. \tag{4.78}$$

So all such automorphisms $k[x] \to k[x]$ are of the form

$$\sum_{i=1}^{n} a_i x^i \mapsto \sum_{i=1}^{n} a_i (ax+b)^i \text{ for } a \in k^\times, b \in k. \tag{4.79}$$

■

Exercise 4.27. Prove that every ideal in $\mathbb{Z}[i]$ contains a nonzero integer.

Solution. Let I be a nonzero ideal. Then pick nonzero $a + ib \in I$. Since I is an ideal, it contains any element of the form xr for $x \in I$ and $r \in R = \mathbb{Z}[i]$. Choosing $x = a + ib$ and $r = a - ib$, we have

$$(a+ib)(a-ib) = a^2 + b^2 \in I. \tag{4.80}$$

$a^2 + b^2$ is an integer, and its nonzero since $a^2 + b^2 = 0$ if and only if $a = b = 0$, which isn't the case as we've picked $a + ib \neq 0$. Therefore, I contains a nonzero integer, namely $a^2 + b^2$. ■

Exercise 4.28. Show that the kernel of the homomorphism $\varphi : \mathbb{C}[x,y] \to \mathbb{C}[t] : x \mapsto t^2,\ y \mapsto t^3$ is the principal ideal generated by $y^2 - x^3$.

Solution. We'll have to show that $\operatorname{Ker}\varphi = \left(y^2 - x^3\right)$. The inclusion $\supseteq$ is clear, since

$$\varphi\left(y^2 - x^3\right) = \varphi(y)^2 - \varphi(x)^3 = \left(t^3\right)^2 - \left(t^2\right)^3 = 0, \tag{4.81}$$

so we have $\left(y^2 - x^3\right) \subseteq \operatorname{Ker}\varphi$.

Let $g(x,y) \in \operatorname{Ker}\varphi$. We consider $\mathbb{C}[x,y] = \mathbb{C}[x][y]$, i.e. we consider g to be a polynomial in y, and we divide by $y^2 - x^3$ (which is a monic polynomial in $\mathbb{C}[x][y]$).

$$g(x,y) = \left(y^2 - x^3\right) q(x,y) + r(x,y), \tag{4.82}$$

with $\deg_y r(x,y) < \deg_y \left(y^2 - x^3\right) = 2$, so $\deg_y r(x,y) \leq 1$. We write

$$r(x,y) = r_0(x) + r_1(x)y, \tag{4.83}$$

where $r_0(x), r_1(x) \in \mathbb{C}[x]$. Now, since $g(x,y), y^2 - x^3 \in \operatorname{Ker}\varphi$, we have $r(x,y) \in \operatorname{Ker}\varphi$. Now,

$$0 = \varphi(r(x,y)) = r_0\left(t^2\right) + r_1(t^2)t^3. \tag{4.84}$$

The terms of $r_0\left(t^2\right)$ are all of the form t^{even} and the terms of $r_1(t^2)t^3$ are all of the form t^{odd}. Therefore, their sum can be 0 if and only if $r_0 = r_1 = 0$, in which case $r(x,y) = 0$. In other words, $g \in \left(y^2 - x^3\right)$, proving that $\operatorname{Ker}\varphi \subseteq \left(y^2 - x^3\right)$. ■

Exercise 4.29. Let R be a ring and I an ideal in R. Let $\sqrt{I} := \{x \in R \mid x^n \in I \text{ for some } n > 0\}$. Show

(a) $\sqrt{I}$ is an ideal in R which contains I. $\sqrt{I}$ is usually called the radical of I.

(b) $\sqrt{\sqrt{I}} = \sqrt{I}$.

Solution. Here, we need the hypothesis that R is commutative. Without that $\sqrt{I}$ fails to be an ideal, since if $I = (0)$, the $\sqrt{I}$ is the set of all nilpotent elements, which need not be closed under addition if R is non-commutative (Exercise 4.6). So we'll assume R is commutative.

(a) It's clear that $I \subseteq \sqrt{I}$. Clearly, $0 \in \sqrt{I}$. To show that it's an additive subgroup, take $x, y \in \sqrt{I}$. Then $x^n, y^m \in I$ for some $m, n > 0$. Then

$$(x-y)^{m+n} = \sum_{i=0}^{m+n} \pm \binom{m+n}{i} x^i y^{m+n-i}. \tag{4.85}$$

If $i < n$, $m+n-i > m$, so $y^{m+n-i} \in I$. Since I is an ideal, $x^i y^{m+n-i} \in I$. On the other hand, if $i \geq n$, $x^i \in I$, so that $x^i y^{m+n-i} \in I$. Therefore, all summands are in I, so $(x-y)^{m+n} \in I$, proving that $x - y \in \sqrt{I}$.

Now, for $x \in \sqrt{I}$ and $r \in R$, $x^n \in I$ for some $n > 0$. Then

$$(rx)^n = r^n x^n \in I, \tag{4.86}$$

so $rx \in \sqrt{I}$. Hence, $\sqrt{I}$ is an ideal of R.

(b) By (a), $\sqrt{I} \subseteq \sqrt{\sqrt{I}}$. Now, let $x \in \sqrt{\sqrt{I}}$.

$$\begin{aligned} x \in\in \sqrt{\sqrt{I}} &\implies x^n \in \sqrt{I} \text{ for some } n > 0 \\ &\implies (x^n)^m = x^{mn} \in I \text{ for some } m, n > 0 \\ &\implies x \in \sqrt{I}. \end{aligned}$$

So $\sqrt{\sqrt{I}} \subseteq \sqrt{I}$.

■

Exercise 4.30. Let R be a ring and let A, B be ideals in R such that $A \cap B = \{0\}$. Show that for all $a \in A$, $b \in B$, $ab = 0$.

Solution. Since A is an ideal and $a \in A$, $ab \in A$. Similarly, since B is an ideal and $b \in B$, $ab \in B$. Therefore, $ab \in A \cap B = \{0\}$, so $ab = 0$. ■

Exercise 4.31. Let R be a ring (commutative with 1) and let S be a subring. The *conductor* $C(S)$ of S in R is the set of all $\alpha \in R$ such that $\alpha R \subseteq S$.

(a) Prove that $C(S)$ is an ideal of R and also an ideal of S.

(b) Prove that $C(S)$ is the largest ideal of S which is also an ideal of R.

(c) Determine $C(S)$ in the following cases:

(i) $R = \mathbb{C}[x]$, $S = \mathbb{C}\left[x^2, x^3\right]$.

(ii) $R = \mathbb{Z}[\omega]$, $\omega = \dfrac{-1+\sqrt{-3}}{2}$, $S = \mathbb{Z}\left[\sqrt{-3}\right]$.

Solution. (a) Clearly $0 \in C(S)$, since $0R = 0 \subseteq S$. To show that it's an additive subgroup, let $\alpha_1, \alpha_2 \in C(S)$. Then for any $r \in R$,

$$(\alpha_1 - \alpha_2) r = \underbrace{\alpha_1 r}_{\in S} - \underbrace{\alpha_2 r}_{\in S} \in S. \tag{4.87}$$

For $\alpha \in C(S)$ and $r \in R$,

$$(r\alpha) R = \alpha (rR) \subseteq \alpha R \subseteq S, \tag{4.88}$$

so $r\alpha \in C(S)$. Therefore, $C(S)$ is an ideal of R. It's also an ideal of S, since $r\alpha \in C(S)$ for every $r \in R$ implies $r\alpha \in C(S)$ for every $r \in S$ as $S \leq R$.

(b) Let $I \trianglelefteq R$ and $I \subseteq S$. We'll prove that $I \subseteq C(S)$. For $a \in I$, $aR \subseteq I$ since $I \trianglelefteq R$. But $I \subseteq S$. So we have

$$aR \subseteq I \subseteq S \implies a \in C(S), \tag{4.89}$$

proving $I \subseteq C(S)$.

(c) By (b), $C(S)$ is the largest ideal of R contained in S.

(i) Here, $S = \mathbb{C}\left[x^2, x^3\right]$ contains all the polynomials whose x^1 term is missing, i.e. the x^1 term has coefficient 0. $C(S)$ is the largest ideal of R contained in S. So let $f(x) \in C(S) \trianglelefteq R = \mathbb{C}[x]$. Since $C(S)$ is an ideal, $xf(x) \in C(S)$.

But $C(S)$ is contained in S, so $xf(x)$ has x^1's coefficient 0. In other words, $f(x)$ has no constant term. But $f(x) \in C(S) \subseteq S$ already implies $f(x)$ has no x^1 term. Therefore,

$$f(x) = a_2 x^2 + a_3 x^3 + \cdots + a_n x^n. \tag{4.90}$$

In other words, $C(S) = x^2\mathbb{C}[x]$.

(ii) Note that $\sqrt{3} = 2\omega + 1$, so

$$S = \mathbb{Z}\left[\sqrt{3}\right] = \mathbb{Z}[2\omega + 1] = \mathbb{Z}[2\omega]. \tag{4.91}$$

$C(S)$ is the largest ideal of $\mathbb{Z}[\omega]$ that is contained in $\mathbb{Z}[2\omega]$. Let $a + b\omega \in C(S)$. Since $C(S) \subseteq \mathbb{Z}[2\omega]$, we must have that b is even.

Since $C(S) \trianglelefteq \mathbb{Z}[\omega]$, $\omega(a + b\omega) \in C(S) \subseteq \mathbb{Z}[2\omega]$.

$$\omega(a + b\omega) = a\omega + b\omega^2 = a\omega + b(-1 - \omega) = -b + (a - b)\omega \tag{4.92}$$

So $a - b$ is even. But we have already proved b is even. So a is even. Therefore, $C(S)$ is $2\mathbb{Z}[\omega]$.

■

Exercise 4.32. Let I, J be ideals of a ring, such that $I + J = R$. Prove that $IJ = I \cap J$.

Solution. R needs to be commutative and have 1, otherwise this fails. Let

$$R = 2\mathbb{Z}, \quad I = 4\mathbb{Z}, \quad J = 6\mathbb{Z}. \tag{4.93}$$

Then $I + J = R$. But $I \cap J = 12\mathbb{Z}$ and $IJ = 24\mathbb{Z}$. For noncommutative R, consider

$$R = k\langle x, y\rangle, \quad I = \langle x\rangle, \quad J = \langle x - 1\rangle. \tag{4.94}$$

Then indeed, $I + J = R$ since it contains 1. But the element $(x-1)yx \neq xy(x-1)$ is in $I \cap J$ but it's not in IJ.

Now, suppose R is a commutative ring with 1. Let $ij \in IJ$, for $i \in I$ and $j \in J$. Then $ij \in I$ since I is an ideal, and $ij \in J$ since J is an ideal. Therefore, $IJ \subseteq I \cap J$.

Now, let $x \in I \cap J$. Since $I + J = R$, there exists $i \in I$ and $j \in J$ such that

$$i + j = 1. \tag{4.95}$$

Multiplying on the left by x,

$$xi + xj = x. \tag{4.96}$$

$x \in J$ implies $xi \in JI = IJ$. $x \in I$ implies $xj \in IJ$. Therefore, $x = xi + xj$ is in IJ, proving $I \cap J \subseteq IJ$. ■

Lothar says,

> From now on until the end of the lectures a ring is a commutative ring with 1 and a homomorphism $\varphi : R \to S$ of rings satisfies $\varphi(1) = 1$.

Exercise 4.33. Let R be a commutative ring with 1. An element $r \in R$ is called *nilpotent* if there exists an $n \in \mathbb{Z}_{>0}$ with $r^n = 0$. Let

$$N(R) := \{r \in R \mid r \text{ is nilpotent}\}. \tag{4.97}$$

(a) Determine $N(\mathbb{Z}/4\mathbb{Z})$.

(b) Show that $N(R)$ is an ideal in R.

(c) Show that $N(R) \subseteq P$ for every prime ideal $P \subseteq R$.

(d) Show that 0 is the only nilpotent element of $R/N(R)$.

Solution. (a) Nilpotents are zero-divisors, in particular. The only zero divisors in $\mathbb{Z}_4$ are $0, 2$. Both of them are nilpotents since $0^1 = 0, 2^2 = 0$. Therefore,

$$N(\mathbb{Z}_4) = \{0, 2\}. \tag{4.98}$$

(b) Since $N(R) = \sqrt{(0)}$, it's clearly an ideal, by Exercise 4.29.

(c) Let $\mathfrak{p}$ be a prime ideal, and let $x \in N(R)$ with $x^n = 0$. Then $x^n = 0 \in \mathfrak{p}$. Let

$$m = \min\left\{i > 0 \mid x^i \in \mathfrak{p}\right\}. \tag{4.99}$$

If $m = 1$, we are done! So assume $m \neq 1$. Then

$$x^m = x \cdot x^{m-1} \in \mathfrak{p}, \tag{4.100}$$

then either $x \in \mathfrak{p}$ or $x^{m-1} \in \mathfrak{p}$. We already assumed $m \neq 1$, so $x \notin \mathfrak{p}$. Therefore, $x^{m-1} \in \mathfrak{p}$, which contradicts the minimality of $\mathfrak{p}$. Hence, $x \in \mathfrak{p}$, i.e. $N(R) \subseteq \mathfrak{p}$.

(d) Let $r + N(R) \in R/N(R)$ be a nilpotent element. Then

$$0 + N(R) = (r + N(R))^n = r^n + N(R) \text{ for some } n > 0. \tag{4.101}$$

This means $r^n \in N(R)$ is a nilpotent, i.e. for some $m > 0$, we have

$$0 = (r^n)^m = r^{mn}. \tag{4.102}$$

So r is a nilpotent, i.e. $r + N(R) = 0 + N(R)$. Hence, 0 is the only nilpotent in $R/N(R)$.

■

Exercise 4.34. Let k be a field and $f \in k[x]$. Let $R := k[x]/\langle f \rangle$. Show that $N(R) = 0$ (see previous exercise) if and only if f is not divisible by the square of a polynomial (aka, square-free).

Solution. Since k is a field, any polynomial $f \in k[x]$ can be written as a product of powers of irreducibles:

$$f = p_1^{e_1} \cdots p_k^{e_k}, \tag{4.103}$$

where p_i's are irreducibles.

Suppose f is square-free. Then $e_i = 1$ for each i. If $g + (f) \in N\left(\frac{k[x]}{(f)}\right)$, then for some $n > 0$, $g^n \in (f)$. Then

$$f = p_1 \cdots p_k \mid g^n. \tag{4.104}$$

Therefore, each $p_i \mid g^n$. But since p_i's are irreducibles (and thus primes, since $k[x]$ is a UFD), $p_i \mid g^n$ if and only if $p_i \mid g$. Therefore, $f \mid g$, i.e.

$$g + (f) = 0 + (f), \quad \text{i.e. } N\left(\frac{k[x]}{(f)}\right) = 0. \tag{4.105}$$

Conversely, suppose $N\left(\frac{k[x]}{(f)}\right) = 0$. Write $f = p_1^{e_1} \cdots p_k^{e_k}$ as a product of irreducibles. Assume for the sake of contradiction that at least one of the e_i's is greater than 1.

Take $g = p_1 \cdots p_k$. Then $g + (f) \neq 0 + (f)$ since $f \nmid g$. Now, let $n = \max\{e_1, \ldots, e_k\} \geq 2$. Then

$$g^n = p_1^n \cdots p_k^n, \tag{4.106}$$

which is divisible by $p_1^{e_1} \cdots p_k^{e_k}$, i.e.

$$[g + (f)]^n = 0 + (f), \tag{4.107}$$

proving that $g + (f) \in N(\frac{k[x]}{(f)})$. But we started with $N\left(\frac{k[x]}{(f)}\right) = 0$. Contradiction! Hence, f must be square-free. ■

Exercise 4.35. Determine the maximal ideals of $\mathbb{R}[x] / \left(x^2\right)$ and $\mathbb{R}[x] / \left(x^2 + x + 1\right)$.

Solution. We'll use Exercise 4.37. An ideal $\mathfrak{m}$ of $\mathbb{R}[x]/(f(x))$ is maximal if and only if $\pi^{-1}(\mathfrak{m})$ (that contains $f(x)$) is maximal in $\mathbb{R}[x]$. A maximal ideal is prime, so it contains all the nilpotents (Exercise 4.33). Therefore,

$$\left[x + \left(x^2\right)\right]^2 = x^2 + \left(x^2\right) = 0 + \left(x^2\right), \tag{4.108}$$

so $\overline{x} = x + \left(x^2\right)$ is a nilpotent in $\mathbb{R}[x]/\left(x^2\right)$. So $\overline{x} \in \mathfrak{m}$, i.e. $\pi^{-1}(\mathfrak{m})$ contains x and x^2. We claim $\pi^{-1}(\mathfrak{m}) = (x)$, because if a nonzero real number $c \in \pi^{-1}(\mathfrak{m})$, then it's the whole $\mathbb{R}[x]$. Therefore, $\pi^{-1}(\mathfrak{m}) = (x)$. So the only maximal ideal of $\mathbb{R}[x]/(x^2)$ is $(\overline{x})$.

We claim that $\mathbb{R}[x] / \left(x^2 + x + 1\right)$ is a field. For $ax + b \in \left(x^2 + x + 1\right)$,

$$(ax + b)(cx + d) = acx^2 + (ad + bc)x + bd = ac(-x - 1) + (ad + bc)x + bd \tag{4.109}$$

$$= (ad + bc - ac)x + (bd - ac). \tag{4.110}$$

(The equality is $\mathbb{R}[x] / \left(x^2 + x + 1\right)$.) Then we can solve for c, d to make this quantity 1:

$$c = \frac{a}{a^2 - ab + b^2}, \quad d = \frac{b - a}{a^2 - ab + b^2}. \tag{4.111}$$

(Of course $a^2 - ab + b^2 \neq 0$ for $a \neq 0$, $b \neq 0$.) Therefore,

$$\frac{a}{a^2 - ab + b^2}x + \frac{b - a}{a^2 - ab + b^2} = (ax + b)^{-1}. \tag{4.112}$$

So $\mathbb{R}[x] / \left(x^2 + x + 1\right)$ is a field, and hence its unique maximal ideal is (0). ■

Exercise 4.36. Prove that the ideal $I = \left\langle x + y^2, y + x^2 + 2xy^2 + y^4 \right\rangle$ in $\mathbb{C}[x, y]$ is a maximal ideal.

Solution. We claim that $I = (x, y)$. Indeed, $x + y^2 \in (x, y)$, and $y + x^2 + 2xy^2 + y^4 \in (x, y)$. Therefore, $I \subseteq (x, y)$.

Now,

$$y + x^2 + 2xy^2 + y^4 = y + \left(x + y^2 - y^2\right)^2 + 2\left(x + y^2 - y^2\right) y^2 + y^4 \tag{4.113}$$

$$= y + \left(x + y^2\right) P - y^4 + (x + y^2)Q - 2y^4 + y^4 \tag{4.114}$$

$$= y + \left(x + y^2\right) P + (x + y^2)Q \tag{4.115}$$

for some $P, Q \in \mathbb{C}[x, y]$. Therefore, $y \in I$. Since $y \in I$, $y^2 \in I$, and hence from $x + y^2 \in I$ we have $x \in I$. As a result, $(x, y) \in I$.

So we have proved $I = (x, y)$. This is a maximal ideal because

$$R/I = \frac{\mathbb{C}\,[x, y]}{(x, y)} \cong \mathbb{C}, \tag{4.116}$$

is a field, so I is a maximal ideal. ■

Exercise 4.37. Let R be a ring, and let I be an ideal of R. Let $\overline{R} := R/I$. For an ideal $M \subseteq R$ containing I let $\overline{M} = M/I$. Show that $\overline{M}$ is maximal in $\overline{R}$ if and only if M is maximal in R.

Solution. Preimage of ideals under a homomorphism is an ideal, and image of ideals under a surjective homomorphism is an ideal. Therefore, if $\overline{M}$ is maximal and M is not, there exists an ideal

$$M \subsetneq L \subsetneq R. \tag{4.117}$$

Taking the image under π, we have

$$M/I \subsetneq L/I \subsetneq R/I, \tag{4.118}$$

contradicting the maximality of $\overline{M} = M/I$. Similarly, if M is maximal but $\overline{M}$ is not, then there exists an ideal

$$\overline{M} \subsetneq \overline{L} \subsetneq \overline{R}. \tag{4.119}$$

Taking the preimage under π, we get

$$M \subsetneq \pi^{-1}\left(\overline{L}\right) \subsetneq R, \tag{4.120}$$

contradicting the maximality of M. ■

Exercise 4.38. Give an example of a ring in which some prime ideal is not a maximal ideal.

Solution. Consider the ring $\mathbb{Z}[x]$. The ideal (x) is prime, since $\mathbb{Z}[x]/(x) \cong \mathbb{Z}$ is an ID. But (x) is not maximal since $\mathbb{Z}[x]/(x) \cong \mathbb{Z}$ is not a field. ■

Exercise 4.39. Let R be a commutative ring with 1 and let I be an ideal that is contained in a finite union $\mathfrak{p}_1 \cup \cdots \cup \mathfrak{p}_n$ of prime ideals of R. Show that $I \subseteq \mathfrak{p}_i$ for some i.

Solution. We shall prove it by induction on n. The base case $n = 1$ is clearly true. Suppose the statement is true for $n - 1$. For each i, if I is contained in $\bigcup_{j \neq i} \mathfrak{p}_j$, then we're done by the inductive hypothesis. So suppose that's not the case, so we can pick $x_i \in I$ such that $x_i \notin \bigcup_{j \neq i} \mathfrak{p}_j$. Then it has to be

$$x_i \in \mathfrak{p}_i. \tag{4.121}$$

Now, take the element

$$y = x_1 + x_2 x_3 \cdots x_n \in I. \tag{4.122}$$

It's in I because each $x_i \in I$. Therefore, $y \in \mathfrak{p}_k$ for some k.

If $y \in \mathfrak{p}_1$, since $x_1 \in \mathfrak{p}_1$, $x_2 x_3 \cdots x_n \in \mathfrak{p}_1$. But since $\mathfrak{p}_1$ is prime, that would imply some $x_i \in \mathfrak{p}_1$. But by construction, $x_i \notin \bigcup_{j \neq i} \mathfrak{p}_j$, so in particular, $x_i \notin \mathfrak{p}_1$.

If $y \in \mathfrak{p}_k$ for $k > 1$, then since $x_k \in \mathfrak{p}_k$, $x_2 x_3 \cdots x_n \in \mathfrak{p}_k$. So we have $x_1 \in \mathfrak{p}_k$. But by construction, $x_1 \notin \bigcup_{j \neq 1} \mathfrak{p}_j$, so in particular, $x_1 \notin \mathfrak{p}_k$.

Therefore, we arrive at a contradiction in all cases! ■

Exercise 4.40. Let $f : R \to S$ be a surjective ring homomorphism with kernel K. Show:

(a) If P is a prime ideal of R containing K, then $f(P)$ is a prime ideal of S.

(b) If Q is a prime ideal of S, then $f^{-1}(Q)$ is a prime ideal of R containing K.

Solution. (a) Since f is a surjective homomorphism, $f(P)$ is an ideal of S.

If $f(P) = S$, that means there exists some $r \in P$ such that $f(r) = 1$. Then

$$1 - r \in \operatorname{Ker} f = K \subseteq P. \tag{4.123}$$

$1 - r \in P$ and $r \in P$ together gives us $1 \in P$, which means $P = R$, which is a contradiction since a prime ideal is never the whole ring. Therefore, $f(P) \neq S$.

Now, let $a, b \in S$ with $ab \in f(P)$. Since f is surjective, $a = f(x)$ and $b = f(y)$ for some $x, y \in R$. Then

$$f(xy) = f(x) f(y) = ab \in f(P). \tag{4.124}$$

This means $xy \in P + K = P$. Therefore, $x \in P$ or $y \in P$. WLOG $x \in P$, so that $f(x) = a \in f(P)$, proving that $f(P)$ is a prime ideal.

(b) $f^{-1}(Q)$ is an ideal since f is a homomorphism. It contains K since Q contains 0, so $f^{-1}(Q)$ contains $f^{-1}(0) = K$.

It's easy to see that $f^{-1}(Q) \neq R$. Since Q is prime, it doesn't contain 1, so $f^{-1}(Q)$ doesn't contain 1 either, so $f^{-1}(Q) \neq R$.

Now, suppose $a, b \in R$ with $ab \in f^{-1}(Q)$. Then $f(ab) \in Q$. But $f(ab) = f(a)f(b)$. Since Q is prime, $f(a) \in Q$ or $f(b) \in Q$. WLOG $f(a) \in Q$, so that $a \in f^{-1}(Q)$. Therefore, Q is prime.

■

Exercise 4.41. Let R be a ring with 1, M an ideal in R. Suppose every element of $R \setminus M$ is a unit in R. Show that M is the unique maximal ideal of R.

Solution. Let $M' \neq M$ be a maximal ideal. Then $M' \not\subseteq M$, so it contains some element not in M, say u. By hypothesis, u is a unit. So

$$1 = u \cdot u^{-1} \in M'. \tag{4.125}$$

But that means for every $r \in R$,

$$r = r \cdot 1 \in M', \tag{4.126}$$

proving $M' = R$. ■

Exercise 4.42. Show: The ring $2\mathbb{Z}$ of even integers contains a maximal ideal M, such that $2\mathbb{Z}/M$ is not a field.

Solution. In $R = 2\mathbb{Z}$, $\mathfrak{m} = 4\mathbb{Z}$ is a maximal ideal. But $R/\mathfrak{m} = \left\{\overline{0}, \overline{2}\right\}$, which is not a field since it has no multiplicative identity. ■

Exercise 4.43. Find the greatest common divisors of the following polynomials in $\mathbb{Q}[x]$.

(a) $x^3 - 6x^2 + x + 4$ and $x^5 - 6x + 1$.

(b) $x^2 + 1$ and $x^6 + x^3 + x + 1$.

Solution. (a) Let's first factorize $f(x) := x^3 - 6x^2 + x + 4$. We can see that 1 is a root of this polynomial. In fact,

$$x^3 - 6x^2 + x + 4 = (x - 1)\left(x^2 - 5x - 4\right). \tag{4.127}$$

$x^2 - 5x - 4$ is irreducible since it has no rational root. Now, 1 is not a root of $g(x) := x^5 - 6x + 1$, so $x - 1$ doesn't divide g. Furthermore, $x^2 - 5x - 4$ also doesn't divide g, as can be checked easily.

Therefore, $\gcd(f, g) = 1$.

(b) Note that

$$x^6 + x^3 + x + 1 = \left(x^2 + 1\right)\left(x^4 - x^2 + x + 1\right), \tag{4.128}$$

so their gcd is $x^2 + 1$.

■

Exercise 4.44. Let k be a field and let $a_0, \ldots, a_n \in k$ be distinct elements and let $b_0, \ldots, b_n \in k$. Show: There is at most one polynomial $f \in k[x]$ of degree n with $f(a_i) = b_i$ for $i = 0, \ldots, n$.

Solution. Suppose there are at least two such polynomaials f, f'. Then $\deg(f - f') \leq n$. But

$$(f - f')(a_i) = f(a_i) - f'(a_i) = b_i - b_i = 0. \tag{4.129}$$

Therefore, $f - f'$ has $n + 1$ roots, while its degree is at most n. This can happen only if it's the zero polynomial, i.e. $f = f'$. ■

Exercise 4.45. Let $a_0, \ldots, a_{n-1} \in \mathbb{Z}$ and let

$$f = x^n + \sum_{i=0}^{n-1} a_i x^i \in \mathbb{Q}[x] \tag{4.130}$$

(note that the leading coefficient is 1). Let $b \in \mathbb{Q}$ be a zero of f. Show that $b \in \mathbb{Z}$.

Solution. Suppose $b = \frac{p}{q}$ for $p, q \in \mathbb{Z}$ and $q \neq 0$, and $\gcd(p, q) = 1$. Then $f(b) = 0$, so we have

$$\sum_{i=0}^{n} a_i \left(\frac{p}{q}\right)^i = 0, \tag{4.131}$$

where $a_n = 1$. Multiplying by q^n, we have

$$\sum_{i=0}^{n} a_i p^i q^{n-i} = 0. \tag{4.132}$$

Except for the $i = n$ term, all the terms are divisible by q, and the RHS is obviously divisible by q. Hence,

$$q \mid a_n p^n = p^n. \tag{4.133}$$

Therefore, $q = 1$, since p and q has no common factor other than 1. So $b = p \in \mathbb{Z}$. ■

Exercise 4.46. Let $f, g \in k[x]$ be polynomials with coefficients in a field k. Assume the both f and g factor into linear factors i.e. $f = (x + a_1) \cdot \cdots \cdot (x + a_n)$, $g = (x + b_1) \cdot \cdots \cdot (x + b_m)$. Then the greatest common divisor is the product of the common linear factors.

Solution. It's clear that the product of the common linear factors is a common divisor of f and g. Another common divisor would be the product of a subcollection of these common linear factors, so that clearly divides the product of all common linear factors. ■

Exercise 4.47. Prove or disprove the following statement: The ring $\mathbb{Z}[x]$ is a principal ideal domain.

Solution. This is false. Because the ideal $(2, x) \trianglelefteq \mathbb{Z}[x]$ is not a principal ideal. If it were, $(2, x) = (f)$, then f must divide 2. But that would mean f is a constant polynomial ± 1 or ± 2. In the first case, $(f) = \mathbb{Z}[x]$, and in the second case, $2 \mid x$, which is not possible. ■

Exercise 4.48. If $a + bi \in \mathbb{Z}[i]$ is not a unit, show that $a^2 + b^2 > 1$.

Solution. If $a^2 + b^2 = 0$, that would just imply $a = b = 0$, which is 0. If $a^2 + b^2 = 1$,

$$1 = a^2 + b^2 = (a + ib)(a - ib), \tag{4.134}$$

proving $a + ib$ is a unit. Therefore, ■

Exercise 4.49. Show that $\mathbb{Z}\left[\sqrt{2}\right]$ is an Euclidean domain.

Solution. Let $N\left(a + b\sqrt{2}\right) = \left|a^2 - 2b^2\right|$ be the norm function. We claim that $\mathbb{Z}[\sqrt{2}]$ of $\mathbb{R}$ is an ED under this function. N is multiplicative, so $N(xy) = N(x)N(y) \geq N(x)$ for $x, y \neq 0$. So now we need to verify the second property of Euclidean function.

Let $\alpha = a + b\sqrt{2}$, and $\beta = c + d\sqrt{2}$. Then $\alpha/\beta \in \mathbb{Q}[\sqrt{2}]$ is given by

$$\frac{\alpha}{\beta} = \frac{ac - 2bd}{c^2 - 2d^2} + \frac{bc - ad}{c^2 - 2d^2}\sqrt{2} = q_1 + q_2\sqrt{2}. \tag{4.135}$$

Let m and n be the integers closest to q_1 and q_2, respectively. Therefore, we have $|m - q_1| \leq \frac{1}{2}$ and $|n - q_2| \leq \frac{1}{2}$.

We take $\gamma = m + n\sqrt{2} \in \mathbb{Z}[\sqrt{2}]$ and

$$\delta = \beta\left((q_1 - m) + \sqrt{2}(q_2 - n)\right) = \beta\left(\frac{\alpha}{\beta} - \gamma\right) = \alpha - \gamma\beta \in \mathbb{Z}[\sqrt{2}]$$

Then we have $\alpha = \beta\gamma + \delta$. Now we need to show that if $N(\delta)$ is nonzero, then $N(\delta) < N(\beta)$.

N is multiplicative not only in $\mathbb{Z}[\sqrt{2}]$, but also in $\mathbb{Q}[\sqrt{2}]$.

$$N(\delta) = N(\beta)N\left((q_1 - m) + \sqrt{2}(q_2 - n)\right) = N(\beta)\left|(q_1 - m)^2 - 2(q_2 - n)^2\right| \tag{4.136}$$

$$\leq N(\beta)\left(|q_1 - m|^2 + 2|q_2 - n|^2\right) \leq N(\beta)\left(\frac{1}{4} + 2 \times \frac{1}{4}\right) \tag{4.137}$$

$$\leq \frac{3}{4}N(\beta) < N(\beta) \tag{4.138}$$

Therefore, N is an euclidean function, and that's why $\mathbb{Z}[\sqrt{2}]$ is an ED. ■

Exercise 4.50. Give an example that division with rest need not be unique in a Euclidean domain.

Solution. $\mathbb{Z}$ is an ED with the norm function $N(a) = |a|$. Then division with rest (remainder) is not unique in $\mathbb{Z}$. For instance, if we divide 7 by 3,

$$7 = 3 \cdot 2 + 1, \tag{4.139}$$

here the remainder is 1, and $N(1) = 1 < 3 = N(3)$. But also,

$$7 = 3 \cdot 3 + (-2), \tag{4.140}$$

here the remainder is -2, and $N(-2) = 2 < 3 = N(3)$. So division with remainder is not unique. ■

Exercise 4.51. Let $m, n \in \mathbb{Z}$. Show that their greatest common divisor in $\mathbb{Z}$ is the same as their greatest common divisor in $\mathbb{Z}[i]$.

Solution. Let $d = \gcd_{\mathbb{Z}}(m, n)$ and $d' = \gcd_{\mathbb{Z}[i]}(m, n)$ (which is not unique, it's unique up to multiplication by a unit, which are $\pm 1, \pm i$ in $\mathbb{Z}[i]$).

Since $\mathbb{Z} \subseteq \mathbb{Z}[i]$, d is a common divisor of m and n in $\mathbb{Z}[i]$. Therefore, $d \mid d'$ in $\mathbb{Z}[i]$.

Since $d = \gcd_{\mathbb{Z}}(m, n)$, by Bezout's identity, there exists integers x, y such that

$$d = mx + ny. \tag{4.141}$$

Since $d' \mid m$ and $d' \mid n$ (in $\mathbb{Z}[i]$), we have that $d' \mid mx + ny = d$ (in $\mathbb{Z}[i]$). In general, any common divisor of m and n in $\mathbb{Z}[i]$ also divides d. Therefore, $d = d'$ up to multiplication by a unit. ■

Exercise 4.52. Prove that

(a) $x^2 + x + 1$ is irreducible in $\mathbb{F}_2[x]$.

(b) $x^2 + 1$ is irreducible in $\mathbb{F}_7[x]$.

Solution. (a) If it wasn't irreducible, then it would split into degree-1 terms, i.e. it would have roots. But $f(0) = f(1) = 1 \neq 0$, so $x^2 + x + 1$ is irreducible in $\mathbb{F}_2[x]$.

(b) If it wasn't irreducible, then it would split into degree-1 terms, i.e. it would have roots. But since 7 is a prime of the form $4k - 1$, there doesn't exist any x with $x^2 + 1 = 0$.

■

Exercise 4.53. Let R be a commutative ring with no nonzero nilpotent elements (i.e. $a^n = 0$ implies $a = 0$). If $f = \sum_{i=0}^{n} a_i x^i \in R[x]$ is a zero divisor, show that there is an element $b \neq 0$ in R with $ba_0 = ba_1 = \cdots = ba_n = 0$.

Solution. In fact, we don't need the condition that R has no nonzero nilpotent. Since f is a zero divisor, let $g \neq 0$ be a polynomial of **minimal degree** such that $fg = 0$. We'll show that $\deg g = 0$.

$$f = \sum_{i=0}^{n} a_i x^i, \quad g = \sum_{i=0}^{m} b_i x^i. \tag{4.142}$$

Then

$$fg = \sum_{k=0}^{m+n} \left(\sum_{i+j=k} a_i b_j \right) x^k = 0. \tag{4.143}$$

Looking at the coefficient of x^{m+n}, we see that $a_n b_m = 0$. Therefore, $\deg(a_n g) < m$. But

$$f \cdot (a_n g) = a_n \cdot (fg) = 0, \tag{4.144}$$

and $a_n g$ is a polynomial of degree less than $\deg g$ with $f \cdot (a_n g) = 0$. This contradicts the minimality of $\deg g$. Hence, $a_n g = 0$.

We'll prove that $a_{n-l} g = 0$ for every $0 \leq l \leq n$. We'll prove it by induction on l. The base case $l = 0$ has already been done. Now suppose

$$a_n g = a_{n-1} g = \cdots = a_{n-l+1} g = 0. \tag{4.145}$$

As a result,

$$a_n b_j = a_{n-1} b_j = \cdots = a_{n-l+1} b_j = 0, \tag{4.146}$$

for every $0 \leq j \leq m$. Let's look at the coefficient of x^{m+n-l} in (4.143).

$$a_n b_{m-l} + a_{n-1} b_{m-l+1} + a_{n-2} b_{m-l+2} + \cdots + a_{n-l} b_m = 0. \tag{4.147}$$

By (4.146), all the terms of (4.147) except the last one is 0. Therefore, the last one $a_{n-l} b_m$ also has to be 0. Therefore, $\deg(a_{n-l} g) < m$. But

$$f \cdot (a_{n-l} g) = a_{n-l} \cdot (fg) = 0, \tag{4.148}$$

and by the minimality of $\deg g$, we must have that $a_{n-l} g = 0$. Therefore, by induction, $a_i g = 0$ for each $i = 0, 1, \ldots, n$. As a result, $a_i b_j = 0$ for each $0 \leq i \leq n, 0 \leq j \leq m$.

Since $g \neq 0$, it has a nonzero coefficient, and we can take that to be b, and we'll have $b a_i = 0$ for each $0 \leq i \leq n$. ■

Exercise 4.54. Let k be a field. Show that $k[x_1, x_2]$ is not a principal ideal domain.

Solution. The ideal (x_1, x_2) is not a principal ideal. If if were, $(x_1, x_2) = (f)$ for some $f \in k[x_1, x_2]$. Then $f \mid x_1$, and $f \mid x_2$ forces $f \in k$. But that would mean $(f) = (0)$ or $(f) = k$. So $(x_1, x_2) \neq (f)$. ■

Exercise 4.55. Prove $1 + x + x^3 + x^4$ is irreducible in $k[x]$ for any field k.

Solution.

$$1+x+x^3+x^4=(1+x)+x^3(1+x)=(1+x)\left(1+x^3\right), \tag{4.149}$$

is not irreducible in any field k. ■

Exercise 4.56. Prove x^4+2x+2 is irreducible in $\mathbb{Q}[x]$.

Solution. We'll apply Eisenstein's criterion. $2 \mid a_0, a_1, a_2, a_3$, $2 \nmid a_4$, $2^2 \nmid a_0$. So by Eisenstein's criterion, x^4+2x+2 is irreducible in $\mathbb{Q}[x]$. ■

Exercise 4.57. Let p be a prime number, $c \in \mathbb{Q}$. Show $x^p - c$ is irreducible over $\mathbb{Q}$ if and only if it has no zero in $\mathbb{Q}$.

Solution. If it's irreducible over $\mathbb{Q}$, it doesn't have any root in $\mathbb{Q}$. Now, conversely, suppose $f(x) = x^p - c$ has no root in $\mathbb{Q}$. Assume for the sake of contradiction that $x^p - c$ is reducible, i.e $x^p - c = fg$ for some $f, g \in \mathbb{Q}[x]$ with $\deg f, \deg g \geq 1$. We assume both f and g are monic, since otherwise their leading terms would be a and a^{-1}, and we can make the leading terms 1 multiplying by a^{-1} and a, respectively.

Now, suppose $\deg f = n \geq 1$. It splits into linear factors in $\mathbb{C}$, say

$$f(x) = \prod_{i=1}^{n} (x - r_i). \tag{4.150}$$

Consider the number

$$r = \prod_{i=1}^{n} r_i. \tag{4.151}$$

It's a product of complex numbers, but it's $\pm$ the constant term of f. So $r \in \mathbb{Q}$. Also,

$$r^p = \prod_{i=1}^{n} r_i^p = \prod_{i=1}^{n} c = c^n. \tag{4.152}$$

Since $n < p$, $\gcd(n, p) = 1$. So we can find $a, b \in \mathbb{Z}$ with $ap + bn = 1$. Then

$$\left(c^a r^b\right)^p = c^{ap}\left(r^p\right)^b = c^{ap}\left(c^n\right)^b = c^{ap+bn} = c. \tag{4.153}$$

This $c^a r^b$ is a rational number, which is a root of $x^p - c$. Therefore, assuming $x^p - c$ is reducible, we got that a root must exist in $\mathbb{Q}$. Hence, if no roots exist in $\mathbb{Q}$, $x^p - c$ must be irreducible. ■

Exercise 4.58. Let k be a field, $a, b \in k$. Show that $f \in k[x]$ is irreducible if and only if $f(ax+b)$ is irreducible.

Solution. Here, we need the condition that $a \neq 0$. For $a \neq 0$, $x \mapsto ax + b$ is an automorphism of $k[x]$ (Exercise 4.26). We'll show that automorphism preserves irreducible elements. Indeed, if

$f = gh$ is reducible, where g and h are not units, then under an automorphism Ψ,

$$\Psi(f) = \Psi(g)\,\Psi(h), \tag{4.154}$$

and $\Psi(g)$ and $\Psi(h)$ are not units, because if they were, then their inverse images would also be units, since Ψ is an automorphism. Therefore, $\Psi(f)$ is also reducible.

Similarly, one can show that $\Psi(f)$ reducible implies f reducible, by taking Ψ^{-1} in the previous argument. Therefore, f irreducible if and only if $\Psi(f)$ irreducible. ■

Exercise 4.59. Prove that the kernel of the homomorphism $\varphi : \mathbb{Z}[x] \to \mathbb{R};\ x \mapsto 1 + \sqrt{2}$ is a principal ideal and find a generator for this ideal.

Solution. For notational simplicity, let's write $\alpha := 1 + \sqrt{2}$. Then $\alpha - 1 = \sqrt{2}$, so that

$$(\alpha - 1)^2 - 2 = 0. \tag{4.155}$$

So $f(\alpha) = 0$ for $f(x) = x^2 - 2x - 1 \in \mathbb{Z}[x]$. Furthermore, this is the minimal polynomial for α, since there's no degree-1 polynomial whose root is α. We claim that $\operatorname{Ker}\varphi = (f)$.

Since φ sends x to α, it sends a general $g(x) \in \mathbb{Z}[x]$ to $g(\alpha)$. Therefore, if $g \in (f)$, then $g = fh$ for some $h \in \mathbb{Z}[x]$. Then

$$\varphi(g) = g(\alpha) = f(\alpha)\,h(\alpha) = 0, \tag{4.156}$$

so $g \in \operatorname{Ker}\varphi$. Therefore, $(f) \subseteq \operatorname{Ker}\varphi$. Now, for $g \in \operatorname{Ker}\varphi$, we perform an Euclidean division on g by f:

$$g = fq + r, \tag{4.157}$$

where $r = 0$ or $\deg r < \deg f = 2$. $fq \in \operatorname{Ker}\varphi$ (as shown above), so that $r \in \operatorname{Ker}\varphi$. But if $r \neq 0$, then $\deg r = 0, 1$, in which case $r(\alpha)$ can't be 0. Hence, $r = 0$, so $g = fq \in (f)$.

Therefore, $\operatorname{Ker}\varphi = (f) = \left(x^2 - 2x - 1\right)$. ■

Exercise 4.60. Prove that the following polynomials are irreducible in $\mathbb{Q}[x]$.

(a) $x^2 + 27x + 213$.

(b) $x^3 + 6x + 12$.

(c) $x^5 - 3x^4 + 3$.

Solution. (a) We'll apply Eisenstein's criterion. $3 \mid a_0, a_1$, $3 \nmid a_2$, $3^2 \nmid a_0$. So by Eisenstein's criterion, $x^2 + 27x + 213$ is irreducible in $\mathbb{Q}[x]$.

(b) $3 \mid a_0, a_1, a_2$, $3 \nmid a_3$, $3^2 \nmid a_0$. So by Eisenstein's criterion, $x^3 + 6x + 12$ is irreducible in $\mathbb{Q}[x]$.

(Choosing $p = 2$ wouldn't have worked here since $2^2 \mid 12$.)

(c) $3 \mid a_0, a_1, a_2, a_3, a_4$, $3 \nmid a_5$, $3^2 \nmid a_0$. So by Eisenstein's criterion, $x^5 - 3x^4 + 3$ is irreducible in $\mathbb{Q}[x]$.

■

Exercise 4.61. Factor $x^5 + 5x + 5$ into irreducible factors in $\mathbb{Q}[x]$ and $\mathbb{F}_2[x]$.

Solution. For $\mathbb{Q}$, we'll apply Eisenstein's criterion. $5 \mid a_0, a_1, a_2, a_3, a_4$, $5 \nmid a_5$, $5^2 \nmid a_0$. So by Eisenstein's criterion, $x^5 + 5x + 5$ is irreducible in $\mathbb{Q}[x]$.

In $\mathbb{F}_2[x]$,

$$f(x) = x^5 + 5x + 5 = x^5 + x + 1. \tag{4.158}$$

$f(0) = f(1) = 1$, so f doesn't have any linear factors. Let's check if it has any quadratic factors. The only irreducible quadratic in $\mathbb{F}_2[x]$ is $x^2 + x + 1$. After a polynomial long division, we find that $x^2 + x + 1$ divides f:

$$x^5 + x + 1 = \left(x^2 + x + 1\right)\left(x^3 + x^2 + 1\right). \tag{4.159}$$

$x^3 + x^2 + 1$ is irreducible because it has no roots in $\mathbb{F}_2$. ■

Exercise 4.62. Suppose that a polynomial $x^4 + bx^2 + c \in \mathbb{Q}[x]$ is the product of two factors of degree 2 in $\mathbb{Q}[x]$. What can one say about the coefficients of these factors?

Solution. Suppose

$$x^4 + bx^2 + c = \left(x^2 + px + q\right)\left(x^2 + rx + s\right) = x^4 + (p + r)x^3 + (q + s + pr)x^2 + (ps + rq)x + qs. \tag{4.160}$$

Since the coefficients of x^3 is 0, we have $r = -p$. Substituting this, we get

$$q + s - p^2 = b \tag{4.161}$$

$$p(s - q) = 0 \tag{4.162}$$

$$qs = c \tag{4.163}$$

Now we have two cases:

- if $p = 0$, then
$$x^4 + bx^2 + c = \left(x^2 + q\right)\left(x^2 + s\right) \tag{4.164}$$
with $q + s = b$ and $qs = c$.
- if $p \neq 0$, then we must have $s - q = 0$. Then,
$$x^4 + bx^2 + c = \left(x^2 + px + q\right)\left(x^2 - px + q\right), \tag{4.165}$$
where $q^2 = c$, $2q - p^2 = b$.

■

Exercise 4.63. Factor the following polynomials in $\mathbb{Q}[x]$

(a) $x^3 + 2x^2 + 3x + 1$.

(b) $x^4 + 2x^3 + 3x^2 + 2x + 1$.

Solution. (i) Reduction modulo 2 gives rise to the polynomial $x^3 + x + 1$. If this were reducible, it would have a linear factor, i.e. it would have a root. Plugging $0, 1$ in $x^3 + x + 1$, we get $1 \neq 0$ both times. So $x^3 + x + 1$ has no roots in $\mathbb{F}_2$. Therefore, it's irreducible in $\mathbb{F}_2[x]$, and as a result, it's irreducible in $\mathbb{Q}[x]$.

(ii) Reduction modulo 2 gives $x^4 + x^2 + 1$, which is (Fresher's dream[1])

$$x^4 + x^2 + 1 = \left(x^2 + x + 1\right)^2. \tag{4.166}$$

In $\mathbb{Q}[x]$, one can check that

$$\left(x^2 + x + 1\right)^2 = x^4 + x^2 + 1 + 2x^3 + 2x^2 + 2x = x^4 + 2x^3 + 3x^2 + 2x + 1. \tag{4.167}$$

$x^2 + x + 1$ can't be factored further since it's irreducible.

■

Exercise 4.64. Let $p \in \mathbb{Z}$ be a prime number. Let $f = \sum_{i=0}^{2n+1} a_i x^i \in \mathbb{Z}[x]$ be a polynomial of odd degree. Suppose $p \nmid a_{2n+1}$, $p^2 \mid a_0, \ldots, a_n$, $p \mid a_{n+1}, \ldots, a_{2n}$ and $p^3 \nmid a_0$. Show that f is irreducible in $\mathbb{Q}[x]$.

Solution. By Gauss' lemma, f is irreducible in $\mathbb{Q}[x]$ if and only if it's irreducible in $\mathbb{Z}[x]$. Assume for the sake of contradiction that f is reducible in $\mathbb{Z}[x]$. Then $f = gh$ for $g, h \in \mathbb{Z}[x]$. Reducing to mod p, we obtain

$$gh \equiv a_{2n+1} x^{2n+1}. \tag{4.168}$$

Therefore, other than the leading terms of g and h, the rest of those polynomaials are divisible by p.

$$g = Ax^{m_1} + pg', \quad h = Bx^{m_2} + ph', \tag{4.169}$$

where $p \nmid A, B$ since $p \nmid AB = a_{2n+1}$. Then

$$f = gh = AB\ x^{m_1+m_2} + Ap\ x^{m_1} h' + Bp\ x^{m_2} g' + p^2\ g'h'. \tag{4.170}$$

WLOG, $m_1 \leq m_2$, then $m_1 \leq n$. Now, the coefficient of x^{m_1} modulo p^2 is

$$a_{m_1} \equiv Ap\ h'(0) \pmod{p^2}. \tag{4.171}$$

[1] Although the term is Freshman's dream, my obsession with using gender-neutral language knows no bounds.

Since $m_1 \leq n$, $a_{m_1} \equiv 0 \pmod{p^2}$. Since $p \nmid A$, then we must have $p \mid h'(0)$. But then the constant term of f would be

$$a_0 = p^2 \; g'(0) \; h'(0), \tag{4.172}$$

divisible by p^3, which is a contradiction! ■

§5 Problem Set 3

All rings in this problem sheet are commutative with 1. Morphisms of rings send 1 to 1.

Exercise 5.1. Let R be the ring $\mathbb{Z} \times \mathbb{Z}$.

(a) Show that every ideal of R is principal.

(b) Describe all prime and maximal ideals of R.

Solution. First, let's show that any ideal in a product ring $R \times S$ is of the form $I \times J$ for $I \trianglelefteq R$ and $J \trianglelefteq S$. Let $\pi_1 : R \times S \to R$ and $\pi_2 : R \times S \to S$ be projections, which are surjective homomorphisms. If $\mathfrak{a} \trianglelefteq R \times S$, then $\pi_i(\mathfrak{a})$ are also ideals, since π_i are surjective homomorphisms. Let

$$I = \pi_1(\mathfrak{a}) \trianglelefteq R, \quad J = \pi_2(\mathfrak{a}) \trianglelefteq S. \tag{5.1}$$

Then $\mathfrak{a} \subseteq I \times J$. Now, given any $(r, s) \in I \times J$, there exists some $s' \in S$ such that $(r, s') \in \mathfrak{a}$, and there exists some $r' \in R$ such that $(r', s) \in \mathfrak{a}$. We have

$$\left(r, ss'\right) = (1, s)\left(r, s'\right) \in \mathfrak{a}, \text{ and } \left(r', ss'\right) = \left(1, s'\right)\left(r', s\right) \in \mathfrak{a}. \tag{5.2}$$

Therefore, their difference, $(r - r', 0) \in \mathfrak{a}$. As a result,

$$\left(r - r', 0\right) + \left(r', s\right) = (r, s) \in \mathfrak{a}. \tag{5.3}$$

Hence, $\mathfrak{a} = I \times J$.

(a) Now, any ideal of $\mathbb{Z} \times \mathbb{Z}$ is of the form $I \times J$ for $I, J \trianglelefteq \mathbb{Z}$. And since $\mathbb{Z}$ is a PID, we have $I = (m)$ and $J = (n)$ for some $m, n \in \mathbb{Z}$. We claim that $I \times J = \langle (m, n) \rangle$.

$(m, n) \in I \times J$, so $\langle (m, n) \rangle \subseteq I \times J$. And any element of $I \times J$ is of the form (am, bn), for some $a, b \in \mathbb{Z}$. Then

$$(am, bn) = (a, b) \cdot (m, n) \in \langle (m, n) \rangle. \tag{5.4}$$

Therefore, $I \times J = \langle m \rangle \times \langle n \rangle = \langle (m, n) \rangle$.

(b) Let $I \times J$ be an ideal of $\mathbb{Z} \times \mathbb{Z}$. Then

$$\frac{\mathbb{Z} \times \mathbb{Z}}{I \times J} \cong \frac{\mathbb{Z}}{I} \times \frac{\mathbb{Z}}{J}. \tag{5.5}$$

The products of two rings is an integral domain if and only if one the products is 0 and the other is an ID. Because otherwise, if both are nonzero, then

$$(1, 0) \cdot (0, 1) = (0, 0) \tag{5.6}$$

would lead to nonzero zero-divisors. Therefore, in order for $\frac{\mathbb{Z}}{I} \times \frac{\mathbb{Z}}{J}$ to be an ID, we must have that one of these is 0, WLOG, $\frac{\mathbb{Z}}{J} = 0$, i.e. $J = \mathbb{Z}$.

Then $\frac{\mathbb{Z}}{I}$ is an ID if and only if $I = (0)$ or $I = (p)$ for prime p. And $\frac{\mathbb{Z}}{I}$ is a field if and only if $I = (p)$ for prime p. Therefore, the prime ideals of $\mathbb{Z} \times \mathbb{Z}$ are

$$(0) \times \mathbb{Z},\ (p) \times \mathbb{Z},\ \mathbb{Z} \times (0),\ \mathbb{Z} \times (p); \tag{5.7}$$

and the maximal ideals are

$$(p) \times \mathbb{Z},\ \mathbb{Z} \times (p). \tag{5.8}$$

■

Exercise 5.2. Let R be a ring. Define

$$J(R) = \bigcap_{\mathfrak{m} \trianglelefteq R \text{ maximal ideal}} \mathfrak{m}. \tag{5.9}$$

This ideal is called **Jacobson radical**.

(a) Show that $x \in J(R)$ if and only if $1 + xy$ is invertible for every $y \in R$.

(b) Let $\varphi : R \to S$ be a surjective homomorphism of rings. Show that $\varphi(J(R)) \subseteq J(S)$.

Solution. (a) Suppose $x \in J(R)$ and Assume for the sake of contradiction that there exists some $y \in R$ such that $1 + xy$ is not invertible. Then $1 + xy$ is contained in a maximal ideal $\mathfrak{m}$ (since the proper ideal $(1 + xy)$ is contained in a maximal ideal, assuming axiom of choice). $x \in J(R)$ implies $x \in \mathfrak{m}$, so we have $xy \in \mathfrak{m}$ since it's an ideal. But that gives us

$$1 = (1 + xy) - xy \in \mathfrak{m}, \tag{5.10}$$

which is not possible since maximal ideals are proper ideals.

Conversely, suppose $1 + xy$ is invertible for every $y \in R$, and assume for the sake of contradiction that $x \notin J(R)$. Then there exists a maximal ideal $\mathfrak{m}$ that doesn't contain x. Now,

$$(x) + \mathfrak{m} \supsetneq \mathfrak{m}, \tag{5.11}$$

so by the maximality of $\mathfrak{m}$, it has to be the whole R. Therefore, there exists $y \in R$ and $m \in \mathfrak{m}$ such that

$$xy + m = 1. \tag{5.12}$$

That would give us $m = 1 - xy$, which is invertible by hypothesis. But that can't happen, since maximal ideals can't contain invertible elements.

(b) Let $y \in \varphi(J(R))$. Then $y = \varphi(x)$ for some $x \in J(R)$. We'll show that $1 + sy$ is invertible for any $s \in S$, which will prove that $y \in J(S)$.

Given $s \in S$, since φ is surjective, there exists $r \in R$ such that $s = \varphi(r)$. Let $u = (1 + rx)^{-1}$. Then

$$f(u)[1 + sy] = f(u) f(1 + rx) = f(u(1 + rx)) = f(1) = 1. \tag{5.13}$$

Hence, $1 + sy$ is invertible, and thus $y \in J(R)$. This proves that $\varphi(J(R)) \subseteq J(S)$.

■

Exercise 5.3. Let R be the ring $\underbrace{\mathbb{Z} \times \mathbb{Z} \times \cdots \times \mathbb{Z}}_{n \text{ times}} =: \mathbb{Z}^n$.

(a) Show that R has a finite number of ring automorphisms.

(b) Find the number of automorphisms.

Solution. Let $e_i = (0, \ldots, 0, 1, 0, \ldots, 0)$ where we have 1 at the i-th component and 0 everywhere else. Note that $e_i^2 = e_i$, so e_i is an idempotent. An automorphism $\mathbb{Z}^n \to \mathbb{Z}^n$ will map idempotents to idempotents. The only idempotents of $\mathbb{Z}^n$ are vectors consisting of 0 and 1 only. Therefore,

$$f(e_i) = \left(c_1^{(i)}, \ldots, c_n^{(i)}\right), \tag{5.14}$$

where $c_j^{(i)}$'s are either 0 or 1.

Furthermore, if $i \neq j$, then $e_i \cdot e_j = 0$. So $f(e_i) \cdot f(e_j) = 0$. Therefore, for $i \neq j$, $f(e_i)$ and $f(e_j)$ can't contain 1 in the same coordinates.

Let $O_i \subseteq \{1, \ldots, n\}$ be the collection of indices where $f(e_i)$'s component is 1, i.e.

$$O_i = \left\{ j \mid c_j^{(i)} = 1 \right\}. \tag{5.15}$$

Then the O_i's are pairwise disjoint, as argued above. Also, each O_i is nonempty, since $\operatorname{Ker} f$ is trivial. Now,

$$\sum_{i=1}^{n} f(e_i) = f\left(\sum_{i=1}^{n} e_i\right) = f(1, \ldots, 1) = (1, \ldots, 1). \tag{5.16}$$

Also, the j-th component of $\sum_{i=1}^{n} f(e_i)$ is

$$\sum_{i=1}^{n} c_j^{(i)} = 1, \tag{5.17}$$

so that means exactly one $c_j^{(i)} = 1$ and the rest are all 0's. Therefore, the set $\{f(e_i)\}_{i=1}^n$ is just a permutation of $\{e_i\}_{i=1}^n$.

Each $f \in \operatorname{Aut}(\mathbb{Z}^n)$ corresponds to one permutation, and each permutation corresponds to one automorphism. Hence, $|\operatorname{Aut}(\mathbb{Z}^n)| = n!$. ■

Exercise 5.4. Let $\varphi : R \to S$ be a morphism of rings. Prove or show a counterexample of the following:

(a) If $\mathfrak{p}_1, \mathfrak{p}_2$ are prime ideals of R, then $\mathfrak{p}_1 + \mathfrak{p}_2$ is a prime ideal of R.

(b) If $\sqrt{I}$ is a prime ideal of R, then I is a prime ideal.

(c) If $\mathfrak{p} \trianglelefteq S$ is a prime ideal, then $\varphi^{-1}(\mathfrak{p}) \trianglelefteq R$ is a prime ideal.

(d) If $\mathfrak{m} \trianglelefteq S$ is a maximal ideal, then $\varphi^{-1}(\mathfrak{m}) \trianglelefteq R$ is a maximal ideal.

Solution. (a) False. Take $\mathfrak{p}_1 = (p_1), \mathfrak{p}_2 = (p_2)$ prime ideals of $R = \mathbb{Z}$, where p_1, p_2 are distinct primes. Then $(p_1) + (p_2) = (1) = \mathbb{Z}$ is the whole ring, which is not a prime ideal.

(b) False. Take $I = \left(p^2\right)$ in $R = \mathbb{Z}$, which is not a prime ideal. Then

$$\sqrt{(p^2)} = \left\{a \in \mathbb{Z} \mid a^n \in \left(p^2\right) \text{ for some } n > 0\right\} = \{a \in \mathbb{Z} \mid a \in (p)\} = (p), \tag{5.18}$$

is a prime ideal.

(c) True. The proof is written in Exercise 4.40(b).

(d) False. Take $\mathbb{Z} \hookrightarrow \mathbb{Q}$. The only maximal ideal of $\mathbb{Q}$ is (0), whose preimage is (0), and it's not maximal in $\mathbb{Z}$.

More generally, we can take any injective homomorphism $R \to \mathbb{F}$ where R is NOT a field and $\mathbb{F}$ is a field.

■

Exercise 5.5. Let R be a ring such that $x = x^2$ for all $x \in R$.

(a) Show that $x + x = 0$ for all $x \in R$.

(b) Show that every finitely generated ideal of R is principal.

Solution. (a)

$$x + x = (x + x)^2 = x^2 + x^2 + x^2 + x^2 = x + x + x + x. \tag{5.19}$$

Cancelling out $x + x$, we get $x + x = 0$.

(b) It suffices to show that every ideal (x, y) generated by two elements is principal. We claim that

$$(x, y) = (x + y + xy). \tag{5.20}$$

Indeed, $x + y + xy \in (x, y)$, so $(x + y + xy) \subseteq (x, y)$. Now,

$$x(x + y + xy) = x^2 + xy + x^2y = x + xy + xy = x. \tag{5.21}$$

So $x \in (x + y + xy)$. Similarly, $y \in (x + y + xy)$. Hence, $(x + y + xy) = (x, y)$.

Then by induction, $(x_1, \ldots, x_n)$ is principal.

■

Exercise 5.6. Let R be a ring.

(a) Assume that every ideal $I \trianglelefteq R$ satisfies $I = \sqrt{I}$. Show that if $J, K \trianglelefteq R$ are ideals and there exists an integer $m > 0$ such that $J^m = K^m$, then $J = K$.

(b) Assume instead that there exists an ideal $I \trianglelefteq R$ such that $I = I^2$ and $I \neq \sqrt{I}$. Show that there exist two distinct ideals $J \neq K$ such that $J^m = K^m$ for all $m > 1$.

(c) Assume now that there exists an ideal $I \trianglelefteq R$ such that $I \neq \sqrt{I}$ and the chain of ideals

$$I \supseteq I^2 \supseteq I^3 \supseteq \cdots \tag{5.22}$$

is stationary. Prove that there exists an ideal $I_0 \trianglelefteq R$ such that $I_0 = I_0^2$, and $\sqrt{I_0} \neq I_0$.

Solution. (a) Firstly, note that $\sqrt{I^m} = \sqrt{I}$. Indeed, since $I^m \subseteq I$, we have $\sqrt{I^m} \subseteq \sqrt{I}$. Now, for $a \in \sqrt{I}$, $a^n \in I$ for some I. Then

$$a^{mn} = a^n \cdot a^n \cdot \cdots \cdot a^n \in I^m, \tag{5.23}$$

so $a \in \sqrt{I^m}$. So $\sqrt{I} \subseteq \sqrt{I^m}$.

Now, since $J^m = K^m$, we have

$$J = \sqrt{J} = \sqrt{J^m} = \sqrt{K^m} = \sqrt{K} = K. \tag{5.24}$$

(b) Since $I \neq \sqrt{I}$, we can find $z \in \sqrt{I} - I$. Then $z^n \in I$, and assume n is the minimal, so $n > 1$. Now, we take

$$x = z^{\lceil \frac{n}{2} \rceil} \notin I. \tag{5.25}$$

Then $x \in \sqrt{I}$ since $x^2 = z^n$ (or z^{n+1}) is in I. We take

$$J = I, \quad K = I + (x)\,. \tag{5.26}$$

Then $J \neq K$, since $x \in K$ and $x \notin J$. We'll prove that $J^m = K^m$ for each $m > 1$. Indeed, since $I^2 = I$, $J^m = I^m = I$ for each $m \geq 2$.

Now, since $I \subseteq K$, $I = I^2 \subseteq K^2$. Also, a general element of K^2 looks like

$$(i + rx)\left(i' + r'x\right) \tag{5.27}$$

for $i, i' \in I$ and $r, r' \in R$. Simplifying, we get

$$(i + rx)\left(i' + r'x\right) = ii' + rx\ i' + r'x\ i + rr'\ x^2, \tag{5.28}$$

and each term is in I, so $K^2 \subseteq I$. Therefore, $K^2 = I = J^2$.

$I \subseteq K$ gives $I = I^m \subseteq K^m$. Suppose $K^{m-1} = I$. Then $K^m = IK \subseteq I$. Therefore, $K^m = I$ provided $K^{m-1} = I$. Since we already proved that $K^2 = I$, we have $K^m = I$ for each $m \geq 2$.

(c) Since the chain $I \supseteq I^2 \supseteq I^3 \supseteq \cdots$ is stationary, there exists some $k \geq 1$ such that

$$I^k = I^{k+1} = I^{k+2} = I^{k+3} = \cdots\,. \tag{5.29}$$

We take $I_0 = I^k$. Then

$$I_0^2 = \left(I^k\right)^2 = I^{2k} = I^k = I_0. \tag{5.30}$$

Since $I \neq \sqrt{I}$, we can find $x \in \sqrt{I}$, $x \notin I$. Then $x \notin I^k = I_0$ since I contains I^k. And $x \in \sqrt{I} = \sqrt{I^k} = \sqrt{I_0}$.

Therefore, we found I_0 such that $I_0 = I_0^2$, and $I_0 \neq \sqrt{I_0}$.

■

Exercise 5.7. Let G be a group and p a prime number, and $|G| = p^n$ for some integer $n > 0$.

(a) Show that G has a subgroup of order p^m for all $0 \leq m \leq n$.

(b) Show that G has a normal subgroup of order p^m for all $0 \leq m \leq n$.

Solution. (a) This is just Exercise 3.23.

(b) We proceed by induction on n. The base case $n = 1$ is trivial. Suppose the statement is true for $1, \ldots, n-1$.

Every p-group has a nontrivial center (this can be seen by the class-equation.) Therefore, $Z(G)$ contains an element of order p (by Cauchy's theorem), say a. Then consider $N = \langle a \rangle$, which is a normal subgroup of order p.

So we can consider the quotient G/N which has order p^{n-1}, and the quotient map $\pi : G \to G/N$. By inductive hypothesis, G/N has a normal subgroup N_i of order p^i for each $0 \leq i \leq n-1$. Then we just consider $\pi^{-1}(N_i) \trianglelefteq G$, which is normal since preimage of normal subgroups are normal.

$$\left|\pi^{-1}(N_i)\right| = |N_i|\,|\mathrm{Ker}\,\pi| = p^i \cdot p = p^{i+1}. \tag{5.31}$$

Therefore, G has a normal subgroup of order p^m for each $1 \leq m \leq n$. THe normal subgroup of order p^0 is the trivial one, $\{e\}$.

■

§6 Problem Set 4

Exercise 6.1. Factor the following polynomials in irreducible factors in $\mathbb{Z}[x]$.

(a) $x^4 + x^2 - x + 2$;

(b) $(x-1)^3 - 2$;

(c) $x^3 + x - 1$.

Solution. (a) By checking $f(\pm 1)$, $f(\pm 2)$, we can see that it has no rational root. So the irreducible factors must have degree 2. But in $\mathbb{F}_2[x]$, the polynomial is

$$x^4 + x^2 + x = x\left(x^3 + x + 1\right), \tag{6.1}$$

and $x^3 + x + 1$ is irreducible in $\mathbb{F}_2$ since it doesn't have any roots. Since in $\mathbb{F}_2[x]$, the polynomial has a degree-1 factor, it would have a degree-1 factor in $\mathbb{Z}[x]$ as well, but that's not the case. Therefore, $x^4 + x^2 - x + 2$ is irreducible.

(b) This is $y^3 - 2$, which is irreducible by Eisenstein's criterion.

(c) We reduce it to $\mathbb{F}_2$. If it was reducible, it would have a degree-1 factor, i.e. a root in $\mathbb{F}_2$. But no element of $\mathbb{F}_2$ is a root of $x^3 + x - 1$, so it's irreducible over $\mathbb{F}_2$, and hence it's irreducible over $\mathbb{Z}$.

■

Exercise 6.2.

Solution. ■

Exercise 6.3.

Solution. ■

Exercise 6.4.

Solution. ■

Exercise 6.5.

Solution. ■

Exercise 6.6.

Solution. ■