

**Table 14-5.02**  
**Incidence of Treatment Emergent Serious Adverse Events by Treatment Group**

| System Organ Class/<br>Preferred Term | Placebo<br>(N=86) |       | Xanomeline<br>Low Dose<br>(N=84) |       | Xanomeline<br>High Dose<br>(N=84) |       | Fisher's Exact<br>p-values |                             |
|---------------------------------------|-------------------|-------|----------------------------------|-------|-----------------------------------|-------|----------------------------|-----------------------------|
|                                       | n(%)              | [AEs] | n(%)                             | [AEs] | n(%)                              | [AEs] | Placebo<br>vs.<br>Low Dose | Placebo<br>vs.<br>High Dose |
| ANY BODY SYSTEM                       | 0                 |       | 1 ( 1.2%)                        | [1]   | 2 ( 2.4%)                         | [2]   | 0.494                      | 0.243                       |
| NERVOUS SYSTEM DISORDERS              | 0                 |       | 1 ( 1.2%)                        | [1]   | 2 ( 2.4%)                         | [2]   | 0.494                      | 0.243                       |
| PARTIAL SEIZURES WITH                 | 0                 |       | 0                                |       | 1 ( 1.2%)                         | [1]   |                            | 0.494                       |
| SECONDARY GENERALISATION              | 0                 |       | 1 ( 1.2%)                        | [1]   | 1 ( 1.2%)                         | [1]   | 0.494                      | 0.494                       |
| SYNCOPE                               | 0                 |       | 1 ( 1.2%)                        | [1]   | 1 ( 1.2%)                         | [1]   | 0.494                      | 0.494                       |

Note: Treatment emergent events are defined as events which start on or after the start of treatment.

Note: Adverse events are coded using MedDRA.

Note: Percentages are based on the number of subjects in the safety population within each treatment group.

Note: P-values are based on Fisher's Exact test for the comparison of placebo versus each active treatment group. An asterisk is appended to p-values that are less than 0.15.

Note: The column [AE] represents the total number of times an event was recorded.

Source: programs/t-14-5-02.R

19:30 Tuesday, June 16, 2020