

Solving Problems with Desmos

A Complete Student Guide – CLC Educational Institute

What We Will Cover Today

01

Getting Started

Interface, navigation, saving & sharing

02

Algebra

Points, lines, equations, inequalities, tables

03

Functions & Graphing

Quadratics, sliders, piecewise, parametric, polar

04

Geometry

Shapes, conics, constructions, Geometry tool

05

Calculus

Derivatives, integrals, limits

06

Statistics

Data, regression, distributions

07

Tips & Tricks

Shortcuts, mobile, pro features

Getting Started

Understanding the Desmos interface


What is Desmos?

- Desmos is a free, browser-based graphing calculator — no download or installation needed.
- Access it at desmos.com/calculator on any computer, tablet, or smartphone.
- It plots points, graphs equations, evaluates functions, tables, and more — all in real time.
- Used by millions of students worldwide and accepted in many standardized exams.
- Also includes a Geometry tool, Scientific Calculator, and classroom activities.
- All your work can be saved and shared with a simple URL — no account required to start.

TIP: Desmos works in any modern browser. Bookmark desmos.com/calculator so you always have it ready.

A Tour of the Desmos Interface

- Expression List (left panel): where you type equations, points, and notes.
- Graph (right): updates instantly as you type.
- Colored circle: click to change color or hide a graph.
- Keyboard (bottom): access math symbols, Greek letters, functions.
- Wrench icon (top-right of graph): change axes, grid, angle units.
- Home button: resets the view to fit all graphs on screen.
- Share button: save and share your graph as a URL or image.



TIP: Click directly on any curve on the graph to see its key points.

Why Use Desmos?

- Completely FREE — no subscription, no login required to start.
- Instant visual feedback: you see the graph the moment you type.
- Interactive sliders let you explore how changing a number changes the graph.
- Handles everything from basic arithmetic to calculus and statistics.
- Works on computers, iPads, and Android tablets — great for home study.
- Accepted on many international assessments and curricula.
- If you can write a math formula, Desmos can graph it.

DESMOS TOOLS

Graphing Calculator
Scientific Calculator
Four-Function Calc
Matrix Calculator
Geometry Tool
Classroom Activities
3D Calculator (beta)

Navigating the Graph

- Pan: click and drag anywhere on the graph to move the view.
- Zoom: use the + and - buttons, or scroll your mouse wheel.
- Home button: press to auto-fit the view so all graphs are visible.
- Set exact range: click the Wrench icon and enter x-min, x-max, y-min, y-max.
- Points of interest appear automatically — hover to see exact coordinates.
- Click a point to pin its label so it stays visible on screen.
- Audio trace: press Alt+T to hear a graph as sound.

ZOOM SHORTCUTS

Scroll wheel = zoom
Ctrl + = zoom in
Ctrl - = zoom out
Home btn = auto-fit
Drag = pan

Wrench = set range

Saving and Sharing Your Work

- Save button (top-left): saves your graph — prompts login if not signed in.
- Share button: generates a unique URL — paste it anywhere to share.
- Export image: save as PNG, SVG, or even Braille embosser format.
- Snapshot: a static image of your current graph view.
- Re-open saved graphs by clicking the folder icon (top-left) after sign-in.
- No account? The URL in your browser bar reflects your latest graph — bookmark it.

TIP: Always copy your graph's URL before closing the tab if you don't have an account.

Essential Keyboard Shortcuts

New Line

Enter

Undo

Ctrl + Z

Redo

Ctrl + Y

Duplicate

Ctrl+Shift+C

Deg / Rad toggle

Alt + D

Superscript (^)

^ key

Fraction

/ key

Show Shortcuts

Ctrl + /

Basic Algebra

Equations, lines, inequalities & tables

Plotting Points


Type a coordinate pair directly into the expression list.

```
(3, 4)      <- plots a point at x=3, y=4  
(-2, 0)     <- plots a point on the x-axis  
  
// Multiple points – list them on one line:  
(-1,1), (0,0), (1,1), (2,4)  
  
// Or use a Table: click + -> Table  
//   and enter x and y columns
```

TIP: Click any plotted point to see its exact coordinates. You can drag points to move them.

Graphing Lines and Finding Intersections


- Type any equation involving x and y — Desmos draws it instantly.
- $y = 2x + 1$ draws a line with slope 2, y -intercept 1.
- $y = -x + 4$ draws a line with slope -1.
- Type TWO equations and Desmos automatically shows their intersection point.
- Click the intersection dot to read the exact (x, y) solution.
- Vertical line: type $x = 3$. Horizontal: type $y = -2$.



TIP: Hover over the intersection point to read the exact coordinates.

Solving Linear Equations Graphically

- Method: type both sides of the equation as $y = \dots$
- Example: solve $2x + 3 = 7$
- Line 1: $y = 2x + 3$
- Line 2: $y = 7$
- The x-coordinate of the intersection = the solution.
- Click the point to confirm: $x = 2$.
- You can check any hand-calculated answer this way instantly.



TIP: All three methods — graphical, table, direct equation — give the same answer.

Systems of Equations


- Type each equation on a separate line — Desmos plots both and marks the intersection.
- The intersection point (x, y) is the solution to the system.
- Click the intersection dot to read the exact coordinates — no calculation needed.
- Works for 2 or more equations simultaneously.
- Parallel lines = no intersection = no solution.
- Overlapping lines = infinitely many solutions.
- Try adding a slider to see how changing a coefficient moves the solution.

TIP: Add $x = \dots$ or $y = \dots$ as a check line to verify each equation in the system.

Graphing Inequalities

- Replace = with $>$ $<$ $>=$ $<=$ — Desmos shades the solution region.
- $y > 2x - 1$ shades ABOVE the dashed line.
- $y \leq x + 3$ shades BELOW with a solid boundary.
- Type two inequalities together — Desmos shows the overlapping region.
- Dashed boundary = strict inequality ($<$ or $>$).
- Solid boundary = includes the line ($\leq$ or $\geq$).
- The shaded overlap is the solution to the system.

TIP: Type $\geq$ for $\geq$ and $\leq$ for $\leq$ — Desmos converts them automatically.



Using Tables in Desmos

- Click + then Table to add a data table to the expression list.
- Enter x values in the left column — y values fill automatically if you set a function.
- Or enter both x and y manually to create a scatter of data points.
- Press Enter inside the table to add more rows.
- Type $y_1 = f(x_1)$ to link a table column to any function you defined.
- Tables are the starting point for regression — enter your data here, then fit a curve.
- You can paste data copied from Excel or Google Sheets directly into a table.

TIP: Use tables to spot patterns: look for constant differences (linear) or constant ratios (exponential).

Functions & Graphing

Defining, evaluating, and transforming functions

Defining and Evaluating Functions

Name your own function — then call it anywhere in the expression list.

```
f(x) = x^2 - 4x + 3    <- define the function

f(2)                  <- evaluates to -1
f(0)                  <- evaluates to 3


// Plot f and a specific point together:
f(x) = x^2 - 4x + 3
(2, f(2))              <- plots the point (2, -1) on the curve
```

TIP: Use any letter: $f(x)$, $g(x)$, $h(x)$. Desmos assigns a different color to each.

Quadratic Functions

- Standard form: $f(x) = ax^2 + bx + c$
- Vertex form: $f(x) = a(x-h)^2 + k$
- Vertex is the highest or lowest point — click it to read (h, k).
- Roots (x-intercepts): click where the curve crosses the x-axis.
- Axis of symmetry: the vertical line $x = h$.
- Add sliders for a, b, c (or a, h, k) — drag and watch the parabola move.
- If the graph never crosses the x-axis, the roots are complex (no real solutions).


TIP: Type $x = (-b)/(2a)$ to plot the axis of symmetry as a dotted line.



Sliders and Transformations


- Type a letter not yet defined — Desmos offers to add a slider.
- Example: type $f(x) = a \cdot x^2 + b$ then press 'add sliders'.
- Drag the slider handle to change the value — the graph updates live.
- Press the Play button on any slider to animate it continuously.
- Set slider range and step in the slider's settings icon.
- Try $f(x) = a \cdot \sin(bx+c)+d$ to explore amplitude, period, phase, midline.
- Sliders build the deepest intuition about how formulas control graphs.

TIP: Use sliders in exams preparation: explore 'what happens if a doubles?' style questions.



Piecewise Functions

- Define functions that behave differently on different intervals.
- Use curly braces $\{ \}$ with comma-separated pieces.
- Syntax: $f(x) = \{\text{condition: expression, condition: expression}\}$
- Example: $f(x) = \{x < 0: -x, 0 \leq x \leq 2: x^2, x > 2: 4\}$
- Desmos draws each piece only in its correct interval.
- Open circles = strict inequality (not included).
- Filled circles = the endpoint is included.




TIP: Press the $\{ \}$ keys on your keyboard directly — no special menu needed.

Parametric Curves


- Graph curves defined by separate $x(t)$ and $y(t)$ expressions.
- Type $x = \cos(t)$ and $y = \sin(t)$ on consecutive lines — Desmos links them.
- A bracket appears on the left — click to set the t domain.
- Circle: $x=\cos(t)$, $y=\sin(t)$ | Spiral: $x=t\cdot\cos(t)$, $y=t\cdot\sin(t)$
- Restrict: add $0 \leq t \leq 2\pi$ to control how much is drawn.
- Parametric curves let you draw shapes impossible in $y = f(x)$ form.

TIP: Any curve that loops or doubles back on itself must be parametric or polar.



Polar Curves

- Graph equations of the form $r = f(\theta)$.
- Enable the polar grid: Wrench icon -> Grid -> Polar.
- Type 'theta' — Desmos converts it to the Greek letter automatically.
- $r = 3$ → circle of radius 3
- $r = 1 + \cos(\theta)$ → cardioid
- $r = \sin(2\theta)$ → 4-petal rose
- $r = \theta$ → Archimedean spiral




TIP: Use $0 \leq \theta \leq 4\pi$ to draw extra loops on spirals and multi-petal roses.

Geometry

Shapes, conics, constructions, and the Geometry tool

Geometry in the Graphing Calculator

- Desmos plots geometric shapes from their algebraic equations.
- Circles, triangles, polygons, and conics — all from standard equations.
- Restrict domains to draw only segments of lines or curves.
- Plot vertex points alongside segments to label shapes.
- Combine algebra and geometry on the same grid.
- Everything is interactive — drag slider points in real time.



TIP: For compass-and-ruler constructions, use the separate desmos.com/geometry tool.

Lines, Rays, and Segments

Restrict the domain of any equation using curly braces to draw just a portion.

```
y = 2x + 1          <- full infinite line
y = 2x + 1 {0 <= x <= 4} <- segment from x=0 to x=4
y = 2x + 1 {x >= 0}   <- ray starting at x=0

// Vertical segment:
x = 3 {1 <= y <= 5}   <- from (3,1) to (3,5)

// Right triangle (3 segments):
y = 0 {0<=x<=4}      x = 0 {0<=y<=3}      y = (-3/4)*x + 3 {0<=x<=4}
```

TIP: Combine multiple segments to draw any polygon — label the vertices as points.

Circles and Polygons


Type standard geometric equations — Desmos draws the shape instantly.

```
// Circle:  $(x-h)^2 + (y-k)^2 = r^2$   
 $x^2 + y^2 = 25$                 <- centre  
(0,0), radius 5  
 $(x-2)^2 + (y-3)^2 = 9$         <- centre (2,3),  
radius 3
```

```
// Triangle (three restricted lines):  
 $y=x$  { $0 \leq x \leq 4$ }   $y=4$  { $0 \leq x \leq 4$ }   $x=0$  { $0 \leq y \leq 4$ }
```

```
// Upper semicircle:  
 $y = \sqrt{16 - x^2}$           <- top half of  
circle  $r=4$ 
```


TIP: Plot vertex points like (0,0), (4,0), (0,4) alongside your segments.



Conic Sections

- Circle: $(x-h)^2 + (y-k)^2 = r^2$
- Ellipse: $(x-h)^2/a^2 + (y-k)^2/b^2 = 1$
- Hyperbola: $(x-h)^2/a^2 - (y-k)^2/b^2 = 1$
- Parabola: $y = a(x-h)^2 + k$ or $x = a(y-k)^2 + h$
- Add sliders for h , k , a , b — drag them to see how parameters change shape.
- Desmos draws the full conic from one equation — no need to split into halves.
- For a vertical hyperbola, swap x and y in the hyperbola formula.

TIP: Conics are cross-sections of a cone — Desmos lets you explore all four in seconds.



Distance, Midpoint, and Slope

Define draggable points with sliders — formulas update live as you move them.

```
// Define two moveable points:  
A = (a1, a2)    <- add sliders for a1 and a2  
B = (b1, b2)    <- add sliders for b1 and b2  
  
// Distance |AB|:  
d = sqrt((b1-a1)^2 + (b2-a2)^2)  
  
// Midpoint M:  
M = ((a1+b1)/2, (a2+b2)/2)  
  
// Slope of AB:  
m = (b2-a2) / (b1-a1)
```

TIP: Drag slider points A and B — watch d, M, and m update instantly. Perfect for verifying formulas.

The Desmos Geometry Tool (desmos.com/geometry)

- A separate compass-and-ruler environment — free, no account needed.
- Draw points, lines, circles, and polygons using geometric tools (not equations).
- Construct midpoints, perpendicular bisectors, angle bisectors, parallel lines.
- Measure lengths, angles, areas, and perimeters with the Measure tool.
- Reflect, rotate, translate, and dilate shapes with transformation tools.
- Add labels and annotations to create clear geometry diagrams for proofs.
- Everything is draggable — pull any point to explore how the construction changes.

TIP: Use the Geometry tool for proof-style constructions; use the Graphing Calculator when you need equations.

Calculus


Derivatives, integrals, and limits

Derivatives

- Type $f'(x)$ to plot the derivative of f — Desmos computes it automatically.
- Example: if $f(x) = x^3 - 3x$, then $f'(x)$ draws $3x^2 - 3$.
- Add $f'(x) = 0$ as a line to find critical points (where $f'(x)$ crosses x-axis).
- Local max: $f'(x)$ changes from + to - (click to read coordinates).
- Local min: $f'(x)$ changes from - to + (click to read coordinates).
- Type $f''(x)$ for the second derivative — find inflection points where $f''=0$.

TIP: Press the apostrophe key (') for prime. Higher orders: $f''(x)$, $f'''(x)$.

Derivatives — $f(x)$ and $f'(x)$ Plotted Together




Definite Integrals and Area

- Access the integral symbol via the Desmos keyboard: tap $f(x)$ -> Calculus.
- The integral shades the region AND shows the numeric value.
- Example: integral from 0 to 3 of $x^2 dx = 9$.
- Area between two curves: integrate $(f(x) - g(x))$ over the interval.
- Negative areas = the curve is below the x-axis (Desmos shows this correctly).
- Click the vertical boundary lines to drag the integration limits interactively.

TIP: Desmos shows the shaded region AND the numeric answer — use this to verify your hand calculations.

Definite Integral — Shaded Area Under the Curve



Exploring Limits Visually

- Desmos does not compute limits symbolically — but you can explore them visually.
- Method 1 — Graph: look for an open circle (hole) at a removable discontinuity.
- Method 2 — Table: set x values approaching the limit (e.g. 1.9, 1.99, 1.999, 2).
- Read the y column — the values converging tell you the limit.
- Example: $f(x) = (x^2 - 4)/(x - 2)$ has a hole at $x=2$, but the limit equals 4.
- Zoom in very close to the point to read the limit numerically from both sides.
- For one-sided limits, restrict domain: $f(x) \{x < 2\}$ and $f(x) \{x > 2\}$ separately.

TIP: If the graph has a hole at $x=a$, the limit still exists — Desmos shows it as an open circle.

Statistics

Data, regression, and distributions

Entering Data in Desmos

- Click + -> Table to create a data table in the expression list.
- Type your x values in the left column (x_1, x_2 ...) and y values on the right.
- Press Enter to add more rows. The scatter plot appears automatically.
- You can paste data copied from Excel or Google Sheets directly into the table.
- Edit cell values directly — the scatter plot updates instantly.
- Click any data point on the graph to see its exact coordinates.
- A table can have more than 2 columns — add a third for a second y-variable or formula.


TIP: Start by pasting your class data into a table — you will have a scatter plot in seconds.

Regression and Best Fit

- Use the $\sim$ (tilde) operator to fit any model to your data.
- After entering data in a table (columns x_1 , y_1):
- $y_1 \sim m \cdot x_1 + b$ linear regression
- $y_1 \sim a \cdot x_1^2 + b \cdot x_1 + c$ quadratic
- $y_1 \sim a \cdot e^{(b \cdot x_1)}$ exponential
- Desmos shows the best-fit parameters AND the R^2 value.
- R^2 closer to 1 = better fit. The fitted curve is drawn over your data.

TIP: Try multiple models (linear, quadratic, exponential) and compare their R^2 values.

Linear Regression — Best Fit Line and R^2 Value



Statistical Functions with Lists

- Define a dataset as a list: `data = [12, 15, 18, 22, 25, 30]`
- `mean(data)` — arithmetic mean | `median(data)` — middle value
- `stdev(data)` — standard deviation | `mad(data)` — mean absolute deviation
- `min(data)` and `max(data)` | `total(data)` — sum | `length(data)` — count
- `sort(data)` — ascending order | `shuffle(data)` — random order
- These functions work on any list — no need to create a table first.
- You can combine them: `mean(sort(data))` or `stdev(data)/mean(data)` (CV).


EXAMPLE

```
scores = [72,85,90,  
          66,88,91]
```

```
mean(scores) = 82  
stdev(scores) = 10.2  
median(scores) = 86.5  
min = 66  
max = 91
```

Probability Distributions

- $f(x) = \text{normaldist}(0, 1)$ → standard normal curve
- $\text{normaldist}(0,1).\text{cdf}(-1, 1) \rightarrow P(-1 \leq Z \leq 1) = 0.683$
- $\text{normaldist}(0,1).\text{inversecdf}(0.95) \rightarrow$ finds the 95th percentile
- $f(x) = \text{binomialdist}(10, 0.5) \rightarrow$ binomial $n=10, p=0.5$
- Desmos also supports: `poissondist`, `uniformdist`, and more.
- Click any point on the distribution curve to read its exact probability.



TIP: Use `.cdf(a, b)` for $P(a \leq X \leq b)$ and `.inversecdf(p)` to find the value at probability p .

Tips, Tricks & Pro Features

Get the most out of Desmos

Organizing Your Expression List

- Add a text note: start any expression line with " (quotation mark) to write a comment.
- Create folders: click + -> Folder to group related expressions.
- Click the folder arrow to collapse or expand it.
- Hide a graph without deleting it: click its colored circle to toggle visibility.
- Duplicate an expression: open Edit List mode (gear icon) and click the duplicate icon.
- Rearrange lines: in Edit List mode, drag the handle on the left to reorder.
- Delete: press the X button, or use Edit List mode for batch deletes.

TIP: Keep your list tidy — folders and text notes make complex graphs much easier to understand.

Using Desmos on iPad and Mobile

- Open desmos.com/calculator in Safari or Chrome — the full calculator works in the browser.
- The free Desmos app is available on the App Store (iOS) and Google Play (Android).
- On touchscreen: pinch to zoom, two-finger drag to pan the graph.
- Tap the keyboard icon (bottom of expression list) to open the math keyboard.
- iPad copy/paste Method 1: three-finger pinch closed = copy, pinch open = paste.
- iPad copy/paste Method 2: tap-and-hold an expression, select Copy from the menu.
- iPad copy/paste Method 3: with a physical keyboard, use Ctrl+C / Ctrl+V as usual.
- Duplicate an expression: tap Edit (gear icon) then tap the overlapping-squares icon.

TIP: The iPad app and browser version are nearly identical — all features work on both.

Pro Tips for Students

- Use Desmos as a checker — solve by hand, then verify with Desmos.
- Label important points: click any graph point and add a custom label.
- Animate sliders with the Play button — great for spotting patterns visually.
- Type π for pi, e for Euler's number, ∞ for infinity.
- Use $f(x) = \{\text{condition: value}\}$ to restrict any function to a domain.
- For scientific notation: type $1e6$ for 1,000,000.
- The Desmos keyboard has Greek letters, absolute value, summation — explore it all.
- Explore Desmos Art online — students have created stunning images with only equations!

TIP: Your teacher can also set up a Desmos Classroom activity for interactive lessons at teacher.desmos.com.

Arithmetic Operations in Desmos

- Basic operators: **+** **-** ***** **/** **^**
- Square root: type **sqrt(9)** = 3 or use the keyboard symbol
- nth root: **nthroot(8,3)** = 2 (cube root of 8)
- Pi: type **pi** → Desmos shows $\pi = 3.14159...$
- Euler's number: type **e** → Desmos uses $e = 2.71828...$
- Absolute value: **abs(-7)** = 7 or use | | from the keyboard
- Logarithms: **log(100)** = 2 | **ln(e)** = 1
- Trig: **sin(pi/6)** = 0.5 (use Wrench to set Radians or Degrees)

EXAMPLES

$$\text{sqrt}(2) * \text{pi} = 4.4429...$$

$$e^2 + \text{sqrt}(5) = 9.6207...$$

$$\text{nthroot}(27, 3) = 3$$

$$\log(1000) = 3$$

$$\sin(\text{pi}/2) = 1$$

$$2^{10} = 1024$$

$$\text{abs}(-5) + \text{sqrt}(16) = 9$$

TIP: Desmos evaluates expressions instantly. Use it as a scientific calculator — no graphing needed!

Regression: Fitting a Curve to Data

- Enter data in a **Table** with columns x_1 and y_1
- Use $\sim$ (tilde) to define your model
- Desmos shows best-fit parameters and **R^2** value (closer to 1 = better fit)

LINEAR REGRESSION (2 points)

Table: $x_1 = [1, 4]$ $y_1 = [3, 9]$

$$y_1 \sim m \cdot x_1 + b$$

Result: $m = 2$, $b = 1$ ($y = 2x + 1$)

$R^2 = 1.0$ (perfect fit)

QUADRATIC REGRESSION (3 points)

Table: $x_1 = [0, 2, 4]$

$y_1 = [1, 5, 17]$

$$y_1 \sim a \cdot x_1^2 + b \cdot x_1 + c$$

Result: $a=1$, $b=1$, $c=1$

$$y = x^2 + x + 1$$

$R^2 = 1.0$ (perfect fit)

OTHER MODELS:

$$y_1 \sim a \cdot e^{(b \cdot x_1)} \text{ exponential}$$

$$y_1 \sim a \cdot \ln(x_1) + b \text{ logarithmic}$$

TIP: Try multiple models and compare R^2 values — the model closest to 1 fits your data best.

Mean and Standard Deviation

- Define your data as a list: **data** = [v1, v2, ...]
- Mean: **mean(data)** = arithmetic average
- Standard deviation: **stdev(data)**
- Median: **median(data)** = middle value
- Min / Max: **min(data)** / **max(data)**
- Total / Count: **total(data)** / **length(data)**
- No table needed — lists work directly in the expression list

EXAMPLE

scores = [72, 85, 90, 66, 88, 91]

mean(scores) = 82

stdev(scores) = 10.2

median(scores) = 86.5

min(scores) = 66

max(scores) = 91

total(scores) = 492

length(scores) = 6

COEFFICIENT OF VARIATION:

stdev(scores)/mean(scores) = 0.124

TIP: On the SAT, mean and standard deviation are frequently tested. Verify your hand calculation instantly with Desmos.

SAT Quick-Reference: Desmos Commands

ARITHMETIC

sqrt(x) square root

nthroot(x,n) nth root

abs(x) $|x|$

pi $\pi = 3.14159...$

e $e = 2.71828...$

log(x) log base 10

ln(x) natural log

REGRESSION

Enter data in a Table

columns x_1 , y_1

then type your model:

$y_1 \sim mx_1 + b$ linear

$y_1 \sim ax_1^2 + bx_1 + c$ quadratic

$y_1 \sim ae^{(bx_1)}$ exponential

$y_1 \sim a \ln(x_1) + b$ logarithmic

Desmos shows R^2 and
best-fit curve automatically

STATISTICS

data = [v1, v2, ...]

then call:

mean(data) average

stdev(data) std dev

median(data) middle

min(data) minimum

max(data) maximum

total(data) sum

length(data) count

TIP: Keep this slide as your Desmos cheat sheet for SAT exam day — all three toolsets in one view!

You are now ready to solve problems with Desmos!

Start exploring today:

desmos.com/calculator

CLC Educational Institute | clc.gr

All rights reserved. For educational use only.