

How fluctuation-dependent species coexistence affects the diversity-stability relationship

Andrew T. Tredennick^{1,*}, Peter B. Adler¹, and Frederick R. Adler²

¹*Department of Wildland Resources and the Ecology Center, Utah State University, Logan, Utah 84322;* ²*Departments of Biology and Mathematics, University of Utah, Salt Lake City, Utah*

Abstract

For biodiversity to increase the temporal stability of ecosystem function in the long-term, coexistence must be stable. Species-specific responses to environmental variation through time is key to understanding fluctuation-dependent coexistence mechanisms and how asynchrony in species dynamics can arise to stabilize ecosystem function. Despite the shared dependence on environmental fluctuations, theory on species coexistence and the relationship between species richness and ecosystem stability have developed independently. To formally link the two bodies of theory, we use consumer-resource models where coexistence between two species utilizing a single resource is maintained by two fluctuation-dependent mechanisms: the storage effect and relative nonlinearity. We examine how the strength of species coexistence relates to the temporal stability of aggregate ecosystem function and how the effect of environmental variability on stability is mediated by the mechanism of coexistence. Blah, blah, blah...

Keywords: coexistence, storage effect, relative nonlinearity, diversity-stability hypothesis, pulsed differential equation, consumer-resource dynamics, synchrony

Introduction

Concern over the rapid rate of contemporary biodiversity loss (Barnosky et al. 2011) prompted 20 years of research on the links between biodiversity and ecosystem functioning (Cardinale et al. 2012). In particular, considerable effort has been dedicated to asking whether communities with more species are more stable through time. The question was not new, having its roots in May's classical work on complexity and stability, but it took on new relevance as local species richness declines became a worldwide phenomenon. At first, theory far outpaced empirical tests, and nearly all theoretical models suggested that the relationship between diversity and stability should be positive, even if the underlying mechanisms were debatable. Nonetheless, theory identified an important mechanism that can stabilize ecosystem functioning: species complementarity.

Empirical tests of the prediction that species richness increases ecosystem stability have relied on both experimental and observational approaches. Experimental tests involve

*Corresponding author; email: atredenn@gmail.com

manipulating the number of species present in a plot by randomly drawing a specific number of species from a pre-defined species pool (Hector et al. 1999, Tilman et al. 2001). Results from such experimental tests, which have focused on plant species richness and the temporal stability of aboveground biomass production, have been surprisingly consistent: temporal stability is greater in communities with more species (Tilman et al. 2006, Hector et al. 2010). Observational tests of the DSR, which require sampling across natural diversity gradients, have been less consistent. Positive, neutral, and negative DSRs have all been reported from observational studies. Indeed, in a meta-analysis of DSRs Jiang and Pu (2009) found no significant evidence for an effect of species richness on ecosystem stability from observational studies in terrestrial ecosystems. Why, given the consistent positive DSRs from biodiversity-ecosystem functioning experiments, do positive DSRs not consistently emerge from observational studies? To answer this question, it is necessary to reconsider the process that maintains biodiversity: species coexistence.

Stable coexistence, however maintained, is a prerequisite for biodiversity to confer stability of ecosystem function in the long term (Turnbull et al. 2013). Among the dizzying array of mechanisms that can maintain diversity, Chesson (2000) formalized two broad classes: fluctuation-independent and fluctuation-dependent mechanisms. Both classes of mechanisms rely on niche differences being greater than relative fitness differences for all species pairs in a community (Chesson 2000, Adler et al. 2007). In a fluctuation-independent case, species coexistence can be maintained by resource partitioning so long as each species is limited by a different resource (Tilman 1982). Much of the theoretical literature aimed at deciphering the mechanisms behind the diversity-stability relationship implicitly assumes fluctuation-independent coexistence (reviewed in Loreau 2010). For example, Lotka-Volterra models have been widely used and include coexistence by keeping competition coefficients less than one (e.g., Lehman and Tilman 2000, Loreau and de Mazancourt 2013). Our focus, on the other hand, is on the interaction between species coexistence and ecosystem stability in communities where coexistence is dependent on environmental fluctuations.

There are two ways that species coexist in temporally fluctuating environments. First, there is the storage effect, where species coexistence is stable if the following three conditions are met: (i) have unique responses to environmental conditions, (ii) have some way to persist in unfavorable years, and (iii) the effects of competition must be greater in ‘good’ years relative to ‘bad’ years (Chesson 2000). Second, there is relative nonlinearity, where species have unique, nonlinear responses to a shared resource that fluctuates through time (Chesson 2000).

Both the storage effect and relative nonlinearity require that species have unique responses to environmental conditions. Not only do such species-specific responses stabilize coexistence, they can also stabilize ecosystem function by creating compensatory dynamics. All else being equal, ecosystem stability is highest when species responses to the environment are perfectly uncorrelated. Under the storage effect, uncorrelated species responses to the environment also results in the most stable conditions for coexistence. It stands to reason, then, that the strength of the storage effect should predict the degree to which species asynchrony stabilizes ecosystem function. However, the exact relationship between storage effect strength and ecosystem stability could take many, unknown, forms. Species-specific responses to

non-constant environmental conditions can stabilize species coexistence (Chesson 2000) and ecosystem function (Loreau 2010). This means that fluctuation-dependent mechanisms of species coexistence are the very same mechanisms that link biodiversity and ecosystem function.

Model and Analysis

To discover how fluctuation-dependent coexistence mechanisms affect the diversity-stability relationship, we developed a semi-discrete consumer-resource model that allows many species to coexist on one resource by either the storage effect or relative nonlinearity. In our model, the consumer can be in one of two-states: a dormant state D and a live state N . The dormant state could represent, for example, the seedbank of an annual plant. Transitions between N and D occur at discrete intervals τ with continuous-time consumer-resource dynamics between discrete transitions. Thus, our model is formulated as “pulsed differential equations” (Pachepsky et al. 2008, Mailleret and Lemesle 2009, Mordecai et al. 2016). For clarity we refer to τ as years and the growing time between years as seasons with daily (t) time steps.

During a growing season, consumer-resource dynamics are modeled as two differential equations:

$$\frac{dN_i}{dt} = N_i \epsilon_i f_i(R), \quad t \neq \tau_k \quad (1)$$

$$\frac{dR}{dt} = - \sum_{i=1,2} f_i(R) N_i, \quad t \neq \tau_k \quad (2)$$

where the discrete transitions between N and D occur between seasons at times τ_k , $k = 1, 2, 3, \dots, K$. The subscript i denotes species, N is the living biomass state, and ϵ_i is each species’ resource-to-biomass conversion efficiency. The growth rate of living biomass is a resource-dependent Hill function, $f_i(R) = r_i R^{a_i} / (b_i^{a_i} + R^{a_i})$, where r is a species’ intrinsic growth rate and a and b define the curvature of the function. Resource depletion is equal to the sum of each species’ consumption.

Along with resource uptake, consumer population growth depends on the production of dormant biomass (D), the activation of dormant biomass to live biomass ($D \rightarrow N$), and the survival of living biomass from one year to the next. The biomass of each species’ states at the start of a growing season are equal to

$$D_i(\tau_k^+) = (1 - \gamma_{i,\tau_k})[\alpha_i N_i(\tau_k) + D_i(\tau_k)](1 - \eta_i) \quad (3)$$

$$N_i(\tau_k^+) = (1 - \alpha_i)N_i(\tau_k) + \gamma_{i,t}[\alpha_i N_i(\tau_k) + D_i(\tau_k)](1 - \eta_i), \quad (4)$$

where $D(\tau_k)$, $N(\tau_k)$, and $R(\tau_k)$ are the abundances of each state at the end of growing season k and τ_k^+ denotes the beginning of growing season $k = 1$. The activation of dormant biomass to live biomass is controlled by γ , which is year (k) and species (i) specific. Dormant biomass

Table 1: Default values of model parameters and their descriptions. Parameters that vary depending on the mode and strength of species coexistence or depending on species competitive hierarchies are labeled as "variable" in parentheses. The dormant-to-live biomass transition fraction (γ) is a function of other parameters, so has no default value.

Parameter	Description	Value
r	maximum per capita growth rate	1 (variable)
a	Hill function rate parameter	2 (variable)
b	Hill function curvature parameter	2.5 (variable)
ϵ	resource-to-biomass conversion efficiency	0.5
α	allocation fraction of live biomass to dormant biomass	0.5 (variable)
γ	dormant-to-live biomass transition fraction	–
ρ	correlation of species' response to the environment	0 (variable)
σ_E	variance of the environmental cue	2 (variable)
η	dormant biomass mortality rate	0.1
$\mu(R^+)$	mean annual resource pulse	20 (non-log scale)
$\sigma(R^+)$	standard deviation of annual resource pulse	0 (variable)

is equal to a constant fraction (α) of live biomass at the end of the previous season ($N_i(\tau_k)$), plus survival ($1 - \eta_i$) of dormant biomass ($D_i(\tau_k)$) at the end of the previous year and dormant biomass remaining after live biomass activation ($D_i(\tau_k)(1 - \gamma_{i,\tau_k})$). Live biomass is equal to newly activated dormant biomass ($\gamma_{i,t}[D_i(\tau_k)]$), minus some fraction of live biomass that is converted to dormant biomass ($(1 - \alpha_i)N_i(\tau_k)$). We assume the resource pool is not replenished within a growing season. Resource replenishment occurs between growing seasons, and the resource pool (R) at the start of the growing season $k+1$ is $R(\tau_k^+) = R^+$, where R^+ is a random resource pulse drawn from a log-normal distribution with mean $\mu(R^+)$ and standard deviation $\sigma(R^+)$. Model parameters and notation are described in table 1.

Implementing the Storage Effect

To make this a storage effect model, we need to satisfy three conditions: (1) the organisms must have a mechanism for persistence under unfavorable conditions, (2) species must respond differently to environmental conditions, and (3) the effects of competition on a species must be more strongly negative in good years relative to unfavorable years. Our model meets condition 1 because we include a dormant stage with very low death rates. We satisfy condition 2 with our model whenever γ is not perfectly correlated between species. Lastly, our model meets condition 3 because condition 2 partitions intraspecific and interspecific competition into different years. Thus, during a high γ year for one species, resource uptake is also inherently high for that species, which increases intraspecific competition relative to interspecific competition. So, given adequate variability in γ , the inferior competitor can persist. We created competitive hierarchies in the storage effect version of the model by altering species' biomass conversion efficiencies (ϵ)

We generated sequences of (un)correlated dormant-to-live state transition rates (γ) for each species by drawing from multivariate normal distributions with mean 0 and a variance-covariance matrix ($\Sigma(\gamma)$) of

$$\Sigma(\gamma) = \begin{bmatrix} \sigma_E^2 & \rho\sigma_E^2 \\ \rho\sigma_E^2 & \sigma_E^2 \end{bmatrix} \quad (5)$$

where σ_E^2 is the variance of the environmental cue and ρ is the correlation between the two species' transition rates. ρ must be less than 1 for stable coexistence. The inferior competitor has the strongest potential to persist when $\rho = -1$ (perfectly uncorrelated transition rates).

Implementing Relative Nonlinearity

When considering consumer-resource dynamics, species coexistence by relative nonlinearity requires that each species has different nonlinear responses to resource availability, and resource availability must fluctuate through time. In a constant resource environment, the species with the lowest R^* will always exclude the other species. To create competitive hierarchies among species we altered species resource uptake curves (Fig. SX). We still allow the germination rate (γ) to vary, but both species are perfectly correlated – that is, $\rho = 1$ (Fig. 1).




Figure 1: Resource uptake functions and example time series of (un)correlated germination fractions for the storage effect (a,b) and relative nonlinearity (c,d) formulations of the consumer-resource model. The resource uptake functions for both species are equivalent for the storage effect, but their germination fractions are uncorrelated in time. The opposite is true for relative nonlinearity: the two species have unique resource uptake functions, but their germination fractions are perfectly correlated in time.

Model Simulations

We simulated the model with four species under two scenarios for each coexistence mechanism. First, we allowed the variance of the environment to determine how many species can coexist,

akin to a community assembly experiment with a species pool of four species. This required simulating communities with all species initially present across a gradient of annual resource variability (for relative nonlinearity) or environmental cue variability (for the storage effect). Second, we chose parameter values that allowed coexistence of all four species and performed species removals. We ran simulations for 10,000 seasons with 20 day growing seasons. We averaged biomass over the growing season. After discarding an initial 1,000 seasons to reduce transient effects on our results, we calculated the coefficient of variation of summed species biomass through time. For simulations where environmental variability determined species coexistence, we calculated species richness as the number of species whose average biomass was greater than 1 over the course of the simulation. Within-season dynamics were solved given initial conditions using the package `deSolve` (Soetaert et al. 2010) in R (Team 2013). All model code has been deposited on Dryad ([link](https://github.com/atredennick/Coexistence-Stability)) and is available on GitHub at <http://github.com/atredennick/Coexistence-Stability>.

Results and Discussion

The direction of the diversity-variability relationship can be positive and negative when species coexistence is maintained by fluctuation-dependent mechanisms (Fig. 2). Ecosystem variability is positively correlated with species richness when species richness is measured across a gradient of environmental variability, which maintains diversity and promotes ecosystem variability (Fig. 2a,b). If environmental conditions are sufficient to maintain coexistence, removing species increases ecosystem variability (Fig. 2c,d). Thus, our results both confirm and contrast with theoretical and experimental findings that diversity begets stability.

When we held environmental variability constant and removed species, we produced the typical negative diversity-variability relationship (Fig. 2c,d), consistent with theoretical expectations from models with species coexistence maintained by fluctuation-dependent mechanisms. Likewise, our results from the species removal simulations are consistent with results from biodiversity-ecosystem functioning experiments showing a negative relationship between species richness and ecosystem variability. This is encouraging because species almost certainly coexist by some combination of fluctuation-independent (e.g., resource partitioning) and fluctuation-dependent mechanisms. By extending theory to communities where species richness is explicitly maintained by temporal variability, we have gained confidence that experimental findings are generalizable to many communities. In other words, in local settings where environmental variability is relatively homogenous, reductions in the number of species will reduce the stability of ecosystem functioning, regardless of how coexistence is maintained.

When we allowed a gradient of environmental variability to determine species coexistence, we discovered a positive relationship between species richness and ecosystem variability (Fig. 2a,b). While surprising when viewed through the lens of previous theory and experimental findings, such a relationship is a direct consequence of how diversity can be maintained in fluctuating environments. The storage effect and relative nonlinearity both require environmental fluctuations to allow niche differentiation between species pairs (Chesson 2000). Therefore, species coexistence gains strength, for both mechanisms, as the environment




Figure 2: Variability of total community as function of species richness when coexistence is maintained by the storage effect (a,c) or relative nonlinearity (b,d). Top panels show results from simulations where environmental or resource variance determine the number species that coexist in a community. Bottom panels show results from simulations where environmental or resource variance is fixed at a level that allows coexistence of all four species, but species are removed to manipulate diversity. In the most general sense, the top panels represent regional diversity-stability relationships across natural diversity gradients, whereas the bottom panels represent local diversity-stability relationships.

becomes more variable (Fig. SX).

Our results may explain why deviations from the negative diversity-variability relationship often come from observational studies. Observational studies must rely on natural diversity gradients, and if species richness depends environmental variability, it is entirely possible to observed positive diversity-variability relationships. For example, Sasaki and Lauenroth (2011) found a negative relationship between species richness and the temporal stability of plant abundance (a positive diversity-variability relationship) in a semi-arid grassland. Their data came from a six sites that were 6 km apart. Thus, it is possible that each site experienced slightly different levels of environmental variability that influenced species coexistence. Indeed, fluctuation-dependent coexistence mechanisms are particularly common in arid and semi-arid environments (CITATIONS).

Acknowledgments

The National Science Foundation provided funding for this work through a Postdoctoral Research Fellowship in Biology to ATT (DBI-#####) and a CAREER award to PBA (DEB-#####).

References

- Adler, P. B., J. HilleRisLambers, and J. M. Levine. 2007. A niche for neutrality. *Ecology Letters* 10:95–104.
- Barnosky, A. D., N. Matzke, S. Tomiya, G. O. U. Wogan, B. Swartz, T. B. Quental, C. Marshall, J. L. McGuire, E. L. Lindsey, K. C. Maguire, B. Mersey, and E. a Ferrer. 2011. Has the Earth’s sixth mass extinction already arrived? *Nature* 471:51–57.
- Cardinale, B. J., J. E. Duffy, A. Gonzalez, D. U. Hooper, C. Perrings, P. Venail, A. Narwani, G. M. Mace, D. Tilman, D. A. Wardle, A. P. Kinzig, G. C. Daily, M. Loreau, and J. B. Grace. 2012. Biodiversity loss and its impact on humanity. *Nature* 489:59–67.
- Chesson, P. 2000. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and Systematics* 31:343–366.
- Hector, A., Y. Hautier, P. Saner, L. Wacker, R. Bagchi, J. Joshi, M. Scherer-Lorenzen, E. M. Spehn, E. Bazeley-White, M. Weilenmann, M. C. Caldeira, P. G. Dimitrakopoulos, J. a. Finn, K. Huss-Danell, A. Jumpponen, and M. Loreau. 2010. General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology* 91:2213–2220.
- Hector, A., C. Schmid, C. Beierkuhnlein, M. C. Caldeira, M. Diemer, P. G. Dimitrakopoulos, J. a. Finn, H. Freitas, P. S. Giller, J. Good, R. Harris, P. Högberg, K. Huss-danell, J. Joshi, A. Jumpponen, C. Körner, P. W. Leadley, M. Loreau, A. Minns, C. P. H. Mulder, G. O’donovan, S. J. Otway, J. S. Pereira, A. Prinz, D. J. Read, M. Scherer-lorenzen, E. D. Schulze, A.-. Siamantziouras, E. M. Spehn, a C. Terry, a Y. Troumbis, F. I. Woodward, S. Yachi, and J. H. Lawton. 1999. Plant diversity and productivity experiments en European grasslands. *Science* 286:1123–1127.
- Jiang, L., and Z. Pu. 2009. Different effects of species diversity on temporal stability in single-trophic and multitrophic communities. *The American Naturalist* 174:651–659.
- Lehman, C. L., and D. Tilman. 2000. Biodiversity, Stability, and Productivity in Competitive Communities. *The American Naturalist* 156:534–552.
- Loreau, M. 2010. From Polutations to Ecosystems: Theoretical Fondations for a New Ecological Synthesis.
- Loreau, M., and C. de Mazancourt. 2013. Biodiversity and ecosystem stability: A synthesis of underlying mechanisms. *Ecology Letters* 16:106–115.
- Mailleret, L., and V. Lemesle. 2009. A note on semi-discrete modelling in the life sciences. *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences* 367:4779–4799.

236 Mordecai, E. A., K. Gross, and C. E. Mitchell. 2016. Within-Host Niche Differences
237 and Fitness Trade-offs Promote Coexistence of Plant Viruses. *The American Naturalist*
238 187:E13–E26.

239 Pachepsky, E., R. M. Nisbet, and W. W. Murdoch. 2008. Between discrete and continuous:
240 Consumer-resource dynamics with synchronized reproduction. *Ecology* 89:280–288.

241 Soetaert, K., T. Petzoldt, and R. W. Setzer. 2010. Package deSolve : Solving Initial Value
242 Differential Equations in R. *Journal Of Statistical Software* 33:1–25.

243 Team, R. 2013. R Development Core Team. R: A Language and Environment for Statistical
244 Computing.

245 Tilman, D. 1982. Resource competition and community structure. Pages 1–296.

246 Tilman, D., P. B. Reich, and J. M. H. Knops. 2006. Biodiversity and ecosystem stability in a
247 decade-long grassland experiment. *Nature* 441:629–632.

248 Tilman, D., P. B. Reich, J. Knops, D. Wedin, T. Mielke, and C. Lehman. 2001. Diversity
249 and productivity in a long-term grassland experiment. *Science* 294:843–845.

250 Turnbull, L. A., J. M. Levine, M. Loreau, and A. Hector. 2013. Coexistence, niches and
251 biodiversity effects on ecosystem functioning. *Ecology Letters* 16:116–127.