

1 **How fluctuation-dependent species coexistence affects the**
2 **diversity-stability relationship**

3 Andrew T. Tredennick¹, Peter B. Adler¹, and Frederick R. Adler²

4 ¹*Department of Wildland Resources and the Ecology Center, Utah State University, Logan, Utah*
5 84322

6 ²*Departments of Biology and Mathematics, University of Utah, Salt Lake City, Utah*

7 **Keywords:** coexistence, storage effect, relative nonlinearity, diversity-stability hypothesis,
8 pulsed differential equation, consumer-resource dynamics

9 **Authorship:** All authors conceived the research and designed the modeling approach; ATT
10 conducted model simulations, with input from PBA and FRA; ATT wrote the manuscript
11 and all authors contributed to revisions.

12 **Running Title:** Environmental variability, ecosystem stability, & species coexistence

13 **Article Type:** Letter

14 **Number of Words:**

15 **Number of References:**

16 **Number of Tables and Figures:**

17 **Corresponding Author:**

18 Andrew Tredennick

19 Department of Wildland Resources and the Ecology Center

20 Utah State University

21 5230 Old Main Hill

22 Logan, Utah 84322 USA

23 Phone: +1-970-443-1599

24 Fax: +1-435-797-3796

25 Email: atredenn@gmail.com

Abstract

Theory relating species richness to ecosystem stability typically ignores interactions between environmental variability and species coexistence, or fluctuation-dependent coexistence. This may explain why we lack general explanations for observed deviations from the predicted positive diversity-stability relationship. It also limits our ability to predict the consequences of increasing environmental variability. We use a consumer-resource model to explore how fluctuation-dependent coexistence via the storage effect and relative nonlinearity affects ecosystem stability. We show that a negative, rather than positive, diversity-stability relationship is possible when ecosystem function is sampled across a natural diversity gradient. We also describe how fluctuation-dependent coexistence can buffer ecosystem functioning against increasing environmental variability by allowing more species to coexist and thus contribute to portfolio effects. Our work provides a general explanation for non-positive diversity-stability relationships and highlights the importance of conserving regional species pools so that species can be added to fluctuation-dependent communities as environmental variability increases.

INTRODUCTION

MacArthur (1955), Elton (1958), and even Darwin (Turnbull et al. 2013) recognized that species can compensate for each other and stabilize functioning in ecosystems subject to temporal variation in environmental conditions. This idea underlies the “insurance hypothesis” (Yachi and Loreau 1999), which suggests stability increases with diversity because species respond dissimilarly to environmental conditions – species A has highest growth rates under conditions X whereas species B has highest growth rates under conditions Y. More species confer temporal stability by broadening the range of conditions under which the community maintains function (Loreau 2010). Diverse models all predict a positive relationship between species richness and ecosystem stability (Lehman and Tilman 2000, Ives and Hughes 2002, Loreau and de Mazancourt 2013), and experimental tests tend to support such a prediction (Tilman et al. 2006, Hector et al. 2010). However, empirical support for a positive diversity-stability relationship is not unequivocal (Jiang and Pu 2009). Furthermore, theory on the relationship between biodiversity and ecosystem stability often ignores the processes that determine species coexistence in variable environments (Loreau 2010, but see Chesson et al. 2001).

Temporally fluctuating environmental conditions are an important ingredient for stable species coexistence, both in theoretical models (Chesson 2000, Chesson et al. 2004) and in natural communities (Cáceres 1997, Descamps-Julien and Gonzalez 2005, Adler et al. 2006, Angert et al. 2009). Such “fluctuation-dependent” coexistence (Chesson 2000) requires that species have unique environmental responses and that environmental conditions vary enough for each coexisting species to experience good and bad conditions. Thus, there is reason to expect environmental variability to promote species richness when coexistence is maintained by a fluctuation-dependent mechanism (Adler and Drake 2008). Of course, increasing environmental variability may also decrease ecosystem stability through time by increasing the fluctuations of individual species, regardless of species richness.

The countervailing effects of environmental variability present an interesting paradox: increasing variability should decrease ecosystem stability, but may also increase richness, which may offset the decrease in stability. Such a paradox complicates predictions about how ecosystems will respond as environmental conditions exceed historical ranges of variability. The unknown net effect of environment variability may be reflected in the mixed results from empirical studies on the diversity-stability relationship. Observational tests of the diversity-stability relationship, which require sampling across natural diversity gradients, have yielded positive (Hautier et al. 2014), neutral (Valone and Hoffman 2003, Cusson et al. 2015), and negative (Sasaki and Lauenroth 2011) relationships. In a meta-analysis of diversity-stability relationships, Jiang and Pu (2009) found no significant evidence for an effect of species richness on ecosystem stability from observational studies in terrestrial ecosystems. Thus, there appears to be a gap between the consistency of theoretical studies and the equivocation of empirical studies.

We argue this gap exists because the two bodies of theory that have developed to explain species coexistence on the one hand, and diversity-stability relationships on the other, have diverged. One reason these two disciplines have diverged is because they have focused on slightly different questions. Biodiversity-ecosystem stability studies typically ask how ecosystem variability responds to different levels of species richness at a given level of environmental variability (reviewed in Kinzig et al. 2001, Loreau 2010), whereas coexistence studies ask how the long term stability of species coexistence responds to different levels of environmental variability (Chesson and Warner 1981).

To reconcile these two bodies of theory, we extend theory on the relationship between species richness and ecosystem stability to cases in which species coexistence explicitly depends on environmental fluctuations and species-specific responses to environmental conditions. We focus on the storage effect and relative nonlinearity using a general consumer-resource model. First, we use model simulations to investigate the diversity-stability relationship across a gradient of environmental variability. Counter to common expectations, we find

that a negative diversity-stability relationship should be expected when sampling occurs over natural diversity gradients and species coexistence is fluctuation dependent. Importantly, and in line with previous theory (Ives and Hughes 2002, Loreau 2010, Mazancourt et al. 2013), at a given level of environmental variability, increasing species richness stabilizes ecosystem function, even when coexistence is fluctuation dependent.

Second, we explore the net effect of increasing environmental variability on ecosystem stability by isolating the gain in stability due to increased richness and the loss in stability due to increased amplitude of species fluctuations. We find that increasing environmental variability does not always lead to a decrease in ecosystem stability: environmental variability promotes species richness, which stabilizes ecosystem functioning via portfolio effects even as the environment becomes more variable. Whether or not increasing environmental variability results in species gains depends on the specific coexistence mechanism, the traits of resident species, and the traits of the species in the regional pool.

MATERIALS AND METHODS

Consumer-resource model

We developed a semi-discrete consumer-resource model that allows many species to coexist on one resource by either the storage effect or relative nonlinearity. In our model, the consumer can be in one of two-states: a dormant state D and a live state N . The dormant state could represent, for example, the seedbank of an annual plant. Transitions between N and D occur at discrete intervals τ with continuous-time consumer-resource dynamics between discrete transitions. Thus, our model is formulated as “pulsed differential equations” (Pachepsky et al. 2008, Mailleret and Lemesle 2009, Mordecai et al. 2016). For clarity we refer to τ as years and the growing time between years as seasons with daily (t) time steps.

During a growing season, consumer-resource dynamics are modeled as two differential equations:

$$\frac{dN_i}{dt} = N_i \epsilon_i f_i(R), \quad t \neq \tau_k \quad (1)$$

$$\frac{dR}{dt} = - \sum_{i=1,2} f_i(R) N_i, \quad t \neq \tau_k \quad (2)$$

where the discrete transitions between N and D occur between seasons at times τ_k , $k = 1, 2, 3, \dots, K$. The subscript i denotes species, N is the living biomass state, and ϵ_i is each species' resource-to-biomass conversion efficiency. The growth rate of living biomass is a resource-dependent Hill function, $f_i(R) = r_i R^{a_i} / (b_i^{a_i} + R^{a_i})$, where r is a species' intrinsic growth rate and a and b define the curvature of the function. Resource depletion is equal to the sum of each species' consumption.

Along with resource uptake, consumer population growth depends on the production of dormant biomass (D), the activation of dormant biomass to live biomass ($D \rightarrow N$), and the survival of living biomass from one year to the next. The biomass of each species' states at the start of a growing season are equal to

$$D_i(\tau_k^+) = (1 - \gamma_{i,\tau_k})[\alpha_i N_i(\tau_k) + D_i(\tau_k)](1 - \eta_i) \quad (3)$$

$$N_i(\tau_k^+) = (1 - \alpha_i)N_i(\tau_k) + \gamma_{i,t}[\alpha_i N_i(\tau_k) + D_i(\tau_k)](1 - \eta_i), \quad (4)$$

where $D(\tau_k)$, $N(\tau_k)$, and $R(\tau_k)$ are the abundances of each state at the end of growing season k and τ_k^+ denotes the beginning of growing season $k = 1$. The activation of dormant biomass to live biomass is controlled by γ , which is year (k) and species (i) specific. Dormant biomass is equal to a constant fraction (α) of live biomass at the end of the previous season ($N_i(\tau_k)$), plus survival ($1 - \eta_i$) of dormant biomass ($D_i(\tau_k)$) at the end of the previous year and dormant biomass remaining after live biomass activation ($D_i(\tau_k)(1 - \gamma_{i,\tau_k})$). Live biomass is equal to newly activated dormant biomass ($\gamma_{i,t}[D_i(\tau_k)]$), minus some fraction of live biomass that is

converted to dormant biomass $((1 - \alpha_i)N_i(\tau_k))$. We assume the resource pool is not replenished within a growing season. Resource replenishment occurs between growing seasons, and the resource pool (R) at the start of the growing season $k+1$ is $R(\tau_k^+) = R^+$, where R^+ is a random resource pulse drawn from a log-normal distribution with mean $\mu(R^+)$ and standard deviation $\sigma(R^+)$. Model parameters and notation are described in Table 1.

Implementing the Storage Effect To make this a storage effect model, we need to satisfy three conditions: (1) the organisms must have a mechanism for persistence under unfavorable conditions, (2) species must respond differently to environmental conditions, and (3) the effects of competition on a species must be more strongly negative in good years relative to unfavorable years. Our model meets condition 1 because we include a dormant stage with very low death rates. We satisfy condition 2 with our model whenever γ is not perfectly correlated between species. Lastly, our model meets condition 3 because condition 2 partitions intraspecific and interspecific competition into different years. Thus, during a high γ year for one species, resource uptake is also inherently high for that species, which increases intraspecific competition relative to interspecific competition. So, given adequate variability in γ , the inferior competitor can persist. We created competitive hierarchies in the storage effect version of the model by altering species' biomass conversion efficiencies (ϵ)

We generated sequences of (un)correlated dormant-to-live state transition rates (γ) for each species by drawing from multivariate normal distributions with mean 0 and a variance-covariance matrix ($\Sigma(\gamma)$) of

$$\Sigma(\gamma) = \begin{bmatrix} \sigma_E^2 & \rho_{1,2}\sigma_E^2 & \rho_{1,3}\sigma_E^2 & \rho_{1,4}\sigma_E^2 \\ \rho_{2,1}\sigma_E^2 & \sigma_E^2 & \rho_{2,3}\sigma_E^2 & \rho_{2,4}\sigma_E^2 \\ \rho_{3,1}\sigma_E^2 & \rho_{3,2}\sigma_E^2 & \sigma_E^2 & \rho_{3,4}\sigma_E^2 \\ \rho_{4,1}\sigma_E^2 & \rho_{4,2}\sigma_E^2 & \rho_{4,3}\sigma_E^2 & \sigma_E^2 \end{bmatrix} \quad (5)$$

where σ_E^2 is the variance of the environmental cue and $\rho_{i,j}$ is the correlation between between the species i 's and species j 's transition rates. ρ must be less than 1 for stable coexistence, and in all simulations we placed that constraint that $\rho_{i,j} = \rho_{j,i}$ for each species pair. The inferior competitor has the strongest potential to persist when $\rho = -1$ (perfectly uncorrelated transition rates). We used the R function `mvrnorm` to generate sequences of (un)correlated variates E that we converted to germination rates in the 0-1 range: $\gamma = e^E / (1 + e^E)$. Note that $\Sigma(\gamma)$ must be positive definite. So, after defining $\Sigma(\gamma)$ with all $\rho_{i,j}$ s and σ_E^2 , we used the `nearPD` function from the `Matrix` package in R to coerce the variance-covariance matrix to be positive definite.

Implementing Relative Nonlinearity When considering consumer-resource dynamics, species coexistence by relative nonlinearity requires that each species has different nonlinear responses to resource availability, and resource availability must fluctuate through time. In a constant resource environment, the species with the lowest R^* will always exclude the other species. To create competitive hierarchies among species we manipulated species resource uptake curves such that the species with the lowest R^* also had the lowest maximum growth rate at high resource levels (e.g., low r , low a , and low b values relative to other species in the Hill equation; Fig. SX). Thus, our simulated species represent a continuum from resource-conservative to resource-acquisitive. We still allow the germination rate (γ) to vary, but both species are perfectly correlated – that is, $\rho = 1$ (Fig. 1).

Numerical simulations

To understand how fluctuation-dependent coexist can affect the diversity-stability relationship, we simulated the model with four species under two scenarios for each coexistence mechanism. First, we allowed the variance of the environment to determine how many species can coexist, akin to a community assembly experiment with a species pool of four species. This required simulating communities with all species initially present across a gradient of annual resource

variability (for relative nonlinearity) or environmental cue variability (for the storage effect). Second, we chose parameter values that allowed coexistence of all four species and performed species removals. The two simulation experiments correspond to (i) sampling ecosystem function across a natural gradient of species richness and (ii) sampling ecosystem function across diversity treatments within a site.

To understand how increasing environmental variability will impact ecosystem stability when coexistence is fluctuation-dependent, we simulated the model over a range of environmental cue variability and species pool sizes. Thus, for each size of species pool (1, 2, 3, or 4 species), we simulated the model at 15 evenly-spaced levels of environmental cue (range = 0.1,2) for the storage effect and 25 evenly-spaced levels of resource variability (range = 0.1,1.4) for relative nonlinearity. We also explored the influence of species asymmetries in competitive ability and species' correlations of environmental responses in the storage effect model. Under the storage effect, if all species are perfectly symmetrical, that is, there is no superior competitor, then coexistence is fluctuation independent. We use one such parameterization of our model to contrast the response of ecosystem stability in fluctuation-dependent and fluctation-independent communities to environmental variation. Likewise, under relative nonlinearity, species' resource response curves (Fig. SX) reflect traits that determine the intrinsic stability of each species. Therefore, we ran two sets of simulations for relative nonlinearity: one where the species pools increased from stable to unstable species and vice versa. For example, if species A is the most stable species and species D is the least stable we ran simulations where the species pool increased from one to four species as A then B then C then D. We then ran simulations with that order reversed.

All simulations were run for 5,000 seasons with 20-day growing seasons. We averaged biomass over the growing season, and those yearly values were used to calculate total community biomass in each year. After discarding an initial 500 seasons to reduce transient effects on our results, we calculated the coefficient of variation (CV) of summed species biomass through time. Therefore, in our results we refer to ecosystem variability, which is

the inverse of ecosystem stability. We calculated species richness as the number of species whose average biomass was greater than 1 over the course of the simulation. Parameter values for specific results are given in figure captions. Within-season dynamics were solved given initial conditions using the package `deSolve` (Soetaert et al. 2010) in R (Team 2013). All model code has been deposited on Figshare (*link*) and is available on GitHub at <http://github.com/atredennick/Coexistence-Stability>.

RESULTS AND DISCUSSION

The diversity-variability relationship

The direction of the diversity-variability relationship can be positive and negative when species coexistence is maintained by fluctuation-dependent mechanisms (Fig. 2). Ecosystem variability is positively correlated with species richness when species richness is measured across a gradient of environmental variability, which maintains diversity and promotes ecosystem variability (Fig. 2A,C). If environmental conditions are sufficient to maintain coexistence, removing species increases ecosystem variability (Fig. 2B,D). Thus, our results both confirm and contrast with theoretical and experimental findings that diversity begets stability.

When we held environmental variability constant and removed species, we produced the typical negative diversity-variability relationship (Fig. 2B,D), consistent with theoretical expectations from models with species coexistence maintained by fluctuation-dependent mechanisms. Likewise, our results from the species removal simulations are consistent with results from biodiversity-ecosystem functioning experiments showing a negative relationship between species richness and ecosystem variability. This is encouraging because species almost certainly coexist by some combination of fluctuation-independent (e.g., resource partitioning) and fluctuation-dependent mechanisms. By extending theory to communities where species richness is explicitly maintained by temporal variability, we have gained confidence that

experimental findings are generalizable to many communities. In other words, in local settings where environmental variability is relatively homogenous, reductions in the number of species will reduce the stability of ecosystem functioning, regardless of how coexistence is maintained.

When we allowed a gradient of environmental variability to determine species coexistence, we discovered a positive relationship between species richness and ecosystem variability (Fig. 2A,C). While surprising when viewed through the lens of previous theory and experimental findings, such a relationship is a direct consequence of how diversity can be maintained in fluctuating environments. The storage effect and relative nonlinearity both require environmental fluctuations to allow niche differentiation between species pairs (Chesson 2000). Therefore, species coexistence gains strength, for both mechanisms, as the environment becomes more variable (Fig. SX).

Our results may explain why deviations from the negative diversity-variability relationship often come from observational studies (Jiang and Pu 2009). Observational studies must rely on natural diversity gradients, and if species richness depends on environmental variability, it is entirely possible to observe positive diversity-variability relationships. For example, Sasaki and Lauenroth (2011) found a negative relationship between species richness and the temporal stability of plant abundance (a positive diversity-variability relationship) in a semi-arid grassland. Their data came from six sites that were 6 km apart. While Sasaki and Lauenroth explained their results in terms of dominant species' effects, it is also possible that each site experienced slightly different levels of environmental variability that influenced species coexistence. DeClerck et al. (2006) also found a positive diversity-variability when sampling conifer richness and the variability of productivity across a large spatial gradient in the Sierra Nevada.

While our modeling results show that fluctuation-dependent coexistence can create positive diversity-variability relationships, whether such trends are detected will depend on the particular traits of the species in the community and the relative influence of fluctuation-

dependent and fluctuation-independent coexistence mechanisms, which are not mutually exclusive. Thus, our results may also help explain observational studies where no relationship between diversity and variability is detected. For example, Cusson et al. (2015) found no relationship between species richness and variability of abundances in several marine macro-benthic ecosystems. Many of their focal ecosystems were from highly variable intertidal environments. If coexistence was at least in part determined by environmental fluctuations, then the confounding effect of variability and species richness could compensate any direct effect of species richness on variability. Previous theoretical work showed how environmental variation can mask the effect of species diversity on ecosystem productivity when sampling across sites (Loreau 1998). Our mechanistic model extends that conclusion to ecosystem stability.

The impact of increasing environmental variability on ecosystem variability

Whether coexistence is fluctuation-independent or fluctuation-dependent becomes especially important when we consider how ecosystem stability responds to increasing environmental variability. In the fluctuation-independent case, species richness is essentially fixed because the species' inequalities that determine coexistence (niche and fitness differences) are not linked to environmental variability. Therefore, increasing environmental variability will always increase ecosystem variability by increasing the fluctuations of individual species' abundances.

When species coexistence is fluctuation-dependent via the storage effect, increasing environmental variability has much more interesting effects on ecosystem variability. In Fig. 3 we show storage effect simulation results where environmental variability determines species coexistence from a regional species pool of four species. We also show results from nested subsets of the four species pool (e.g., only two species in the pool instead of four) to show the trajectory of ecosystem variability if new species are not present to join the local community. In accordance with coexistence theory, we find that realized species richness increases with

environmental variability and, in some cases, increasing variability can actually completely temper the effect of increasing environmental variability. More species rich communities are less variable on average (e.g., lower intercepts in log-log space; Fig. SX) and increase in ecosystem CV at a slower rate (e.g., lower slopes in log-log space; Fig. SX).

The dampening effect of fluctuation-dependent coexistence on increasing environmental variability depends on the specific traits (parameter values) of the species in the regional pool. Asymmetric competition makes it more difficult for new species to enter the local community (Fig. 3; compare top and bottom panels). Relatively high competition also increases the rate at which ecosystem CV increases with environmental variance (Fig. SX). This is because the abundance of inferior competitors is reduced and therefore does not influence ecosystem CV as much as when competition is symmetric. The correlation of species' environmental responses also mediates the relationship between environmental variance, species richness, and ecosystem CV : lower correlations make it easier for new species to coexist and unique environmental responses are always stabilizing (Fig. 3).

In communities where species coexist via relative nonlinearity, whether or not the direct impact of environmental variability on ecosystem variability is tempered by species additions depends on the species traits of immigrating species. When additional species, which immigrate from the regional pool, are less intrinsically stable than the resident species, ecosystem variability increases at a constant rate even as species are added (Fig. 4A). On the contrary, if more stable species are added, species additions buffer the ecosystem from increasing environmental variability (Fig. 4B). The stability of individual species in our relative nonlinearity model is determined by their respective resource response curves (Fig. SX). Under relative nonlinearity, we find that the buffering effect of species additions depends on species traits, and the order in which species enter the local community. Indeed, if all species in the regional pool are less stable than the resident species, then no stabilization occurs as species are added (Fig. 4A).

Our simulation results lead to two conclusions. First, when predicting the impacts of increasing environmental variability on ecosystem stability, the mechanism of coexistence in the community matters. Fluctuation-dependent coexistence can buffer ecosystems from increasing environmental variability by allowing for species additions. As shown in previous work (Loreau and de Mazancourt 2013), the stabilizing effect of species additions depends on the correlations of their environmental responses (Fig. 3e-f). Whether our theoretical predictions hold in real communities is unknown and requires empirical tests. Doing so would require manipulating environmental variability in communities where coexistence is known to be fluctuation-dependent, at least in part. Such data do exist (Angert et al. 2009), and a coupled modeling-experimental approach could determine if our predictions hold true in real communities.

Second, whether local fluctuation-dependent communities can receive the benefit of additional species depends on a diverse regional species pool. If the regional pool is not greater in size than the local species pool, than ecosystem stability will decline with environmental variability in a similar manner as in fluctuation-independent communities because species richness will be fixed (Fig. 5A,B). Metacommunity theory has made clear the importance of rescue effects to avoid species extinctions (Brown and Kodric-Brown 1997, Leibold et al. 2004). Here, instead of local immigration by a resident species working to rescue a species from extinction, immigration to the local community by a new species rescues ecosystem processes from becoming less stable (Fig. 5C,D). Thus, our results reinforce the importance of both local and regional biodiversity conservation. Just as declines in local species richness can destabilize ecosystem functioning (Tilman et al. 2006, Hector et al. 2010, Hautier et al. 2014), species losses at larger spatial scales can also weaken stability. Wang and Loreau (2014) show that regional ecosystem stability depends on regional biodiversity through its effects on beta diversity and, in turn, the asynchrony of functioning in local communities. Our results show that, when coexistence is fluctuation-dependent, regional biodiversity declines could also affect local ecosystem functioning by limiting local species additions that could be

possible under scenarios of increasing environmental variability (Fig. 5).

Species coexistence in real ecological communities probably emerges from some combination of fluctuation-independent and fluctuation-dependent mechanisms (Chesson 2000, Clark et al. 2010). Likewise, environmental conditions in real ecosystems are unlikely to change only in their variability without an associated change in the mean (Avolio et al. 2015). Therefore, environmental change has the potential to alter the niche and fitness differences among species in multiple ways, some of which were not present in our current analysis. Mean changes in environmental conditions could reorder competitive hierarchies (Klanderud and Totland 2005) and/or alter the availability of niches (Harpole et al. 2016). Associated changes in ecosystem stability will depend upon the magnitude of environmental change, each species' response to the particular environmental driver, and biotic interactions (Hallett et al. 2014). Thus, it is becoming clear that understanding how ecosystem stability will respond to global change will require a trait-based approach.

CONCLUSIONS

How does fluctuation-dependent coexistence affect the diversity-stability relationship? At a given level of environmental variability, the typical positive diversity-stability relationship holds because having more species always stabilizes ecosystem functioning. However, counter to other theoretical studies, we found that a negative diversity-stability relationship is also possible if sampling occurs across a natural diversity gradient and species coexistence is dependent on environmental fluctuations. We also found that fluctuation-dependent species coexistence may help buffer ecosystems from increasing environmental variability because environmental variability promotes species richness, which, in turn, promotes stability. Where fluctuation-dependent species coexistence prevails and environmental variability is projected to increase, our findings suggest that conserving regional species pools and dispersal corridors between local communities will be important.

ACKNOWLEDGMENTS

The National Science Foundation provided funding for this work through a Postdoctoral Research Fellowship in Biology to ATT (DBI-1400370) and a CAREER award to PBA (DEB-1054040).

TABLE

Table 1 Default values of model parameters and their descriptions. Parameters that vary depending on the mode and strength of species coexistence or depending on species competitive hierarchies are labeled as "variable" in parentheses. The dormant-to-live biomass transition fraction (γ) is a function of other parameters, so has no default value.

Parameter	Description	Value
r	maximum per capita growth rate	1 (variable)
a	Hill function rate parameter	2 (variable)
b	Hill function curvature parameter	2.5 (variable)
ϵ	resource-to-biomass conversion efficiency	0.5
α	allocation fraction of live biomass to dormant biomass	0.5 (variable)
γ	dormant-to-live biomass transition fraction	–
ρ	correlation of species' response to the environment	0 (variable)
σ_E	variance of the environmental cue	2 (variable)
η	dormant biomass mortality rate	0.1
$\mu(R^+)$	mean annual resource pulse	20 (non-log scale)
$\sigma(R^+)$	standard deviation of annual resource pulse	0 (variable)

FIGURES




Figure 1 Resource uptake functions and example time series of (un)correlated germination fractions for the storage effect (A,B) and relative nonlinearity (C,D) formulations of the consumer-resource model. The resource uptake functions for both species are equivalent for the storage effect, but their germination fractions are uncorrelated in time. The opposite is true for relative nonlinearity: the two species have unique resource uptake functions, but their germination fractions are perfectly correlated in time.




Figure 2 Variability of total community as function of species richness when coexistence is maintained by the storage effect (A,B) or relative nonlinearity (C,D). Top panels show results from simulations where environmental or resource variance determine the number species that coexist in a community. Bottom panels show results from simulations where environmental or resource variance is fixed at a level that allows coexistence of all four species, but species are removed to manipulate diversity. The top panels represent regional diversity-stability relationships across natural diversity gradients, whereas the bottom panels represent local diversity-stability relationships.




Figure 3 The effect of environmental variability on ecosystem variability with associated effects of species richness when species coexist via the storage effect. Panels (A-C) show simulation results where species have slightly asymmetrical competitive effects, whereas panels (D-F) show results when competition is more asymmetric. We show results for different levels of correlations of species' environmental responses, ρ .




Figure 4 The effect of environmental variability on ecosystem variability with associated effects of species richness when species coexist via relative nonlinearity. (A) The species pool increases from 1-4 four species, with the fourth species being most unstable. Increasing environmental variability (the SD of annual resource availability) allows for greater species richness, but species additions do not modulate the effect of environmental variability on ecosystem variability. (B) The species pool increases from 1-4 four species, with the fourth species being most stable (though, the fourth species was unable to coexist under these parameter values). In this case increasing environmental variability allows for greater realized species richness and can temper the effect of environmental variability.




Figure 5 Example of how species additions under increasing environmental variability can buffer ecosystem stability when species coexistence is fluctuation-dependent via the storage effect. Environmental variability (σ_E^2) increases linearly with time. (A) Time series of species' biomasses (colored lines) in a closed community where colonization of new species is not possible and (B) its associated coefficient of variation (Rolling CV; calculated over 100-yr moving window) through time. (C) Time series of species' biomasses in an open community where colonization by new species from the regional pool of four species becomes possible as environmental variation increases. The trajectory of total biomass CV in the open community (D) asymptotes at lower variability than in the closed community (B) due to the buffering effect of species richness.

REFERENCES

- Adler, P. B., and J. M. Drake. 2008. Environmental variation, stochastic extinction, and competitive coexistence. *The American Naturalist* 172:186–195.
- Adler, P. B., J. HilleRisLambers, P. C. Kyriakidis, Q. Guan, and J. M. Levine. 2006. Climate variability has a stabilizing effect on the coexistence of prairie grasses. *Proceedings of the National Academy of Sciences* 103:12793–12798.
- Angert, A. L., T. E. Huxman, P. Chesson, and D. L. Venable. 2009. Functional tradeoffs determine species coexistence via the storage effect. *Proceedings of the National Academy of Sciences of the United States of America* 106:11641–11645.
- Avolio, M. L., K. J. L. Pierre, G. R. Houseman, S. E. Koerner, E. Grman, F. Isbell, D. S. Johnson, and K. R. Wilcox. 2015. A framework for quantifying the magnitude and variability of community responses to global change drivers. *Ecosphere* 6:1–14.
- Brown, J. H., and A. Kodric-Brown. 1997. Turnover Rates in Insular Biogeography : Effect of Immigration on Extinction. *Ecology* 58:445–449.
- Cáceres, C. E. 1997. Temporal variation, dormancy, and coexistence: a field test of the storage effect. *Proceedings of the National Academy of Sciences* 94:9171–9175.
- Chesson, P. 2000. Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and Systematics* 31:343–366.
- Chesson, P. L., and R. R. Warner. 1981. Environmental Variability Promotes Coexistence in Lottery Competitive Systems. *The American Naturalist* 117:923–943.
- Chesson, P., R. L. E. Gebauer, S. Schwinning, N. Huntly, K. Wiegand, M. S. K. Ernest, A. Sher, A. Novoplansky, and J. F. Weltzin. 2004. Resource pulses, species interactions, and diversity maintenance in arid and semi-arid environments. *Oecologia* 141:236–253.
- Chesson, P., S. W. Pacala, and C. Neuhauser. 2001. Environmental Niches and Ecosystem Functioning. Pages 213–245 in A. P. Kinzig, S. W. Pacala, and D. Tilman, editors. *The functional consequences of biodiversity: Empirical progress and theoretical extensions*. Princeton University Press, Princeton.
- Clark, J. S., D. Bell, C. Chu, B. Courbaud, M. Dietze, M. Hersh, J. HilleRisLambers, I. Ibáñez, S. LaDeau, S. McMahon, J. Metcalf, J. Mohan, E. Moran, L. Pangle, S. Pearson, C. Salk, Z. Shen, D. Valle, and P. Wyckoff. 2010. High-dimensional coexistence based on individual variation: a synthesis of evidence. *Ecological Monographs* 80:569–608.
- Cusson, M., T. P. Crowe, R. Araújo, F. Arenas, R. Aspden, F. Bulleri, D. Davoult, K. Dyson, S. Frascchetti, K. Herkül, C. Hubas, S. Jenkins, J. Kotta, P. Kraufvelin, A. Migné, M. Molis, O. Mulholland, L. M.-L. Noël, D. M. Paterson, J. Saunders, P. J. Somerfield, I. Sousa-Pinto, N. Spilmont, A. Terlizzi, and L. Benedetti-Cecchi. 2015. Relationships between biodiversity and the stability of marine ecosystems: Comparisons at a European scale using meta-analysis. *Journal of Sea Research* 98:5–14.
- DeClerck, F. A. J., M. G. Barbour, and J. O. Sawyer. 2006. Species richness and stand stability in conifer forests of the Sierra Nevada. *Ecology* 87:2787–2799.

- Descamps-Julien, B., and A. Gonzalez. 2005. Stable coexistence in a fluctuating environment: An experimental demonstration. *Ecology* 86:2815–2824.
- Elton, C. 1958. *The Ecology of Invasions by Animals and Plants*. Pages 1689–1699. University of Chicago Press, Chicago.
- Hallett, L. M., J. S. Hsu, E. E. Cleland, S. L. Collins, T. L. Dickson, E. C. Farrer, L. A. Gherardi, K. L. Gross, R. J. Hobbs, L. Turnbull, and K. N. Suding. 2014. Biotic mechanisms of community stability shift along a precipitation gradient. *Ecology* 95:1693–1700.
- Harpole, W. S., L. L. Sullivan, E. M. Lind, J. Firn, P. B. Adler, E. T. Borer, J. Chase, P. A. Fay, Y. Hautier, H. Hillebrand, A. S. MacDougall, E. W. Seabloom, R. Williams, J. D. Bakker, M. W. Cadotte, E. J. Chaneton, C. Chu, E. E. Cleland, C. D’Antonio, K. F. Davies, D. S. Gruner, N. Hagenah, K. Kirkman, J. M. H. Knops, K. J. La Pierre, R. L. McCulley, J. L. Moore, J. W. Morgan, S. M. Prober, A. C. Risch, M. Schuetz, C. J. Stevens, and P. D. Wragg. 2016. Addition of multiple limiting resources reduces grassland diversity. *Nature* 537:93–96.
- Hautier, Y., E. W. Seabloom, E. T. Borer, P. B. Adler, W. S. Harpole, H. Hillebrand, E. M. Lind, A. S. MacDougall, C. J. Stevens, J. D. Bakker, Y. M. Buckley, C. Chu, S. L. Collins, P. Daleo, E. I. Damschen, K. F. Davies, P. a Fay, J. Firn, D. S. Gruner, V. L. Jin, J. a Klein, J. M. H. Knops, K. J. La Pierre, W. Li, R. L. McCulley, B. a Melbourne, J. L. Moore, L. R. O’Halloran, S. M. Prober, A. C. Risch, M. Sankaran, M. Schuetz, and A. Hector. 2014. Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature* 508:521–5.
- Hector, A., Y. Hautier, P. Saner, L. Wacker, R. Bagchi, J. Joshi, M. Scherer-Lorenzen, E. M. Spehn, E. Bazeley-White, M. Weilenmann, M. C. Caldeira, P. G. Dimitrakopoulos, J. a. Finn, K. Huss-Danell, A. Jumpponen, and M. Loreau. 2010. General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology* 91:2213–2220.
- Ives, A. R., and J. B. Hughes. 2002. General relationships between species diversity and stability in competitive systems. *The American naturalist* 159:388–395.
- Jiang, L., and Z. Pu. 2009. Different effects of species diversity on temporal stability in single-trophic and multitrophic communities. *The American Naturalist* 174:651–659.
- Kinzig, A. P., S. W. Pacala, and D. Tilman (Eds.). 2001. *The functional consequences of biodiversity: Empirical progress and theoretical extensions*. Pages i–365. Princeton University Press, Princeton.
- Klanderud, K., and Ø. Totland. 2005. Simulated climate change altered dominance hierarchies and diversity of an alpine biodiversity hotspot. *Ecology* 86:2047–2054.
- Lehman, C. L., and D. Tilman. 2000. Biodiversity, Stability, and Productivity in Competitive Communities. *The American Naturalist* 156:534–552.
- Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D. Holt, J. B. Shurin, R. Law, D. Tilman, M. Loreau, and A. Gonzalez. 2004. The metacommunity concept: A framework for multi-scale community ecology.

- Loreau, M. 1998. Biodiversity and ecosystem functioning: a mechanistic model. *Proceedings of the National Academy of Sciences of the United States of America* 95:5632–5636.
- Loreau, M. 2010. From Polutations to Ecosystems: Theoretical Fondations for a New Ecological Synthesis.
- Loreau, M., and C. de Mazancourt. 2013. Biodiversity and ecosystem stability: A synthesis of underlying mechanisms. *Ecology Letters* 16:106–115.
- MacArthur, R. 1955. Fluctuations of Animal Populations and a Measure of Community Stability. *Ecology* 36:533–536.
- Mailleret, L., and V. Lemesle. 2009. A note on semi-discrete modelling in the life sciences. *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences* 367:4779–4799.
- Mazancourt, C. de, F. Isbell, A. Larocque, F. Berendse, E. De Luca, J. B. Grace, B. Haegeman, H. Wayne Polley, C. Roscher, B. Schmid, D. Tilman, J. van Ruijven, A. Weigelt, B. J. Wilsey, and M. Loreau. 2013. Predicting ecosystem stability from community composition and biodiversity. *Ecology Letters* 16:617–625.
- Mordecai, E. A., K. Gross, and C. E. Mitchell. 2016. Within-Host Niche Differences and Fitness Trade-offs Promote Coexistence of Plant Viruses. *The American Naturalist* 187:E13–E26.
- Pachepsky, E., R. M. Nisbet, and W. W. Murdoch. 2008. Between discrete and continuous: Consumer-resource dynamics with synchronized reproduction. *Ecology* 89:280–288.
- Sasaki, T., and W. K. Lauenroth. 2011. Dominant species, rather than diversity, regulates temporal stability of plant communities. *Oecologia* 166:761–768.
- Soetaert, K., T. Petzoldt, and R. W. Setzer. 2010. Package deSolve : Solving Initial Value Differential Equations in R. *Journal Of Statistical Software* 33:1–25.
- Team, R. 2013. R Development Core Team. R: A Language and Environment for Statistical Computing.
- Tilman, D., P. B. Reich, and J. M. H. Knops. 2006. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* 441:629–632.
- Turnbull, L. A., J. M. Levine, M. Loreau, and A. Hector. 2013. Coexistence, niches and biodiversity effects on ecosystem functioning. *Ecology Letters* 16:116–127.
- Valone, T. J., and C. D. Hoffman. 2003. A mechanistic examination of diversity-stability relationships in annual plant communities. *Oikos* 103:519–527.
- Wang, S., and M. Loreau. 2014. Ecosystem stability in space: α , β and γ variability. *Ecology Letters* 17:891–901.
- Yachi, S., and M. Loreau. 1999. Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proceedings of the National Academy of Sciences of the United States of America* 96:1463–1468.