

# Ecosystem functional response across precipitation extremes in a sagebrush steppe

Andrew T. Tredennick<sup>1,\*</sup>, Andrew R. Kleinbesselink<sup>1,2,\*</sup>, J. Bret Taylor<sup>3</sup>, and Peter B. Adler<sup>1</sup>

<sup>1</sup>Department of Wildland Resources and the Ecology Center, Utah State University, Logan, Utah 84322

<sup>2</sup>Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los Angeles, California 90095

<sup>3</sup>United States Department of Agriculture, Agricultural Research Service, U.S. Sheep Experiment Station, Dubois, Idaho 83423

Corresponding author:

Andrew T. Tredennick<sup>1,\*</sup>

Email address: atredenn@gmail.com

## ABSTRACT

**Background.** Precipitation is predicted to become more variable in the western United States, meaning years of above and below average precipitation will become more common. Periods of extreme precipitation are major drivers of interannual variability in ecosystem functioning in water limited communities, but how ecosystems respond to these extremes over the long-term may shift with precipitation means and variances. Long-term changes in ecosystem functional response could reflect compensatory changes in species composition or species reaching physiological thresholds at extreme precipitation levels.

**Methods.** We conducted a five year precipitation manipulation experiment in a sagebrush steppe ecosystem in Idaho, United States. We used drought and irrigation treatments (approximately 50% decrease/increase) to investigate whether ecosystem functional response remains consistent under sustained high or low precipitation. We recorded data on aboveground net primary productivity (ANPP), species abundance, and soil moisture. We fit a generalized linear mixed effects model to determine if the relationship between ANPP and soil moisture differed among treatments. We used nonmetric multidimensional scaling to quantify community composition over the five years.

**Results.** Ecosystem functional response, defined as the relationship between soil moisture and ANPP was similar among drought and control treatments, but the irrigation treatment had a lower slope than the control treatment. However, all estimates for the effect of soil moisture on ANPP overlapped zero, indicating the relationship is weak and uncertain regardless of treatment. There was also large spatial variation in ANPP within-years, which contributes to the uncertainty of the soil moisture effect. Plant community composition was remarkably stable over the course of the experiment and did not differ among treatments.

**Discussion.** Despite some evidence that ecosystem functional response became less sensitive under sustained wet conditions, the response of ANPP to soil moisture was consistently weak and community composition was stable. The similarity of ecosystem functional responses across treatments was not related to compensatory shifts at the plant community level, but instead may reflect the insensitivity of the dominant species to soil moisture. These species may be successful precisely because they have evolved life history strategies which buffer them against precipitation variability.

# 1 INTRODUCTION

Water availability is a major driver of annual net primary productivity (ANPP) in grassland ecosystems (Huxman et al., 2004; Hsu et al., 2012). Therefore, projected changes in precipitation regimes, associated with global climate change, will impact grassland ecosystem functioning. At any given site, the relationship between ANPP and water availability (e.g., soil moisture) can be characterized by regressing historical observations of ANPP on observations of soil moisture. Fitted functional responses can then be used to infer how ANPP may change under future precipitation regimes (e.g., Hsu et al., 2012). A problem with this approach is that it requires extrapolation if future precipitation falls outside the historical range of variability (Smith, 2011; Peters et al., 2012). For example, the soil moisture-ANPP relationship may be linear within the historical range of interannual variation, but could saturate at higher levels of soil moisture. Saturating relationships are actually common (Hsu et al., 2012; Gherardi and Sala, 2015b), perhaps because other resources, like nitrogen, become more limiting in wet years than dry years. Failure to accurately estimate the curvature of the soil moisture-ANPP relationship will lead to over- or underprediction of ANPP under extreme precipitation (Peters et al., 2012).

Another problem with relying on historical ecosystem functional responses to predict impacts of altered precipitation regimes is that these relationships themselves might shift over the long-term. Shifts in species identities and/or abundances can alter an ecosystem's functional response to water availability because different species have different physiological thresholds. Smith et al. (2009) introduced the 'Hierarchical Response Framework' for understanding the interplay of community composition and ecosystem functioning in response to long-term shifts in resources. In the near term, ecosystem functioning such as ANPP will reflect the physiological responses of individual species to the manipulated resource level. For example, ANPP may decline under simulated drought because the initial community consisted of drought-intolerant species (Hoover et al., 2014). Ecosystem functioning may recover over longer time spans as new species colonize or initial species reorder in relative abundance. It is also possible that ecosystem functioning shifts to a new mean state, reflecting the suite of species in the new community (Knapp et al., 2012).

Experimental manipulations of limiting resources, like precipitation, offer a route to understanding how ecosystems will respond to resource levels that fall outside the historical range of variability (Avolio et al., 2015; Gherardi and Sala, 2015a; Knapp et al., 2017). Altering the amount of precipitation over many years should provide insight into the time scales at which water-limited ecosystems respond to chronic resource alteration. We propose four alternative scenarios for the effect of precipitation manipulation on the ecosystem functional response to soil moisture based on the Hierarchical Response Framework (Fig. 1). We define 'ecosystem functional response' as the relationship between available soil moisture and ANPP. The four scenarios are based on possible outcomes at the community (e.g., community composition) and ecosystem (e.g., soil moisture-ANPP regression) levels.

First, altered precipitation might have no effect on either ecosystem functional response or community composition (Fig. 1, top left). In this case, changes in ANPP would be well predicted by the current, observed soil moisture-ANPP relationship. This corresponds to the early phases of the Hierarchical Response Framework, where ecosystem response follows the physiological responses of individual species. Second, the ecosystem functional response might change while community composition remains the same (Fig. 1, top right). A saturating soil moisture-ANPP response fits this scenario, where individual species hit physiological thresholds or are limited by some other resource. Third, the ecosystem functional response might be constant but community composition changes (Fig. 1, bottom left). In this case, changes in species' identities and/or abundances occur in response to altered precipitation levels and species more suited to the new conditions compensate for reduced function of initial species. Fourth, and last, both ecosystem functional response and community composition could change (Fig. 1, bottom right). New species, or newly abundant species, with different physiological responses completely reshape the ecosystem functional response.

All four outcomes are possible in any given ecosystem, but the time scales at which the different scenarios play out likely differ (Smith et al., 2009; Wilcox et al., 2016; Knapp et al., 2017). To determine these time scales, we need to amass information on how quickly ecosystem functional responses change in different ecosystems. We also need to understand whether changes at the ecosystem level are driven by community level changes or individual level responses.

To that end, here we report the results of a five-year precipitation manipulation experiment in a

sagebrush steppe grassland. We imposed drought and irrigation treatments (approximately  $\pm 50\%$ ) and measured ecosystem (ANPP) and community (species composition) responses. We focus on how the drought and irrigation treatments affect the relationship between interannual variation in available soil moisture and interannual variation in ANPP, and if community dynamics underlie the ecosystem responses. In particular, we are interested in the consistency of the soil moisture-ANPP relationship among treatments. Is the relationship steeper under the drought treatment at low soil moisture? Does the relationship saturate under the irrigation treatment at high soil moisture? To answer these questions we fit a generalized linear mixed effects model to test whether the regressions differed among treatments. We also analyzed community composition and the sensitivity of ANPP to drought and irrigation treatments over time, allowing us to place our experimental results within the framework our scenarios (Fig. 1).

## METHODS

### 2.1 Study Area

We conducted our precipitation manipulation experiment in a sagebrush steppe community at the USDA, U.S. Sheep Experiment Station (USSES) near Dubois, Idaho (44.2° N, 112.1° W), 1500 m above sea level. The plant community is dominated by the shrub *Artemesia tripartita*, the perennial forb *Balsamorhiza sagittata*, and three perennial bunchgrasses, *Pseudoroegneria spicata*, *Poa secunda*, and *Hesperostipa comata* (see Appendix 1 for rank abundance curves). During the period of our experiment (2011 – 2016), mean annual precipitation was 250 mm year<sup>-1</sup> and mean monthly temperature ranged from -5.2°C in January to 21.8°C in July. Between 1926 and 1932, range scientists at the USSES established 26 permanent 1 m<sup>2</sup> quadrats to track vegetation change over time. In 2007, we relocated 14 of the original quadrats in permanent livestock enclosures, which we used as control plots (i.e. ambient precipitation) in the experiment described below. We used the relocated plots as our controls because collecting demographic data is time consuming and it was already being collected in these plots for other studies.

In spring 2011, we established 16 new 1 m<sup>2</sup> plots located in the largest enclosure at USSES, which also contained six of our control plots. We avoided areas on steep hill slopes, areas with greater than 20% cover of bare rock, and areas with greater than 10% cover of the shrubs *Purshia tridentata* and/or *Amelanchier utahensis*. We established the new plots in pairs and randomly assigned each plot in a pair to receive a “drought” or “irrigation” treatment. Thus, our experiment consisted of  $n = 14$  control plots,  $n = 8$  irrigation plots, and  $n = 8$  irrigation plots, for a total of 30 plots.

### 2.2 Precipitation Experiment

Drought and irrigation treatments were designed to decrease and increase the amount of ambient precipitation by 50%. To achieve this, we used a system of rain-out shelters and automatic irrigation (Gherardi and Sala, 2013). The rain-out shelters consisted of transparent acrylic shingles 1-1.5 m above the ground that covered an area of 2.5 × 2 m. The shingles intercepted approximately 50% of incoming rainfall, which was channeled into 75 liter containers. Captured rainfall was then pumped out of the containers and sprayed on to the adjacent irrigation plot via two suspended sprinklers. Pumping was triggered by float switches once water levels reached about 20 liters. We disconnected the irrigation pumps each October and reconnected them each April. The rain-out shelters remained in place throughout the year.

We monitored soil moisture in four of the drought-irrigation pairs using Decagon Devices (Pullman, Washington) 5TM and EC-5 soil moisture sensors. We installed four sensors around the edges of each 1x1 m census plot, two at 5 cm soil depth and two at 25 cm soil depth. We also installed four sensors in areas nearby the four selected plot pairs to measure ambient soil moisture at the same depths. Soil moisture measurements were automatically logged every four hours. We coupled this temporally intensive soil moisture sampling with spatially extensive readings taken with a handheld EC-5 sensor at six points within all 16 plots and associated ambient measurement areas. These snapshot data were collected on 06-06-2012, 04-29-2015, 05-07-2015, 06-09-2015, and 05-10-2016<sup>1</sup>.

Analyzing the response to experimental treatments was complicated by the fact that we did not directly monitor soil moisture in each plot on each day of the experiment. Only a subset of plots were equipped

<sup>1</sup>Dates formatted as: mm-dd-yyyy.

with soil moisture sensors, and within those plots, one or more of the sensors frequently failed to collect data. Therefore, we used a statistical model to estimate average daily soil moisture values for the ambient, drought, and irrigation treatments during the course of the experiment.

We first averaged the observed soil moisture readings for each day ( $d$ ) and plot ( $i$ ),  $x_{i,d}$ . Experimental plots were organized in groups of two, with each group ( $g$ ) containing a drought and irrigated plot. In addition a nearby area outside the drought or irrigated plots was monitored for local ambient soil moisture conditions. Within each group of plots we standardized the irrigation and drought effects on soil moisture relative to the ambient soil moisture conditions. Specifically we subtracted the ambient daily soil moisture from the soil moisture measured within the drought and irrigation plots within each group and then divided by the standard deviation of daily soil moisture values measured in the ambient conditions ( $\Delta x_{g,d,\text{trt.}} = (x_{g,d,\text{trt.}} - x_{g,d,\text{ambient}}) / \text{s.d.}(x_{g,d,\text{ambient}})$ ) where  $\Delta x$  is the standardized treatment effect and  $x_{g,d,\text{trt.}}$  is the raw soil moisture measure for plot group  $g$  on day  $d$  and treatment  $\text{trt.}$ . These transformations ensured that the treatment effects in each plot were appropriately scaled by the local ambient conditions within each plot group.

We then modeled these daily deviations ( $\Delta x_{g,d,\text{trt.}}$ ) from ambient conditions using a linear mixed effects model with independent variables for treatment (irrigation or drought), season (winter, spring, summer, fall), rainfall, and all two-way interactions. Using the local daily weather station data, we recorded rainy days as any day with measureable precipitation or the day after such a day and with average temperatures above 3°C (to exclude days with snowfall). We fit the model using the `lme4::lmer()` function (Bates et al., 2015) in R (R Core Team, 2016), with random effects for plot group and date. We weighted observations by the number of unique sensors or spot measurements that were taken in each plot on that day. We then used the model to predict the deviations from ambient conditions produced in the treated plots on each day of the experiment. We added these predicted deviations to the average daily ambient soil moisture to generate predictions for daily soil moisture in all of the treated plots:  $\bar{x}_{d,\text{trt.}} = \Delta \bar{x}_{d,\text{trt.}} + \bar{x}_{d,\text{ambient}}$ , where  $\bar{x}_{d,\text{trt.}}$  is the average predicted daily soil moisture in the treated plots and  $\bar{x}_{d,\text{ambient}}$  is the daily ambient soil moisture averaged across all control plots. We could only predict soil moisture in the treated plots on days for which we took at least one ambient soil moisture measurement. This procedure allowed us to predict daily soil moisture conditions for all plots, even on those days when some of our direct treatment measurements were missing due to malfunction of the sensors or dataloggers. See Kleinhesselink (2017) for more details on our approach to estimating soil moisture.

Following the above procedure, we still lacked soil moisture data for March 2012 (observations did not start until April) and for a string of days in 2013 during which the soil moisture probes failed take readings. To fill in these gaps, we used a version of the SOILWAT soil moisture model (Sala et al., 1992) that has been specifically designed for semiarid shrublands and grasslands (Bradford et al., 2014). The model was parameterized using generic sagebrush steppe parameters and local soil texture, soil bulk density, and precipitation data (Kleinhesselink, 2017). We used the locally parameterized SOILWAT model to generate daily soil moisture predictions for the duration of the experiment, but only used SOILWAT predictions where there were gaps in our data (Fig. 2B and Appendix 2). SOILWAT predicted daily soil moisture under ambient conditions similar to our control plots. We applied the same statistical model and procedure described above to estimate soil moisture in drought and irrigation plots based on control plot conditions.

## 2.3 Data Collection

We estimated aboveground net primary productivity (ANPP) using a radiometer to relate ground reflectance to plant biomass (see Byrne et al., 2011, for a review). We recorded ground reflectance at four wavelengths, two associated with red reflectance (626 nm and 652 nm) and two associated with near-infrared reflectance (875 nm and 859 nm). At each plot in each year, we took four readings of ground reflectances at the above wavelengths. We also took readings in 12 (2015), 15 (2012, 2013, 2014), or 16 (2016) calibration plots adjacent to the experimental site, in which we harvested all aboveground biomass produced in the current year (we excluded litter and standing dead material), dried it to a constant weight at 60°C, and weighed it to estimate ANPP. We harvested at peak biomass each year, typically in late June.

For each plot and year, we averaged the four readings for each wavelength and then calculated a greenness index based on the same bands used to calculate NDVI using the MODIS (Moderate Resolution Imaging Spectroradiometer) and AVHRR (Advanced Very High Resolution Radiometer) bands for NDVI. We regressed the greenness index against the dry biomass weight from the ten calibration plots to convert

the greenness index to ANPP. We fit regressions to a MODIS-based index and an AVHRR-based index for each year and retained the regression with the better fit based on  $R^2$  values. We then predicted ANPP using the best regression equation for each year (Appendix 3). Our analysis, described below, can be done using NDVI directly, but we focus on ANPP to more easily relate our findings to previous work and because our results do not change if we analyze NDVI directly (Appendix 5).

Species composition data came from two sources: yearly census maps for each plot made using a pantograph (Hill, 1920) and yearly counts of annual species in each plot. From these sources, we determined the density of all annuals and perennials forbs, the basal cover of perennial grasses, and the canopy cover of shrubs. We made a large plot-treatment-year by species matrix, where columns were filled with either basal cover or density, depending on the measurement made for the particular species. We standardized the values in each column so we could directly compare species quantified with different metrics (density, basal cover, and canopy cover). This puts all abundance values on the same scale, meaning that common and rare species are weighted equally. Assuming that rare species will respond to treatments more than common ones, our approach is biased towards detecting compositional changes. We limited our community level data to observations from the permanent enclosure with our drought and irrigation treatments for the community composition analyses described below (**Community composition over time**). This enclosure included six of the control plots and all of the treatment plots, for a total of 22 plots. The other eight control plots are in other pastures, which would add a spatial bias to our community analysis focused on temporal dynamics.

## 2.4 Data Analysis

### 2.4.1 Ecosystem functional response

Our main goal was to test whether the relationship between ANPP and soil moisture differed among the drought, control, and irrigation treatments. Based on our own observations and previous work at our study site (Blaisdell, 1958; Dalgleish et al., 2011; Adler et al., 2012), we chose to use cumulative volumetric water content from March through June as our metric of soil moisture (hereafter referred to as ‘VWC’). To achieve this goal, we fit a generalized linear mixed effects regression model with  $\log(\text{ANPP})$  as the response variable and VWC and treatment as fixed effects. Plot and year of treatment were included as random effects to account for non-independence of observations, as described below. We log-transformed ANPP to account for heteroscedasticity. Both  $\log(\text{ANPP})$  and VWC were standardized to have mean 0 and unit variance before fitting the model [i.e.,  $(x_i - \bar{x})/\sigma_x$ ].

Our model is defined as follows:

$$\mu_i = \beta \mathbf{x}_i + \gamma_{j(i)} \mathbf{z}_i + \eta_t, \quad (1)$$

$$\mathbf{y} \sim \text{Normal}(\boldsymbol{\mu}, \sigma^2), \quad (2)$$

where  $\mu_i$  is the deterministic prediction from the regression model for observation  $i$ , which is associated with plot  $j$  and treatment year  $t$ .  $\beta$  is the vector of coefficients for the fixed effects in the design matrix  $\mathbf{X}$ . Each row of the design matrix represents a single observation ( $\mathbf{x}_i$ ) and is a vector with the following elements: 1 for the intercept, a binary 0 or 1 if the treatment is “drought”, a binary 0 or 1 if the treatment is “irrigation”, the scaled value of VWC, binary “drought” value times VWC, and binary “irrigation” value times VWC. Thus, our model treats “control” observations as the main treatment and then estimates intercept and slope offsets for the “drought” and “irrigation” treatments. We use our model to test two statistical hypotheses based on the questions in our Introduction:

**H1.** The coefficient for drought  $\times$  VWC is positive and different from zero.

**H2.** The coefficient for irrigation  $\times$  VWC is negative and different from zero.

These hypotheses are based on evidence that precipitation-ANPP relationships often saturate with increasing precipitation (Hsu et al., 2012; Gherardi and Sala, 2015b).

We include two random effects to account for the fact that observations within plots and years are not independent. Specifically, we include plot-specific offsets ( $\gamma$ ) for the intercept and slope terms and year-specific intercept offsets ( $\eta_t$ ). The covariate vector  $\mathbf{z}_i$  for each observation  $i$  has two elements: a 1 for the intercept and the scaled value of VWC for that plot and year. The plot-specific coefficients are



modeled hierarchically, where plot level coefficients are drawn from a multivariate normal distribution with mean 0 and a variance-covariance structure that allows the intercept and slope terms to be correlated:

$$\boldsymbol{\gamma}_{j(i)} \sim \text{MVN}(0, \Sigma), \quad (3)$$

where  $\Sigma$  is the variance-covariance matrix and  $j(i)$  reads as “plot  $j$  associated with observation  $i$ ”. The random year effects ( $\boldsymbol{\eta}$ ) are drawn from a normal prior with mean 0 and standard deviation  $\sigma_{\text{year}}$ , which was drawn from a half-Cauchy distribution. We used a vague prior distribution for each  $\beta$ :  $\beta \sim \text{Normal}(0, 5)$ . A full description of our model and associated R (R Core Team, 2016) code is in Appendix 4.

We fit the model using a Bayesian approach and obtained posterior estimates of all unknowns via the No-U-Turn Hamiltonian Monte Carlo sampler in Stan (Stan Development Team, 2016b). We used the R package ‘rstan’ (Stan Development Team, 2016a) to link R (R Core Team, 2016) to Stan. We obtained samples from the posterior distribution for all model parameters from four parallel MCMC chains run for 10,000 iterations, saving every 10<sup>th</sup> sample. Trace plots of all parameters were visually inspected to ensure well-mixed chains and convergence. We also made sure all scale reduction factors ( $\hat{R}$  values) were less than 1.1 (Gelman and Hill, 2009).

## 2.4.2 Sensitivity of ANPP over time

Our data do not allow us to directly test whether the ecosystem functional response to precipitation may have changed over time since the start of the experiment in each of the treatments. This is because we lack sufficient within-year and within-treatment variation of soil moisture (i.e., within a year and treatment each plot shares the same value of VWC). However, we did use a separate analysis to test whether the sensitivity of ANPP to treatment changed over time.

We define ‘sensitivity’ as  $\frac{\text{ANPP}_{\text{control}} - \text{ANPP}_{\text{treatment}}}{\text{VWC}_{\text{ambient}} - \text{VWC}_{\text{treatment}}}$ , following Wilcox et al. (2017). This metric of sensitivity scales the difference between ANPP in the treated and control plots by the change in soil moisture in the treated plots. Because our treated plots are not directly paired with individual control plots, we compare ANPP in each treated plot to the average ANPP of the control plots in each year. After calculating sensitivity for each drought and irrigation plot in each year, we regressed sensitivity against year of treatment using the `lm()` function in R (R Core Team, 2016). This analysis also allows us to link particularly sensitive treatment years to changes in community composition.

## 2.4.3 Community composition over time

We used nonmetric multidimensional scaling (NMDS) based on Bray-Curtis distances to identify temporal changes in community composition among treatments. We first calculated Bray-Curtis distances among all plots for each year of the experiment and then extracted those distances for use in the NMDS. Some values of standardized species’ abundances were negative, which is not allowed for calculating Bray-Curtis distances. We simply added ‘2’ to each abundance value to ensure all values were greater than zero. We plotted the first two axes of NMDS scores to see if community composition overlapped, or not, among treatments in each year. We used the `vegan::metaMDS()` function (Oksanen, 2016) to calculate Bray-Curtis distances and then to run the NMDS analysis. We used the `vegan::adonis()` function (Oksanen, 2016) to perform permutational multivariate analysis of variance to test whether treatment plots formed distinct groupings. To test whether treatment plots were equally dispersed, or not, we used the `vegan::betadisper()` function (Oksanen, 2016).

We conducted the above analysis for all species, and then conducted a separate analysis for annual species only (Appendix 1). Annual species have shorter life spans, so conducting a separate analysis allowed us to discover if community responses are restricted to short-lived species that might respond more quickly to altered precipitation regimes. Given the dominance of perennial species in our system (Appendix 1), shifts in the annual plant community could be difficult to detect in the analysis of the full community.

## 2.4.4 Reproducibility

All R code and data necessary to reproduce our analysis has been archived on Figshare (*link here after acceptance*) and released on GitHub ([https://github.com/atredennick/usses\\_water/](https://github.com/atredennick/usses_water/))

releases/v0.1). We also include annotated Stan code in our model description in Appendix 4.

### 3 RESULTS

Ambient precipitation was variable over the five years of the study (Fig. 2A), meaning our experiment captured a range of environmental conditions. ANPP varied from a minimum of  $77.4 \text{ g m}^{-2}$  in 2014 to a maximum of  $239.3 \text{ g m}^{-2}$  in 2016 when averaged across treatments (Fig. 2C). ANPP was slightly higher in irrigation plots (on average 15% higher) and slightly lower in drought plots (on average 25% lower) relative to control plots (Fig. 2C), corresponding to observed and estimated soil volumetric water content (VWC) differences among treatments (Fig. 2B). March-June VWC in drought plots was 12% less than in control plots on average, and March-June VWC in irrigation plots was 19% higher than in control plots on average across the years of the experiment. Such differences in soil VWC indicate our treatment infrastructure was successful. ANPP was highly variable across plots within years, as indicated by the large and overlapping standard deviations (Fig. 2C).

Cumulative March-June soil moisture had a weak positive effect on ANPP (Table 1; Fig. 3B). The effect of soil moisture for each treatment is associated with high uncertainty, however, with 95% Bayesian credible intervals that broadly overlap zero (Table 1). Although the parameter estimates for the effect of soil moisture overlap zero, the posterior distributions of the slopes all shrank and shifted to more positive values relative to the prior distributions (Fig. A3-2), which indicates the data did influence parameter estimates beyond the information from the uninformative priors. Ecosystem functional response was similar among treatments (Table 1; Fig. 3B), but there is evidence that the slope for the drought treatment is greater than the slope for the control treatment. This evidence comes from interpreting the posterior distributions of the slope offsets for the treatments. From these distributions, we calculate a 58% one-tailed probability that the estimate is greater than zero for the irrigation treatment and a 100% one-tailed probability that the estimate is greater than zero for the drought treatment (Fig. 3A, right panel).

Sensitivity of ANPP to irrigation was constant over the course of our experiment (Fig. 3C). Sensitivity of ANPP to drought, however, grew over time ( $P = 0.0007$ ; Fig. 3C).

Community composition was similar among treatments. The multidimensional space of community composition overlapped among treatments in all years and was equally dispersed in all years (Table 2; Fig. 4). Community composition was also remarkably stable over time, with no evidence of divergence among treatments (Table 2; Fig. 4). There were also no changes in the dominant species over time in any treatment (Appendix 1). Analyzing annual species on their own produced similar results. There is some evidence that the annual community changed in response to treatment in two years (Fig. A1-6), but these responses appear to come from very rare species since our analysis weights common and rare species equally (Fig. A1-4). By definition, these rare species contribute little to ANPP and thus any compositional changes have little influence on ecosystem functional response.

### 4 DISCUSSION

Ecosystem response to a new precipitation regime depends on the physiological responses of constituent species and the rate at which community composition shifts to favor species better able to take advantage of, or cope with, new resource levels (Smith et al., 2009). Previous work has shown that community compositional shifts can be both rapid, on the order of years (Hoover et al., 2014), and slow, on order of decades (Knapp et al., 2012; Wilcox et al., 2016). A lingering question is how the time scales of ecosystem response and community change vary among ecosystems. Precipitation manipulation experiments can help answer this question, especially if they push water availability outside the historical range of variability for long periods.

The results of our five year experiment in a sagebrush steppe conforms to one of our four predictions. Ecosystem functional response under chronic drought was different from the control treatment, but community composition remained unchanged (Fig. 3A, Fig. 4), representing the top right scenario in Fig. 1. The increase in the slope of the  $\text{VWC} \times \text{productivity}$  relationship in the drought plots is consistent with a stronger response of productivity to additional water in this ecosystem. This suggests that if average soil moisture were pushed consistently lower than currently observed ambient conditions, there would be a stronger relationship between precipitation and productivity in this system.

343 However, we do not want to over-interpret the significance of the slope offsets given that the full  
344 distributions of the slopes of the VWC  $\times$  productivity relationship, not just the offsets, were similar  
345 among treatments (Table 1). We therefore conclude that ecosystem functional response is consistent  
346 (similar values) and weak (all broadly overlapping zero) across all precipitation treatments.

347 The similarity of ecosystem functional response (Fig. 3) and community composition (Fig. 4) among  
348 treatments is surprising because grasslands generally, and sagebrush steppe specifically, are considered  
349 water-limited systems. For example, Huxman et al. (2004) and Knapp et al. (2015) showed that semi-arid  
350 sites are more sensitive to drought than mesic sites, and Wilcox et al. (2017) found that semi-arid sites are  
351 particularly sensitive to irrigation treatments. Based on these findings, we expected ecosystem functional  
352 response, community composition, or both to change under our treatments. Why did our treatments fail  
353 to induce ecosystem or community responses? We can think of three reasons. Two are limitations of  
354 our study, and one is the life history traits of the species in our focal communities. We first discuss the  
355 potential limitations of our study, and then discuss the biological explanation.

356 First, it could be that our precipitation manipulations were not large enough to induce a response. That  
357 is, a 50% decrease in any given year may not be abnormal given our site's historical range of variability  
358 (Knapp et al., 2017). We cannot definitively rule out this possibility, but we have reason to believe our  
359 treatments *should* have been large enough. Using the methods described by Lemoine et al. (2016), we  
360 calculated the percent reduction and increase of water-year precipitation necessary to reach the 1% and  
361 99% extremes of the historical precipitation regime at our site (Fig. A5-1). The 1% quantile of water-year  
362 precipitation at our site is 78 mm, a 26% reduction. Thus, our drought treatment represented extreme precipitation  
363 amounts, and that is the treatment where we observed a small effect on the slope between soil moisture  
364 and ANPP (Fig. 3A). The irrigation treatment may not have been extreme enough to induce a response.

365 Second, ANPP at our site may be influenced by additional factors, not only the cumulative March-June  
366 soil moisture covariate we included in our statistical model. For example, La Pierre et al. (2016) found  
367 that site-scale ANPP is better predicted by abiotic factors such as nutrient availability than precipitation.  
368 Moreover, temperature can impact ANPP directly (Epstein et al., 1997) and by exacerbating the effects  
369 of soil moisture (De Boeck et al., 2011). Measurements of soil moisture likely contain a signal of  
370 temperature, through its impact on evaporation and infiltration, but the measurements will not capture  
371 the direct effect of temperature on metabolic and physiological processes. We also did not redistribute  
372 snow across our treatments in the winter, and snow melt may spur early spring growth. Failure to account  
373 for potentially important covariates could explain the within-year spread of ANPP (Fig 2C, Fig. 3B) and  
374 the resulting uncertain relationship we observed between soil moisture and ANPP across all treatments  
375 (Table 1, Fig. A3-2). Unfortunately, we did not collect data on additional factors because our focus was  
376 on precipitation.

377 Third, the life history traits of the dominant species in our study ecosystem may explain the consistently  
378 positive, but weak and uncertain, effect of soil moisture on ANPP (Table 1, Fig. 3). Species that live  
379 in variable environments, such as cold deserts, must have strategies to ensure long-term success as  
380 conditions vary. One strategy is bet hedging, where species forego short-term gains to reduce the variance  
381 of long-term success (Seger, 1987). In other words, species follow the same conservative strategy every  
382 year, designed to minimize response to environmental conditions. The dry and variable environment of  
383 the sagebrush steppe has likely selected for bet hedging species that can maintain function at low water  
384 availability and have weak responses to high water availability. In so doing, the dominant species in our  
385 ecosystem avoid “boom and bust” cycles, which corresponds to the weak effect of soil moisture on ANPP  
386 (i.e., the Bayesian credible intervals for the slopes overlapping zero; Table 1).

387 Another strategy to deal with variable environmental conditions is avoidance, which would also result  
388 in a consistent ecosystem functional response between drought and control treatments. For example,  
389 many of the perennial grasses in our focal ecosystem avoid drought stress by growing early in the growing  
390 season (Blaisdell, 1958, A.R. Kleinhesselink, personal observation). Furthermore, the dominant shrub in  
391 our focal ecosystem, *Artemisia tripartita*, has access to water deep in the soil profile thanks to a deep  
392 root system (Kulmatiski et al., 2017). The dominance of our site by the shrub *A. tripartita* (Appendix 1)  
393 may explain why our results do not conform to broader patterns of grassland sensitivity to precipitation  
394 manipulations (e.g., Huxman et al., 2004; Knapp et al., 2015; Wilcox et al., 2017).

395 We found that ANPP became more sensitive to the drought over time (Fig. 3C). In other words,  
396 relative to the effect of the drought shelters on soil moisture, ANPP was more reduced by the drought  
397 conditions later in the experiment than early on in the experiment. We interpret this increase in sensitivity



398 as the effect of cumulative impacts of the drought on dominant species. Plants may not have shown  
399 a large response in terms of ANPP in the first years of the experiment if they could grow from stored  
400 carbohydrate reserves or if they abstained from flowering and reproduction. As the drought progressed,  
401 these same plants may have started to reduce in size or die. Given the long lived perennial species at  
402 this site, this increase in sensitivity may indicate a larger future change in community composition and  
403 ecosystem functional response.

## 404 5 CONCLUSIONS

405 Our results suggest the species in our focal plant community are insensitive to to changes in precipitation  
406 regime, [at least over the five years of our experiment](#). Such insensitivity can buffer species against  
407 precipitation variability in this semi-arid ecosystem, making them successful in the long run. Longer,  
408 chronic precipitation alteration might reveal plant community shifts that we did not observe (e.g., Wilcox  
409 et al., 2016), [and the increased sensitivity of ANPP to drought in the final years of our experiment](#)  
410 [may portend such a shift](#). Likewise, a long-term increase in water availability could allow species that  
411 do not bet hedge to gain prominence and dominate the ecosystem functional response. [The length](#)  
412 [of the perturbation may be especially relevant in our focal ecosystem where the perennial species are](#)  
413 [long-lived, meaning compositional turnover may take many more years than we report on here](#). Our  
414 results suggest compositional shifts would have the largest impact at high rainfall because the current  
415 community maintained consistent ecosystem functional response at very low water availability. [However,](#)  
416 [the increased sensitivity of ANPP to drought conditions suggests a shift in ecosystem functional response](#)  
417 [may be near, and simply takes more time than we observed here](#).

## 418 6 ACKNOWLEDGEMENTS

419 We thank the many summer research technicians who collected the data reported in this paper and the  
420 USDA, U.S. Sheep Experiment Station (Dubois, ID) for facilitating work on their property. We also  
421 thank Susan Durham for clarifying our thinking on the statistical analyses and Kevin Wilcox for helpful  
422 discussions on analyzing community composition data. John Bradford and Caitlin Andrews generously  
423 provided SOILWAT model output, [Two anonymous reviewers and Elsie Denton provided thoughtful](#)  
424 [reviews that improved our paper](#).

## 425 7 FUNDING

426 This research was supported by the Utah Agricultural Experiment Station, Utah State University, and  
427 approved as journal paper number 9035. The research was also supported by the National Science  
428 Foundation, through a Postdoctoral Research Fellowship in Biology and Mathematics to ATT (DBI-  
429 1400370), a Graduate Research Fellowship to ARK, and grants DEB-1353078 and DEB-1054040 to  
430 PBA.

## 431 8 AUTHOR CONTRIBUTIONS

- 432 • Andrew T. Tredennick collected data, analyzed the data, wrote the paper, prepared figures and/or  
433 tables, reviewed drafts of the paper.
- 434 • Andrew R. Kleinhesselink conceived and designed the experiments, performed the experiments,  
435 collected data, analyzed the data, reviewed drafts of the paper.
- 436 • J. Bret Taylor contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- 437 • Peter B. Adler conceived and designed the experiments, performed the experiments, collected data,  
438 analyzed the data, reviewed drafts of the paper.

## 439 **9 SUPPLEMENTAL INFORMATION**

440 **Appendix 1.** Additional information on plant community composition and dynamics, Table A1-1, Table  
441 A1-2, Table A1-3, Fig. A1-1, Fig. A1-2, Fig. A1-3, Fig. A1-4, Fig. A1-5, and Fig. A1-6.

442 **Appendix 2.** Additional details on soil moisture modeling with SOILWAT and Fig. A2-1.

443 **Appendix 3.** Additional methods and information on estimating aboveground net primary productivity,  
444 Table A3-1, and Table A3-2.

445 **Appendix 4.** Details of the hierarchical Bayesian model, Fig. A4-1, Fig. A4-2, and Fig. A4-3.

446 **Appendix 5.** Results from analysis of NDVI without conversion to ANPP, Fig. A5-1.

447 **Appendix 6.** Details on analysis of precipitation historical range of variability and Fig. A6-1.


## 10 TABLES

**Table 1.** Summary statistics from the posterior distributions of coefficients for each treatment ( $\beta$  coefficients in equation 1). The ‘Intercept’ and ‘Slope’ summaries reported here for drought and irrigation are from the posterior distributions of the intercept and slope for the control treatment plus the offsets for each treatment. Posterior distributions of the offsets are in Figure 3A.


Coefficient	Treatment	Posterior Mean	Posterior Median	Lower 95% BCI	Upper 95% BCI
Intercept	Control	0.14	0.14	-1.24	1.50
Intercept	Drought	0.67	0.65	-2.01	3.36
Intercept	Irrigation	-0.39	-0.34	-3.51	2.48
Slope	Control	0.53	0.52	-1.44	2.66
Slope	Drought	1.02	1.00	-1.04	3.25
Slope	Irrigation	0.56	0.55	-1.43	2.71

**Table 2.** Results from statistical tests for clustering and dispersion of community composition among precipitation treatments. ‘adonis’ tests whether treatments form unique clusters in multidimensional space; ‘betadisper’ tests whether treatments have similar dispersion. For both tests,  $P$  values greater than 0.05 indicate there is no support that the treatments differ.

Year	Test	n	d.f.	$F$	$P$
2011	adonis	22	2	1.04	0.41
2011	betadisper	22	2	2.22	0.14
2012	adonis	22	2	1.09	0.32
2012	betadisper	22	2	0.21	0.81
2013	adonis	22	2	1.26	0.14
2013	betadisper	22	2	0.43	0.66
2014	adonis	22	2	0.96	0.52
2014	betadisper	22	2	0.36	0.70
2015	adonis	22	2	1.08	0.31
2015	betadisper	22	2	1.96	0.17
2016	adonis	22	2	1.12	0.27
2016	betadisper	22	2	0.27	0.76




**Figure 1.** Possible outcomes of chronic resource alteration based on the 'Hierarchical Response Framework' (Smith et al. 2009).




**Figure 2.** (A) Probability density of historical water-year (Oct.–Sept.) precipitation from 1927–2016, with the years of the experiment shown with arrows on the  $x$ -axis. (B) Statistically estimated (solid lines) and SOILWAT-generated (dashed lines) daily soil volumetric water content (VWC) in each of the three treatments during the course of the experiment. We used the estimates from our statistical model, with gaps filled in by SOILWAT predictions, to calculate cumulative March–June VWC as used in our analysis of treatment effects on ecosystem functional response. (C) Mean (points) ANPP and its standard deviation (error bars) for each year of the experiment. Colors in panels B and C identify the treatment, as specified in the legend of panel C.





**Figure 3.** Results from the generalized linear mixed effects model (A-B) and the sensitivity analysis (C). (A) Posterior distributions for the effects of drought and irrigation on the intercept and slope of the productivity-soil moisture relationship. Treatment effects show the difference between the coefficients estimated in the treated plots and the control plots. Probabilities (“Pr ≤ 0 =”) for each coefficient indicate the one-tailed probability that the coefficient is less than or greater than zero, depending on whether the median of the distribution is less than or greater than zero. The posterior densities were smoothed for visual clarity by increasing kernel bandwidth by a factor of five. (B) Scatterplot of the data and model estimates shown as solid lines. Model estimates come from treatment level coefficients (colored lines). Note that we show log(ANPP) on the y-axis of panel B; this same plot can be seen on the arithmetic scale in supporting material Fig. A2-1. (C) Regression of sensitivity against time for each treatment. Each point represents the sensitivity of ANPP in a plot relative to the mean of the control plots in that year. Only the significant regression for the drought treatment is plotted ( $P = 0.0007$ ).



**Figure 4.** Nonmetric multidimensional scaling scores representing plant communities in each plot, colored by treatment. *Each point represents a plot and the axes represent species composition.*

## 450 References

- 451 Adler, P. B., Dalgleish, H. J., and Ellner, S. P. (2012). Forecasting plant community impacts of climate  
452 variability and change: when do competitive interactions matter? *Journal of Ecology*, 100:478–487.
- 453 Avolio, M. L., Pierre, K. J. L., Houseman, G. R., Koerner, S. E., Grman, E., Isbell, F., Johnson, D. S.,  
454 and Wilcox, K. R. (2015). A framework for quantifying the magnitude and variability of community  
455 responses to global change drivers. *Ecosphere*, 6(12):1–14.
- 456 Bates, D., Maechler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using  
457 lme4.
- 458 Blaisdell, J. P. (1958). Seasonal development and yield of native plants on the upper snake river plains  
459 and their relation to certain climate factors. *United States Department of Agriculture Technical Bulletin*  
460 *No. 1190*.
- 461 Bradford, J. B., Schlaepfer, D. R., and Lauenroth, W. K. (2014). Ecohydrology of Adjacent Sagebrush  
462 and Lodgepole Pine Ecosystems: The Consequences of Climate Change and Disturbance. *Ecosystems*,  
463 17(4):590–605.
- 464 Byrne, K. M., Lauenroth, W. K., Adler, P. B., and Byrne, C. M. (2011). Estimating Aboveground Net  
465 Primary Production in Grasslands: A Comparison of Nondestructive Methods. *Rangeland Ecology and*  
466 *Management*, 64(5):498–505.
- 467 Dalgleish, H. J., Koons, D. N., Hooten, M. B., Moffet, C. A., and Adler, P. B. (2011). Climate influences  
468 the demography of three dominant sagebrush steppe plants. *Ecology*, 92(1):75–85.
- 469 De Boeck, H. J., Dreesen, F. E., Janssens, I. A., and Nijs, I. (2011). Whole-system responses of  
470 experimental plant communities to climate extremes imposed in different seasons. *New Phytologist*,  
471 189(3):806–817.
- 472 Epstein, H. E., Lauenroth, W. K., and Burke, I. C. (1997). Effects of temperature and soil texture on  
473 ANPP in the U.S. Great plains. *Ecology*, 78(8):2628–2631.
- 474 Gelman, A. and Hill, J. (2009). *Data analysis using regression and multilevel/hierarchical models*.  
475 Cambridge University Press, Cambridge.
- 476 Gherardi, L. A. and Sala, O. E. (2013). Automated rainfall manipulation system: a reliable and inexpensive  
477 tool for ecologists. *Ecosphere*, 4(2):1–10.
- 478 Gherardi, L. A. and Sala, O. E. (2015a). Enhanced interannual precipitation variability increases  
479 plant functional diversity that in turn ameliorates negative impact on productivity. *Ecology Letters*,  
480 18(12):1293–1300.
- 481 Gherardi, L. A. and Sala, O. E. (2015b). Enhanced precipitation variability decreases grass- and increases  
482 shrub-productivity. *Proceedings of the National Academy of Sciences*, 112(41):12735–12740.
- 483 Hill, R. R. (1920). Charting Quadrats with a Pantograph. *Ecology*, 1(4):270–273.
- 484 Hoover, D. L., Knapp, A. K., and Smith, M. D. (2014). Resistance and resilience of a grassland ecosystem  
485 to climate extremes. *Ecology*, 95(9):2646–2656.
- 486 Hsu, J. S., Powell, J., and Adler, P. B. (2012). Sensitivity of mean annual primary production to  
487 precipitation. *Global Change Biology*, 18(7):2246–2255.
- 488 Huxman, T. E., Smith, M. D., Fay, P. A., Knapp, A. K., Shaw, M. R., Loik, M. E., Smith, S. D., Tissue,  
489 D. T., Zak, J. C., Weltzin, J. F., Pockman, W. T., Sala, O. E., Haddad, B. M., Harte, J., Koch, G. W.,  
490 Schwinning, S., Small, E. E., and Williams, D. G. (2004). Convergence across biomes to a common  
491 rain-use efficiency. *Nature*, 429(6992):651–654.
- 492 Kleinhesselink, A. R. (2017). *Direct and indirect effects of climate change on plant populations and*  
493 *communities in sagebrush steppe*. Dissertation, Utah State University.
- 494 Knapp, A. K., Avolio, M. L., Beier, C., Carroll, C. J. W., Collins, S. L., Dukes, J. S., Fraser, L. H., Griffin-  
495 Nolan, R. J., Hoover, D. L., Jentsch, A., Loik, M. E., Phillips, R. P., Post, A. K., Sala, O. E., Slette, I. J.,  
496 Yahdjian, L., and Smith, M. D. (2017). Pushing precipitation to the extremes in distributed experiments:  
497 recommendations for simulating wet and dry years. *Global Change Biology*, 23(5):1774–1782.
- 498 Knapp, A. K., Briggs, J. M., and Smith, M. D. (2012). Community stability does not preclude ecosystem  
499 sensitivity to chronic resource alteration. *Functional Ecology*, 26(6):1231–1233.
- 500 Knapp, A. K., Carroll, C. J., Denton, E. M., La Pierre, K. J., Collins, S. L., and Smith, M. D. (2015).  
501 Differential sensitivity to regional-scale drought in six central US grasslands. *Oecologia*, 177(4):949–  
502 957.
- 503 Kulmatiski, A., Adler, P. B., Stark, J. M., and Tredennick, A. T. (2017). Water and nitrogen uptake are

504 better associated with resource availability than root biomass. *Ecosphere*, 8(3).

505 La Pierre, K. J., Blumenthal, D. M., Brown, C. S., Klein, J. A., and Smith, M. D. (2016). Drivers of  
 506 Variation in Aboveground Net Primary Productivity and Plant Community Composition Differ Across  
 507 a Broad Precipitation Gradient. *Ecosystems*, 19(3):521–533.

508 Lemoine, N. P., Sheffield, J., Dukes, J. S., Knapp, A. K., and Smith, M. D. (2016). Terrestrial Precipitation  
 509 Analysis (TPA): A resource for characterizing long-term precipitation regimes and extremes. *Methods*  
 510 *in Ecology and Evolution*, 7(11):1396–1401.

511 Oksanen, J. (2016). *Vegan: ecological diversity*.

512 Peters, D. P. C., Yao, J., Sala, O. E., and Anderson, J. P. (2012). Directional climate change and potential  
 513 reversal of desertification in arid and semiarid ecosystems. *Global Change Biology*, 18(1):151–163.

514 R Core Team (2016). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.

515 Sala, O. E., Lauenroth, W. K., and Parton, W. J. (1992). Long-term soil water dynamics in the shortgrass  
 516 steppe. *Ecology*, 73(4):1175–1181.

517 Seger, J. (1987). What is bet-hedging? In Harvey, P. and Partridge, L., editors, *Oxford surveys in*  
 518 *evolutionary biology*, pages 182–211. Oxford University Press, Oxford.

519 Smith, M. (2011). An ecological perspective on extreme climatic events: A synthetic definition and  
 520 framework to guide future research. *Journal of Ecology*, 99(3):656–663.

521 Smith, M. D., Knapp, A. K., and Collins, S. L. (2009). A framework for assessing ecosystem dynamics in  
 522 response to chronic resource alterations induced by global change. *Ecology*, 90(12):3279–3289.

523 Stan Development Team (2016a). Rstan: the R interface to Stan, Version 2.14.1.

524 Stan Development Team (2016b). Stan: A C++ Library for Probability and Sampling, Version 2.14.1.

525 Wilcox, K. R., Blair, J. M., Smith, M. D., and Knapp, A. K. (2016). Does ecosystem sensitivity to  
 526 precipitation at the site-level conform to regional-scale predictions? *Ecology*, 97(3):561–568.

527 Wilcox, K. R., Shi, Z., Gherardi, L. A., Lemoine, N. P., Koerner, S. E., Hoover, D. L., Bork, E.,  
 528 Byrne, K. M., Cahill, J., Collins, S. L., Evans, S., Gilgen, A. K., Holub, P., Jiang, L., Knapp, A. K.,  
 529 LeCain, D., Liang, J., Garcia-Palacios, P., Peñuelas, J., Pockman, W. T., Smith, M. D., Sun, S., White,  
 530 S. R., Yahdjian, L., Zhu, K., and Luo, Y. (2017). Asymmetric responses of primary productivity  
 531 to precipitation extremes: A synthesis of grassland precipitation manipulation experiments. *Global*  
 532 *Change Biology*, 23(10):4376–4385.