

Ecosystem functional response across precipitation extremes in a sagebrush steppe

Andrew T. Tredennick^{1,*}, Andrew R. Kleinbesselink^{1,2,*}, J. Bret Taylor³, and Peter B. Adler¹

¹Department of Wildland Resources and the Ecology Center, Utah State University, Logan, Utah 84322

²Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los Angeles, California 90095

³United States Department of Agriculture, Agricultural Research Service, U.S. Sheep Experiment Station, Dubois, Idaho 83423

Corresponding author:

Andrew T. Tredennick^{1,*}

Email address: atredenn@gmail.com

ABSTRACT

Background. Precipitation is predicted to become more variable in the western United States, meaning years of above and below average precipitation will become more common. Periods of extreme precipitation are major drivers of interannual variability in ecosystem functioning in water limited communities, but how ecosystems respond to these extremes over the long-term may shift with precipitation means and variances. Long-term changes in ecosystem functional response could reflect compensatory changes in species composition or species reaching physiological thresholds at extreme precipitation levels.

Methods. We conducted a five year precipitation manipulation experiment in a sagebrush steppe ecosystem in Idaho, United States. We used drought and irrigation treatments (approximately 50% decrease/increase) to investigate whether ecosystem functional response remains consistent under sustained high or low precipitation. We recorded data on aboveground net primary productivity (ANPP), species abundance, and soil moisture. We fit a generalized linear mixed effects model to determine if the relationship between ANPP and soil moisture differed among treatments. We used nonmetric multidimensional scaling to quantify community composition over the five years.

Results. Ecosystem functional response, defined as the relationship between soil moisture and ANPP was similar among drought and control treatments, but the irrigation treatment had a lower slope than the control treatment. However, all estimates for the effect of soil moisture on ANPP overlapped zero, indicating the relationship is weak and uncertain regardless of treatment. There was also large spatial variation in ANPP within-years, which contributes to the uncertainty of the soil moisture effect. Plant community composition was remarkably stable over the course of the experiment and did not differ among treatments.

Discussion. Despite some evidence that ecosystem functional response became less sensitive under sustained wet conditions, the response of ANPP to soil moisture was consistently weak and community composition was stable. The similarity of ecosystem functional responses across treatments was not related to compensatory shifts at the plant community level, but instead may reflect the insensitivity of the dominant species to soil moisture. These species may be successful precisely because they have evolved life history strategies which buffer them against precipitation variability.

1 INTRODUCTION

Water availability is a major driver of annual net primary productivity (ANPP) in grassland ecosystems (Huxman et al., 2004; Hsu et al., 2012). Therefore, projected changes in precipitation regimes, associated with global climate change, will impact grassland ecosystem functioning. Increases in precipitation variability will result in prolonged periods of extremely wet or dry years, and predicting the consequences of such chronic resource alterations requires understanding how ecosystems respond to above and below average precipitation levels.

At any given site, the relationship between ANPP and water availability (e.g., soil moisture) can be characterized by regressing historical observations of ANPP on observations of soil moisture. Fitted functional responses can then be used to infer how ANPP may change under future precipitation regimes (e.g., Hsu et al., 2012). A problem with this approach is that it requires extrapolation if future precipitation falls outside the historical range of variability (Smith, 2011; Peters et al., 2012). For example, the soil moisture-ANPP relationship may be linear within the historical range of interannual variation, but could saturate at higher levels of soil moisture. Saturating relationships are actually common (Hsu et al., 2012; Gherardi and Sala, 2015b), perhaps because other resources, like nitrogen, become more limiting in wet years than dry years. Failure to accurately estimate the curvature of the soil moisture-ANPP relationship will lead to over- or underprediction of ANPP under extreme precipitation (Peters et al., 2012).

Another problem with relying on historical ecosystem functional responses to predict impacts of altered precipitation regimes is that these relationships themselves might shift over the long-term. Shifts in species identities and/or abundances can alter an ecosystem's functional response to water availability because different species have different physiological thresholds. Smith et al. (2009) introduced the 'Hierarchical Response Framework' for understanding the interplay of community composition and ecosystem functioning in response to long-term shifts in resources. In the near term, ecosystem functioning such as ANPP will reflect the physiological responses of individual species to the manipulated resource level. For example, ANPP may decline under simulated drought because the initial community consisted of drought-intolerant species (Hoover et al., 2014). Ecosystem functioning may recover over longer time spans as new species colonize or initial species reorder in relative abundance. It is also possible that ecosystem functioning shifts to a new mean state, reflecting the suite of species in the new community (Knapp et al., 2012).

Experimental manipulations of limiting resources, like precipitation, offer a route to understanding how ecosystems will respond to resource levels that fall outside the historical range of variability (Avolio et al., 2015; Gherardi and Sala, 2015a; Knapp et al., 2017). Altering the amount of precipitation over many years should provide insight into the time scales at which water-limited ecosystems respond to chronic resource alteration. We propose four alternative scenarios for the effect of precipitation manipulation on the ecosystem functional response to soil moisture based on the Hierarchical Response Framework (Fig. 1). We define 'ecosystem functional response' as the relationship between available soil moisture and ANPP. The four scenarios are based on possible outcomes at the community (e.g., community composition) and ecosystem (e.g., soil moisture-ANPP regression) levels.

First, altered precipitation might have no effect on either ecosystem functional response or community composition (Fig. 1, top left). In this case, changes in ANPP would be well predicted by the current, observed soil moisture-ANPP relationship. This corresponds to the early phases of the Hierarchical Response Framework, where ecosystem response follows the physiological responses of individual species. Second, the ecosystem functional response might change while community composition remains the same (Fig. 1, top right). A saturating soil moisture-ANPP response fits this scenario, where individual species hit physiological thresholds or are limited by some other resource. Third, the ecosystem functional response might be constant but community composition changes (Fig. 1, bottom left). In this case, changes in species' identities and/or abundances occur in response to altered precipitation levels and species more suited to the new conditions compensate for reduced function of initial species. Fourth, and last, both ecosystem functional response and community composition could change (Fig. 1, bottom right). New species, or newly abundant species, with different physiological responses completely reshape the ecosystem functional response.

All four outcomes are possible in any given ecosystem, but the time scales at which the different scenarios play out likely differ (Smith et al., 2009; Wilcox et al., 2016; Knapp et al., 2017). To determine these time scales, we need to amass information on how quickly ecosystem functional responses change

in different ecosystems. We also need to understand whether changes at the ecosystem level are driven by community level changes or individual level responses.

To that end, here we report the results of a five-year precipitation manipulation experiment in a sagebrush steppe grassland. We imposed drought and irrigation treatments (approximately $\pm 50\%$) and measured ecosystem (ANPP) and community (species composition) responses. We focus on how the drought and irrigation treatments affect the relationship between interannual variation in available soil moisture and interannual variation in ANPP, and if community dynamics underlie the ecosystem responses. In particular, we are interested in the consistency of the soil moisture-ANPP relationship among treatments. Is the relationship steeper under the drought treatment at low soil moisture? Does the relationship saturate under the irrigation treatment at high soil moisture? To answer these questions we fit a generalized linear mixed effects model to test whether the regressions differed among treatments. We also analyzed community composition and the sensitivity of ANPP to drought and irrigation treatments over time, allowing us to place our experimental results within the framework our scenarios (Fig. 1).

METHODS

2.1 Study Area

We conducted our precipitation manipulation experiment in a sagebrush steppe community at the USDA, U.S. Sheep Experiment Station (USSES) near Dubois, Idaho (44.2° N, 112.1° W), 1500 m above sea level. The plant community is dominated by the shrub *Artemesia tripartita*, the perennial forb *Balsamorhiza sagittata*, and three perennial bunchgrasses, *Pseudoroegneria spicata*, *Poa secunda*, and *Hesperostipa comata* (see Appendix 1 for rank abundance curves). During the period of our experiment (2011 – 2015), average mean annual precipitation was 265 mm year⁻¹ and mean monthly temperature ranged from -5.2°C in January to 21.8°C in July. Between 1926 and 1932, range scientists at the USSES established 26 permanent 1 m² quadrats to track vegetation change over time. In 2007, we relocated 14 of the original quadrats in permanent livestock exclosures, which we used as control plots (i.e. ambient precipitation) in the experiment described below. We used the relocated plots as our controls because collecting demographic data is difficult and it is already being collected in these plots for other ongoing studies.

In spring 2011, we established 16 new 1 m² plots located in the largest exclosure at USSES, which also contained six of our control plots. We avoided areas on steep hill slopes, areas with greater than 20% cover of bare rock, and areas with greater than 10% cover of the shrubs *Purshia tridentata* and/or *Amelanchier utahensis*. We established the new plots in pairs and randomly assigned each plot in a pair to receive a “drought” or “irrigation” treatment. Thus, our experiment consisted of $n = 14$ control plots, $n = 8$ irrigation plots, and $n = 8$ drought plots, for a total of 30 plots.

2.2 Precipitation Experiment

Drought and irrigation treatments were designed to decrease and increase the amount of ambient precipitation by 50%. To achieve this, we used a system of rain-out shelters and automatic irrigation (Gherardi and Sala, 2013). The rain-out shelters consisted of transparent acrylic shingles 1-1.5 m above the ground that covered an area of 2.5×2 m. The shingles intercepted approximately 50% of incoming rainfall, which was channeled into 75 liter containers. Captured rainfall was then pumped out of the containers and sprayed on to the adjacent irrigation plot via two suspended sprinklers. Pumping was triggered by float switches once water levels reached about 20 liters. We disconnected the irrigation pumps each October and reconnected them each April. The rain-out shelters remained in place throughout the year.

We monitored soil moisture in four of the drought-irrigation pairs using Decagon Devices (Pullman, Washington) 5TM and EC-5 soil moisture sensors. We installed four sensors around the edges of each 1x1 m census plot, two at 5 cm soil depth and two at 25 cm soil depth. We also installed four sensors in areas nearby the four selected plot pairs to measure ambient soil moisture at the same depths. Soil moisture measurements were automatically logged every four hours. We coupled this temporally intensive soil moisture sampling with spatially extensive readings taken with a handheld EC-5 sensor at six points within all 16 plots and associated ambient measurement areas. These snapshot data were collected on

06-06-2012, 04-29-2015, 05-07-2015, 06-09-2015, and 05-10-2016¹.

Analyzing the response to experimental treatments was complicated by the fact that we did not directly monitor soil moisture in each plot on each day of the experiment. Only a subset of plots were equipped with soil moisture sensors, and within those plots, one or more of the sensors frequently failed to collect data. To remedy these problems, and to produce average daily soil moisture values for the ambient, drought, and irrigation treatments, we used a statistical model to predict the average treatment effects on soil moisture during the course of the experiment.

We first averaged the observed soil moisture readings for each day (d) and plot (i), $x_{i,d}$. Plots were organized in groups of three, with each group (g) containing consisting of a control, a drought, and an irrigation plot. Within each group of plots we standardized the irrigation and drought effects on soil moisture relative to the ambient soil moisture conditions in the control plot. Specifically we subtracted the ambient daily soil moisture measured in the control plot from the soil moisture measured within the drought and irrigation plots within each group and then divided by the standard deviation of daily soil moisture values measured in the control plot ($\Delta x_{g,d,\text{trt.}} = (x_{g,d,\text{trt.}} - x_{g,d,\text{control}}) / \text{s.d.}(x_{g,d,\text{control}})$) where Δx is the standardized treatment effect and $x_{g,d,\text{trt.}}$ is the raw soil moisture measure for plot group g on day d and treatment trt. . These transformations ensured that the treatment effects in each plot were appropriately scaled by the local ambient conditions within each plot group.

We then modeled these daily deviations ($\Delta x_{g,d,\text{trt.}}$) from ambient conditions using a linear mixed effects model with independent variables for treatment (irrigation or drought), season (winter, spring, summer, fall), rainfall, and all two-way interactions. Rainy days were defined as any day in which precipitation was recorded and average temperature was above 3°C. The day immediately following rainfall was also classified as rainy. We fit the model using the `lme4::lmer()` function (Bates et al., 2015) in R (R Core Team, 2016), with random effects for plot group and date. We weighted observations by the number of unique sensors or spot measurements that were taken in each plot on that day. We then used the model to predict the deviations from ambient conditions produced in the treated plots on each day of the experiment. We added these predicted deviations to the average daily ambient soil moisture to generate predictions for daily soil moisture in all of the treated plots: $\bar{x}_{d,\text{trt.}} = \Delta \bar{x}_{d,\text{trt.}} + \bar{x}_{d,\text{control}}$, where $\bar{x}_{d,\text{trt.}}$ is the average predicted daily soil moisture in the treated plots and $\bar{x}_{d,\text{control}}$ is the daily ambient soil moisture averaged across all control plots. We could only predict soil moisture in the treated plots on days for which we took at least one ambient soil moisture measurement. This procedure allowed us to predict daily soil moisture conditions for all plots, even on those days when some of our direct treatment measurements were missing due to malfunction of the sensors or dataloggers. See Kleinhesslink (2017) for more details on our approach to estimating soil moisture.

2.3 Data Collection

We estimated aboveground net primary productivity (ANPP) using a radiometer to relate ground reflectance to plant biomass (see Byrne et al., 2011, for a review). We recorded ground reflectance at four wavelengths, two associated with red reflectance (626 nm and 652 nm) and two associated with near-infrared reflectance (875 nm and 859 nm). At each plot in each year, we took four readings of ground reflectances at the above wavelengths. We also took readings in 12 (2015), 15 (2012, 2013, 2014), or 16 (2016) calibration plots adjacent to the experimental site, in which we harvested all aboveground biomass produced in the current year (we excluded litter and standing dead material), dried it to a constant weight at 60°C, and weighed it to estimate ANPP. We harvested at peak biomass each year, typically in late June.

For each plot and year, we averaged the four readings for each wavelength and then calculated a greenness index based on the same bands used to calculate NDVI using the MODIS (Moderate Resolution Imaging Spectroradiometer) and AVHRR (Advanced Very High Resolution Radiometer) bands for NDVI. We regressed the greenness index against the dry biomass weight from the ten calibration plots to convert the greenness index to ANPP. We fit regressions to a MODIS-based index and an AVHRR-based index for each year and retained the regression with the better fit based on R^2 values. We then predicted ANPP using the best regression equation for each year (Appendix 2). Our analysis, described below, can be done using NDVI directly, but we focus on ANPP to more easily relate our findings to previous work and because our results do not change if we analyze NDVI directly (Appendix 4).

¹Dates formatted as: mm-dd-yyyy.

Species composition data came from two sources: yearly census maps for each plot made using a pantograph (Hill, 1920) and yearly counts of annual species in each plot. From these sources, we determined the density of all annuals and perennials forbs, the basal cover of perennial grasses, and the canopy cover of shrubs. We made a large plot-treatment-year by species matrix, where columns were filled with either basal cover or density, depending on the measurement made for the particular species. We standardized the values in each column so we could directly compare species quantified with different metrics (density, basal cover, and canopy cover). This puts all abundance values on the same scale, meaning that common and rare species are weighted equally. Assuming that rare species will respond to treatments more than common ones, our approach is biased towards detecting compositional changes.

2.4 Data Analysis

2.4.1 Ecosystem functional response

Our main goal was to test whether the relationship between ANPP and soil moisture differed among the drought, control, and irrigation treatments. Based on our own observations and previous work at our study site (Blaisdell, 1958; Dalgleish et al., 2011; Adler et al., 2012), we chose to use cumulative volumetric water content from March through June as our metric of soil moisture (hereafter referred to as ‘VWC’). To achieve this goal, we fit a generalized linear mixed effects regression model with $\log(\text{ANPP})$ as the response variable and VWC and treatment as fixed effects. Plot and year of treatment were included as random effects to account for non-independence of observations, as described below. We log-transformed ANPP to account for heteroscedasticity. Both $\log(\text{ANPP})$ and VWC were standardized to have mean 0 and unit variance before fitting the model [i.e., $(x_i - \bar{x})/\sigma_x$].

Our model is defined as follows:

$$\mu_i = \beta \mathbf{x}_i + \gamma_{j(i)} \mathbf{z}_i + \eta_t, \quad (1)$$

$$\mathbf{y} \sim \text{Normal}(\boldsymbol{\mu}, \sigma^2), \quad (2)$$

where μ_i is the deterministic prediction from the regression model for observation i , which is associated with plot j and treatment year t . β is the vector of coefficients for the fixed effects in the design matrix \mathbf{X} . Each row of the design matrix represents a single observation (\mathbf{x}_i) and is a vector with the following elements: 1 for the intercept, a binary 0 or 1 if the treatment is “drought”, a binary 0 or 1 if the treatment is “irrigation”, the scaled value of VWC, binary “drought” value times VWC, and binary “irrigation” value times VWC. Thus, our model treats “control” observations as the main treatment and then estimates intercept and slope offsets for the “drought” and “irrigation” treatments. We use our model to test two statistical hypotheses based on the questions in our Introduction:

H1. The coefficient for drought \times VWC is positive and different from zero.

H2. The coefficient for irrigation \times VWC is negative and different from zero.

These hypotheses are based on evidence that precipitation-ANPP relationships often saturate with increasing precipitation (Hsu et al., 2012; Gherardi and Sala, 2015b).

We include two random effects to account for the fact that observations within plots and years are not independent. Specifically, we include plot-specific offsets (γ) for the intercept and slope terms and year-specific intercept offsets (η_t). The covariate vector \mathbf{z}_i for each observation i has two elements: a 1 for the intercept and the scaled value of VWC for that plot and year. The plot-specific coefficients are modeled hierarchically, where plot level coefficients are drawn from a multivariate normal distribution with mean 0 and a variance-covariance structure that allows the intercept and slope terms to be correlated:

$$\gamma_{j(i)} \sim \text{MVN}(0, \Sigma), \quad (3)$$

where Σ is the variance-covariance matrix and $j(i)$ reads as “plot j associated with observation i ”. The random year effects (η) are drawn from a normal prior with mean 0 and standard deviation σ_{year} , which was drawn from a half-Cauchy distribution. We used a vague prior distribution for each β : $\beta \sim \text{Normal}(0, 5)$. A full description of our model and associated R (R Core Team, 2016) code is in Appendix 3.

We fit the model using a Bayesian approach and obtained posterior estimates of all unknowns via the No-U-Turn Hamiltonian Monte Carlo sampler in Stan (Stan Development Team, 2016b). We used the R package ‘rstan’ (Stan Development Team, 2016a) to link R (R Core Team, 2016) to Stan. We obtained samples from the posterior distribution for all model parameters from four parallel MCMC chains run for 10,000 iterations, saving every 10th sample. Trace plots of all parameters were visually inspected to ensure well-mixed chains and convergence. We also made sure all scale reduction factors (\hat{R} values) were less than 1.1 (Gelman and Hill, 2009).

2.4.2 Sensitivity of ANPP over time

Unfortunately, our data does not allow us to include the effect of time on ecosystem functional response in the model described above. This is because we lack sufficient within-year and within-treatment variation of soil moisture (i.e., within a year and treatment each plot shares the same value of VWC). Therefore, we conducted an independent analysis to determine if the sensitivity of ANPP to treatment changed over time.

We define ‘sensitivity’ as $\frac{ANPP_{control} - ANPP_{treatment}}{VWC_{control} - VWC_{treatment}}$, following Wilcox et al. (2017). This metric of sensitivity measures the response of ANPP standardized by the magnitude of the treatment. Because our treatment and control plots are not paired, we conducted this analysis by comparing each treatment plot to the mean of the control plots in each year. Then, independently for each treatment, we regressed sensitivity against year of treatment using the `lm()` function in R (R Core Team, 2016). This analysis also allows us to link particularly sensitive treatment years to changes in community composition.

2.4.3 Community composition over time

We used nonmetric multidimensional scaling (NMDS) based on Bray-Curtis distances to identify temporal changes in community composition among treatments. We first calculated Bray-Curtis distances among all plots for each year of the experiment and then extracted those distances for use in the NMDS. Some values of standardized species’ abundances were negative, which is not allowed for calculating Bray-Curtis distances. We simply added ‘2’ to each abundance value to ensure all values were greater than zero. We plotted the first two axes of NMDS scores to see if community composition overlapped, or not, among treatments in each year. We used the `vegan::metaMDS()` function (Oksanen, 2016) to calculate Bray-Curtis distances and then to run the NMDS analysis. We used the `vegan::adonis()` function (Oksanen, 2016) to perform permutational multivariate analysis of variance to test whether treatment plots formed distinct groupings. To test whether treatment plots were equally dispersed, or not, we used the `vegan::betadisper()` function (Oksanen, 2016).

We conducted the above analysis for all species, and then conducted a separate analysis for annual species only (Appendix 1). Annual species have shorter life spans, so conducting a separate analysis allowed us to discover if community responses are restricted to short-lived species that might respond more quickly to altered precipitation regimes. Given the dominance of perennial species in our system (Appendix 1), shifts in the annual plant community could be difficult to detect in the analysis of the full community.

2.4.4 Reproducibility

All R code and data necessary to reproduce our analysis has been archived on Figshare (*link here after acceptance*) and released on GitHub (https://github.com/atredennick/usses_water/releases/v0.1). We also include annotated Stan code in our model description in Appendix 3.

3 RESULTS

Ambient precipitation was variable over the five years of the study (Fig. 2A), meaning our experiment captured a range of environmental conditions. ANPP varied from a minimum of 74.5 g m⁻² in 2014 to a maximum of 237.1 g m⁻² in 2016 when averaged across treatments (Fig. 2C). ANPP was slightly higher in irrigation plots (on average 17% higher) and slightly lower in drought plots (on average 23% lower) relative to control plots (Fig. 2C), corresponding to estimated soil volumetric water content (VWC)

differences among treatments (Fig. 2B). VWC in drought plots was 12% less than in control plots on average, and VWC in irrigation plots was 19% higher than in control plots on average. Such differences in soil VWC indicate our treatment infrastructure was successful. ANPP was highly variable across plots within years, as indicated by the large and overlapping standard deviations (Fig. 2C).

Cumulative March-June soil moisture had a weak positive effect on ANPP (Table 1; Fig. 3B). The effect of soil moisture for each treatment is associated with high uncertainty, however, with 95% Bayesian credible intervals that broadly overlap zero (Table 1). Although the parameter estimates for the effect of soil moisture overlap zero, the posterior distributions of the slopes all shrank and shifted to more positive values relative to the prior distributions (Fig. A3-2), which indicates the data did influence parameter estimates beyond the information from the uninformative priors. Ecosystem functional response was similar among treatments (Table 1; Fig. 3B), but there is evidence that the slopes for the drought and irrigation treatments are less than the slope for the control treatment. This evidence comes from interpreting the posterior distributions of the slope offsets for the treatments. From these distributions, we calculate a 91% one-tailed probability that the estimate is less than zero for the irrigation treatment and a 96% one-tailed probability that the estimate is less than zero for the drought treatment (Fig. 3A, right panel).

Sensitivity of ANPP to irrigation was constant over the course of our experiment (Fig. 3C). Sensitivity of ANPP to drought, however, grew over time ($P = 0.0002$; Fig. 3C).

Community composition was similar among treatments. The multidimensional space of community composition overlapped among treatments in all years and was equally dispersed in all years (Table 2; Fig. 4). Community composition was also remarkably stable over time, with no evidence of divergence among treatments (Table 2; Fig. 4). There were also no changes in the dominant species over time in any treatment (Appendix 1). Splitting the analysis apart for annuals led to similar results. There is some evidence that the annual community changed in response to treatment in two years (Fig. A1-6), but these responses appear to come from very rare species since our analysis weights them equally (Fig. A1-4). By definition, these rare species contribute little to ANPP and thus any compositional changes have little influence on ecosystem functional response.

4 DISCUSSION

Ecosystem response to a new precipitation regime depends on the physiological responses of constituent species and the rate at which community composition shifts to favor species better able to take advantage of, or cope with, new resource levels (Smith et al., 2009). Previous work has shown that community compositional shifts can be both rapid, on the order of years (Hoover et al., 2014), and slow, on order of decades (Knapp et al., 2012; Wilcox et al., 2016). A lingering question is how the time scales of ecosystem response and community change vary among ecosystems. Precipitation manipulation experiments can help answer this question, especially if they push water availability outside the historical range of variability for long periods.

The results of our five year experiment in a sagebrush steppe conforms to one of our four predictions. Ecosystem functional response under chronic drought and irrigation was different from the control treatment, but community composition remained unchanged (Fig. 3A, Fig. 4), representing the top right scenario in Fig. 1. The decrease in the slope of the VWC \times productivity relationship in the irrigated plots is consistent with a weaker response of productivity to additional water in this ecosystem. This suggests that if average soil moisture were pushed consistently higher than currently observed ambient conditions, there would be a weaker relationship between precipitation and productivity in this system.

The decrease in the slope of the VWC \times productivity relationship in the droughted plots is puzzling because it is counter to our expectation that the slope would be higher than the control plots. However, we believe the significance of the slope offsets has more to do with noise than signal. This is because the full distributions of the slopes of the VWC \times productivity relationship, not just the offsets, were similar among treatments (Table 1). We therefore conclude that ecosystem functional response is consistent (similar values) and weak (all broadly overlapping zero) across all precipitation treatments.

The similarity of ecosystem functional response (Fig. 3) and community composition (Fig. 4) among treatments is surprising because grasslands generally, and sagebrush steppe specifically, are considered water-limited systems. For example, Huxman et al. (2004) and Knapp et al. (2015) showed that semi-arid sites are more sensitive to drought than mesic sites, and Wilcox et al. (2017) found that semi-arid sites are

341 particularly sensitive to irrigation treatments. Based on these findings, we expected ecosystem functional
342 response, community composition, or both to change under our treatments. Why did our treatments fail
343 to induce ecosystem or community responses? We can think of three reasons. Two are limitations of
344 our study, and one is the life history traits of the species in our focal communities. We first discuss the
345 potential limitations of our study, and then discuss the biological explanation.

346 First, it could be that our precipitation manipulations were not large enough to induce a response. That
347 is, a 50% decrease in any given year may not be abnormal given our site's historical range of variability
348 (Knapp et al., 2017). We cannot definitively rule out this possibility, but we have reason to believe our
349 treatments *should* have been large enough. Using the methods described by Lemoine et al. (2016), we
350 calculated the percent reduction and increase of mean growing season precipitation necessary to reach the
351 1% and 99% extremes of the historical precipitation regime at our site (Fig. A5-1). The 1% quantile of
352 precipitation at our site is 110 mm, a 47% reduction from the mean, and the 99% quantile is 414 mm,
353 a 77% increase from mean growing season precipitation (Appendix 5). Thus, our drought treatment
354 represented extreme precipitation amounts, especially in years where ambient precipitation was below
355 average (Fig. 2A). The irrigation treatment may not qualify as extreme, yet that is the treatment where we
356 did observe an effect (Fig. 3A).

357 Second, ANPP at our site may be influenced by additional factors, not only the cumulative March-June
358 soil moisture covariate we included in our statistical model. For example, La Pierre et al. (2016) found
359 that site-scale ANPP is better predicted by abiotic factors such as nutrient availability than precipitation.
360 Moreover, temperature can impact ANPP directly (Epstein et al., 1997) and by exacerbating the effects
361 of soil moisture (De Boeck et al., 2011). Measurements of soil moisture likely contain a signal of
362 temperature, through its impact on evaporation and infiltration, but the measurements will not capture
363 the direct effect of temperature on metabolic and physiological processes. We also did not redistribute
364 snow across our treatments in the winter, and snow melt may spur early spring growth. Failure to account
365 for potentially important covariates could explain the within-year spread of ANPP (Fig 2C, Fig. 3B) and
366 the resulting uncertain relationship we observed between soil moisture and ANPP across all treatments
367 (Table 1, Fig. A3-2). Unfortunately, we did not collect data on additional factors because our focus was
368 on precipitation.

369 Third, the life history traits of the dominant species in our study ecosystem may explain the consistently
370 positive, but weak and uncertain, effect of soil moisture on ANPP (Table 1, Fig. 3). Species that live
371 in variable environments, such as cold deserts, must have strategies to ensure long-term success as
372 conditions vary. One strategy is bet hedging, where species forego short-term gains to reduce the variance
373 of long-term success (Seger, 1987). In other words, species follow the same conservative strategy every
374 year, designed to minimize response to environmental conditions. The dry and variable environment of
375 the sagebrush steppe has likely selected for bet hedging species that can maintain function at low water
376 availability and have weak responses to high water availability. In so doing, the dominant species in our
377 ecosystem avoid “boom and bust” cycles, which corresponds to the weak effect of soil moisture on ANPP
378 (i.e., the Bayesian credible intervals for the slopes overlapping zero; Table 1).

379 Another strategy to deal with variable environmental conditions is avoidance, which would also result
380 in a consistent ecosystem functional response between drought and control treatments. For example,
381 many of the perennial grasses in our focal ecosystem avoid drought stress by growing early in the growing
382 season (Blaisdell, 1958, A.R. Kleinhesselink, personal observation). Furthermore, the dominant shrub in
383 our focal ecosystem, *Artemisia tripartita*, has access to water deep in the soil profile thanks to a deep
384 root system (Kulmatiski et al., 2017). The dominance of our site by the shrub *A. tripartita* (Appendix 1)
385 may explain why our results do not conform to broader patterns of grassland sensitivity to precipitation
386 manipulations (e.g., Huxman et al., 2004; Knapp et al., 2015; Wilcox et al., 2017).

387 We did find that ANPP became more sensitive to drought over time (Fig. 3C). The latter years of
388 our experiment were relatively wet, meaning that the drought treatment could have the largest absolute
389 reduction of soil moisture in those years. But our metric of sensitivity accounts for differences among years
390 because ANPP differences is standardized by the difference in soil moisture within the year. Therefore, it
391 is likely that the increased sensitivity to drought over time is due to the cumulative impact of drought on
392 the dominant species. Ecosystem functional response may soon shift if sensitivity continues to increase.

393 5 CONCLUSIONS

394 Our results suggest the species in our focal plant community are insensitive to to changes in precipitation
395 regime, [at least over the five years of our experiment](#). Such insensitivity can buffer species against
396 precipitation variability in this semi-arid ecosystem, making them successful in the long run. Longer,
397 chronic precipitation alteration might reveal plant community shifts that we did not observe (e.g., Wilcox
398 et al., 2016), [and the increased sensitivity of ANPP to drought in the final years of our experiment may](#)
399 [portend such a shift](#). Likewise, a long-term increase in water availability could allow species that do not bet
400 hedge to gain prominence and dominate the ecosystem functional response. [Time *per se* may be especially](#)
401 [relevant in our focal ecosystem where the perennial species are long-lived, meaning compositional](#)
402 [turnover may take many more years than we report on here](#). Our results suggest compositional shifts
403 would have the largest impact at high rainfall because the current community maintained consistent
404 ecosystem functional response at very low water availability. [However, the increased sensitivity of ANPP](#)
405 [to drought conditions suggests a shift in ecosystem functional response may be near, and simply takes](#)
406 [more time than we observed here](#).

407 6 ACKNOWLEDGEMENTS

408 We thank the many summer research technicians who collected the data reported in this paper and the
409 USDA, U.S. Sheep Experiment Station (Dubois, ID) for facilitating work on their property. We also
410 thank Susan Durham for clarifying our thinking on the statistical analyses and Kevin Wilcox for helpful
411 discussions on analyzing community composition data. [Two anonymous reviewers and Elsie Denton](#)
412 [provided thoughtful reviews that improved our paper](#).

413 7 FUNDING

414 This research was supported by the Utah Agricultural Experiment Station, Utah State University, and
415 approved as journal paper number 9035. The research was also supported by the National Science
416 Foundation, through a Postdoctoral Research Fellowship in Biology and Mathematics to ATT (DBI-
417 1400370), a Graduate Research Fellowship to ARK, and grants DEB-1353078 and DEB-1054040 to
418 PBA.

419 8 AUTHOR CONTRIBUTIONS

- 420 • Andrew T. Tredennick collected data, analyzed the data, wrote the paper, prepared figures and/or
421 tables, reviewed drafts of the paper.
- 422 • Andrew R. Kleinhesselink conceived and designed the experiments, performed the experiments,
423 collected data, analyzed the data, reviewed drafts of the paper.
- 424 • J. Bret Taylor contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- 425 • Peter B. Adler conceived and designed the experiments, performed the experiments, collected data,
426 analyzed the data, reviewed drafts of the paper.

427 9 SUPPLEMENTAL INFORMATION

428 **Appendix 1.** [Additional information on plant community composition and dynamics, Table A1-1, Table](#)
429 [A1-2, Table A1-3, Fig. A1-1, Fig. A1-2, Fig. A1-3, Fig. A1-4, Fig. A1-5, and Fig. A1-6.](#)

430 **Appendix 2.** Additional methods and information on estimating aboveground net primary productivity,
431 Table A2-1, and Table A2-2.

432 **Appendix 3.** Details of the hierarchical Bayesian model, Fig. A2-1, Fig. A2-2, and Fig. A2-3.

433 **Appendix 4.** [Results from analysis of NDVI without conversion to ANPP, Fig. A3-1.](#)

434 **Appendix 5.** Details on analysis of precipitation historical range of variability and Fig. A4-1.

10 TABLES

Table 1. Summary statistics from the posterior distributions of coefficients for each treatment (β coefficients in equation 1). The ‘Intercept’ and ‘Slope’ summaries reported here for drought and irrigation are from the posterior distributions of the intercept and slope for the control treatment plus the offsets for each treatment. Posterior distributions of the offsets are in Figure 3A.

Coefficient	Treatment	Posterior Mean	Posterior Median	Lower 95% BCI	Upper 95% BCI
Intercept	Control	0.16	0.14	-1.22	1.69
Intercept	Drought	0.13	0.05	-1.69	2.39
Intercept	Irrigation	-0.24	-0.18	-2.11	1.53
Slope	Control	0.94	0.86	-0.51	2.69
Slope	Drought	0.65	0.57	-0.99	2.62
Slope	Irrigation	0.72	0.65	-0.54	2.26

Table 2. Results from statistical tests for clustering and dispersion of community composition among precipitation treatments. ‘adonis’ tests whether treatments form unique clusters in multidimensional space; ‘betadisper’ tests whether treatments have similar dispersion. For both tests, P values greater than 0.05 indicate there is no support that the treatments differ.

Year	Test	n	d.f.	F	P
2011	adonis	21	2	1.02	0.45
2011	betadisper	21	2	2.22	0.14
2012	adonis	22	2	1.09	0.34
2012	betadisper	22	2	0.21	0.81
2013	adonis	22	2	1.25	0.13
2013	betadisper	22	2	0.44	0.65
2014	adonis	22	2	0.95	0.56
2014	betadisper	22	2	0.35	0.71
2015	adonis	21	2	1.05	0.39
2015	betadisper	21	2	3.01	0.07
2016	adonis	21	2	1.07	0.36
2016	betadisper	21	2	0.49	0.62




Figure 1. Possible outcomes of chronic resource alteration based on the 'Hierarchical Response Framework' (Smith et al. 2009).




Figure 2. (A) Probability density of historical precipitation from 1926-2016, with the years of the experiment shown with arrows on the x -axis. (B) Estimated daily soil volumetric water content (VWC) in each of the three treatments during the course of the experiment. (C) Mean (points) ANPP and its standard deviation (error bars) for each year of the experiment. Colors in panels B and C identify the treatment, as specified in the legend of panel C.




Figure 3. Results from the generalized linear mixed effects model (A-B) and the sensitivity analysis (C). (A) Posterior distributions for the effects of drought and irrigation on the intercept and slope of the productivity-soil moisture relationship. Treatment effects show the difference between the coefficients estimated in the treated plots and the control plots. Probabilities (“Pr ≤ 0 =”) for each coefficient indicate the one-tailed probability that the coefficient is less than or greater than zero, depending on whether the median of the distribution is less than or greater than zero. The posterior densities were smoothed for visual clarity by increasing kernel bandwidth by a factor of five. (B) Scatterplot of the data and model estimates shown as solid lines. Model estimates come from treatment level coefficients (colored lines). Note that we show log(ANPP) on the y-axis of panel B; this same plot can be seen on the arithmetic scale in supporting material Fig. A2-1. (C) Regression of sensitivity against time for each treatment. Each point represents the sensitivity of ANPP in a plot relative to the mean of the control plots in that year. Only the significant regression for the drought treatment is plotted ($P = 0.0002$).




Figure 4. Nonmetric multidimensional scaling scores representing plant communities in each plot, colored by treatment. *Each point represents a plot and the axes represent species composition.*

References

- Adler, P. B., Dalgleish, H. J., and Ellner, S. P. (2012). Forecasting plant community impacts of climate variability and change: when do competitive interactions matter? *Journal of Ecology*, 100:478–487.
- Avolio, M. L., Pierre, K. J. L., Houseman, G. R., Koerner, S. E., Grman, E., Isbell, F., Johnson, D. S., and Wilcox, K. R. (2015). A framework for quantifying the magnitude and variability of community responses to global change drivers. *Ecosphere*, 6(12):1–14.
- Bates, D., Maechler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4.
- Blaisdell, J. P. (1958). Seasonal development and yield of native plants on the upper snake river plains and their relation to certain climate factors. *United States Department of Agriculture Technical Bulletin No. 1190*.
- Byrne, K. M., Lauenroth, W. K., Adler, P. B., and Byrne, C. M. (2011). Estimating Aboveground Net Primary Production in Grasslands: A Comparison of Nondestructive Methods. *Rangeland Ecology and Management*, 64(5):498–505.
- Dalgleish, H. J., Koons, D. N., Hooten, M. B., Moffet, C. A., and Adler, P. B. (2011). Climate influences the demography of three dominant sagebrush steppe plants. *Ecology*, 92(1):75–85.
- De Boeck, H. J., Dreesen, F. E., Janssens, I. A., and Nijs, I. (2011). Whole-system responses of experimental plant communities to climate extremes imposed in different seasons. *New Phytologist*, 189(3):806–817.
- Epstein, H. E., Lauenroth, W. K., and Burke, I. C. (1997). Effects of temperature and soil texture on ANPP in the U.S. Great plains. *Ecology*, 78(8):2628–2631.
- Gelman, A. and Hill, J. (2009). *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press, Cambridge.
- Gherardi, L. A. and Sala, O. E. (2013). Automated rainfall manipulation system: a reliable and inexpensive tool for ecologists. *Ecosphere*, 4(2):1–10.
- Gherardi, L. A. and Sala, O. E. (2015a). Enhanced interannual precipitation variability increases plant functional diversity that in turn ameliorates negative impact on productivity. *Ecology Letters*, 18(12):1293–1300.
- Gherardi, L. A. and Sala, O. E. (2015b). Enhanced precipitation variability decreases grass- and increases shrub-productivity. *Proceedings of the National Academy of Sciences*, 112(41):12735–12740.
- Hill, R. R. (1920). Charting Quadrats with a Pantograph. *Ecology*, 1(4):270–273.
- Hoover, D. L., Knapp, A. K., and Smith, M. D. (2014). Resistance and resilience of a grassland ecosystem to climate extremes. *Ecology*, 95(9):2646–2656.
- Hsu, J. S., Powell, J., and Adler, P. B. (2012). Sensitivity of mean annual primary production to precipitation. *Global Change Biology*, 18(7):2246–2255.
- Huxman, T. E., Smith, M. D., Fay, P. A., Knapp, A. K., Shaw, M. R., Loik, M. E., Smith, S. D., Tissue, D. T., Zak, J. C., Weltzin, J. F., Pockman, W. T., Sala, O. E., Haddad, B. M., Harte, J., Koch, G. W., Schwinning, S., Small, E. E., and Williams, D. G. (2004). Convergence across biomes to a common rain-use efficiency. *Nature*, 429(6992):651–654.
- Kleinhesselink, A. R. (2017). *Direct and indirect effects of climate change on plant populations and communities in sagebrush steppe*. Dissertation, Utah State University.
- Knapp, A. K., Avolio, M. L., Beier, C., Carroll, C. J. W., Collins, S. L., Dukes, J. S., Fraser, L. H., Griffin-Nolan, R. J., Hoover, D. L., Jentsch, A., Loik, M. E., Phillips, R. P., Post, A. K., Sala, O. E., Slette, I. J., Yahdjian, L., and Smith, M. D. (2017). Pushing precipitation to the extremes in distributed experiments: recommendations for simulating wet and dry years. *Global Change Biology*, 23(5):1774–1782.
- Knapp, A. K., Briggs, J. M., and Smith, M. D. (2012). Community stability does not preclude ecosystem sensitivity to chronic resource alteration. *Functional Ecology*, 26(6):1231–1233.
- Knapp, A. K., Carroll, C. J., Denton, E. M., La Pierre, K. J., Collins, S. L., and Smith, M. D. (2015). Differential sensitivity to regional-scale drought in six central US grasslands. *Oecologia*, 177(4):949–957.
- Kulmatiski, A., Adler, P. B., Stark, J. M., and Tredennick, A. T. (2017). Water and nitrogen uptake are better associated with resource availability than root biomass. *Ecosphere*, 8(3).
- La Pierre, K. J., Blumenthal, D. M., Brown, C. S., Klein, J. A., and Smith, M. D. (2016). Drivers of Variation in Aboveground Net Primary Productivity and Plant Community Composition Differ Across

491 a Broad Precipitation Gradient. *Ecosystems*, 19(3):521–533.

492 Lemoine, N. P., Sheffield, J., Dukes, J. S., Knapp, A. K., and Smith, M. D. (2016). Terrestrial Precipitation
 493 Analysis (TPA): A resource for characterizing long-term precipitation regimes and extremes. *Methods*
 494 *in Ecology and Evolution*, 7(11):1396–1401.

495 Oksanen, J. (2016). *Vegan: ecological diversity*.

496 Peters, D. P. C., Yao, J., Sala, O. E., and Anderson, J. P. (2012). Directional climate change and potential
 497 reversal of desertification in arid and semiarid ecosystems. *Global Change Biology*, 18(1):151–163.

498 R Core Team (2016). *R: A Language and Environment for Statistical Computing*. Vienna, Austria.

499 Seger, J. (1987). What is bet-hedging? In Harvey, P. and Partridge, L., editors, *Oxford surveys in*
 500 *evolutionary biology*, pages 182–211. Oxford University Press, Oxford.

501 Smith, M. (2011). An ecological perspective on extreme climatic events: A synthetic definition and
 502 framework to guide future research. *Journal of Ecology*, 99(3):656–663.

503 Smith, M. D., Knapp, A. K., and Collins, S. L. (2009). A framework for assessing ecosystem dynamics in
 504 response to chronic resource alterations induced by global change. *Ecology*, 90(12):3279–3289.

505 Stan Development Team (2016a). Rstan: the R interface to Stan, Version 2.14.1.

506 Stan Development Team (2016b). Stan: A C++ Library for Probability and Sampling, Version 2.14.1.

507 Wilcox, K. R., Blair, J. M., Smith, M. D., and Knapp, A. K. (2016). Does ecosystem sensitivity to
 508 precipitation at the site-level conform to regional-scale predictions? *Ecology*, 97(3):561–568.

509 Wilcox, K. R., Shi, Z., Gherardi, L. A., Lemoine, N. P., Koerner, S. E., Hoover, D. L., Bork, E.,
 510 Byrne, K. M., Cahill, J., Collins, S. L., Evans, S., Gilgen, A. K., Holub, P., Jiang, L., Knapp, A. K.,
 511 LeCain, D., Liang, J., Garcia-Palacios, P., Peñuelas, J., Pockman, W. T., Smith, M. D., Sun, S., White,
 512 S. R., Yahdjian, L., Zhu, K., and Luo, Y. (2017). Asymmetric responses of primary productivity
 513 to precipitation extremes: A synthesis of grassland precipitation manipulation experiments. *Global*
 514 *Change Biology*, 23(10):4376–4385.