

# Regression versus State-Space

## FISH 507 – Applied Time Series Analysis

Eli Holmes

2 Mar 2021

## Fixed & random effects

Let's go back to Mark's lecture on DFA

$$y_t = \underbrace{\alpha + \beta x_t}_{\text{fixed}} + \underbrace{f_t + e_t}_{\text{random}}$$

Regression (linear or non-linear) with correlated errors gets your the fixed effects while properly taking into account the correlated random effects.

State-space model allows you to model the  $f_t$ .

# Things to think about

$$y_t = \underbrace{\alpha + \beta x_t}_{\text{fixed}} + \underbrace{f_t + e_t}_{\text{random}}$$

- ▶ What if you want  $f_t$ , the hidden random walk
- ▶ What if you want  $E[\alpha + \beta x_t + f_t]$
- ▶ What if you want the  $E[y_t | y_{1:t-1}]$
- ▶ What if we want to forecast  $y_t$ ?

Let's simulate some data



$$x_t = t$$

$$f_t = \phi f_{t-1} + w_t, w_t \sim N(0, 1)$$

$$y_t = \beta x_t + f_t + v_t, v_t \sim N(0, \sqrt{0.2})$$

# Simulated data

```
set.seed(123)
N <- 100; h <- 100; x <- 1:N
sim1 <- sim.data(N, h=h, phi=0.8)
sim2 <- sim.data(N, h=h, phi=1)
```



## Fit with arima(y, xreg=x)



Fits a linear regression with ARMA errors.

```
library(forecast)
fit <- auto.arima(sim1$y, xreg=x)
fr1 <- forecast(fit, xreg=(N+1):(N+h))
fit <- auto.arima(sim2$y, xreg=x)
fr2 <- forecast(fit, xreg=(N+1):(N+h))
```

## Plot forecasts versus true

I created simulations from the true process to get truth.

Plot shows prediction intervals (future  $y$ ). Red lines are true 80% intervals.




## Fitted

You need to be careful to think about what you mean by `fitted()`


- ▶ `glS(y~x)` and similar would return  $E[\alpha + \beta x_t]$
- ▶ `arima(y, xreg=x)` returns  $E[\alpha + \beta x_t + e_t | y_{1:t-1}]$
- ▶ state-space model would get you  $E[\alpha + \beta x_t + f_t | y_{1:N}]$



**gls fitted  
fixed effects**



**arima fitted  
 $E[y|y(1:t-1)]$**



# Regression w ARMA errors vs ARMAX

These are different models.

Regression w AR1 errors

$$e_t = \phi e_{t-1} + w_t$$

$$y_t = \alpha + \beta x_t + e_t$$

ARMAX: In this case, AR(1)-X

$$y_t = \phi y_{t-1} + \alpha + \beta x_t + w_t$$