

Is Achieving Carbon Neutrality Possible in Healthcare? Comparing the Environmental Costs of Differentiated Service Delivery of Antiretroviral Therapy for People Living with HIV in Rural South Africa

Ashley S. Tseng^{1,2}, Adam A. Szpiro³, Jesse Heitner⁴, Alastair van Heerden^{5,6}, Xolani Ntinga⁵, Meighan L. Krows², Torin T. Schaafsma², Ruanne V. Barnabas^{4,7}

1 Department of Epidemiology, University of Washington, Seattle, Washington, USA

2 Department of Global Health, University of Washington, Seattle, Washington, USA

3 Department of Biostatistics, University of Washington, Seattle, Washington, USA

4 Division of Infectious Diseases, Massachusetts General Hospital, Boston, Massachusetts, USA

5 Center for Community Based Research, Human Sciences Research Council, Sweetwaters, KwaZulu-Natal, South Africa

6 South African Medical Research Council/Wits Developmental Pathways for Health Research Unit, Department of Paediatrics, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, Gauteng, South Africa

7 Department of Medicine, Harvard Medical School, Boston, Massachusetts, USA

Background:

- People living with HIV require reliable access to antiretroviral therapy (ART) to maintain viral suppression
- Standard-of-care for people living with HIV in South Africa to refill ART is going in-person to clinics
- Retention in HIV care is a complex public health problem
- Differentiated service delivery: client-centered approach that adapts HIV services across the health care cascade to reflect the preferences, expectations, and needs of people living with HIV while reducing unnecessary burdens on the health system
- The Deliver Health Study (2019-20): Adults living with HIV in rural KwaZulu-Natal, South Africa were enrolled and randomized to receive standard clinic-based care or 3-month ART refills and monitoring at home
- Proportion of clients who were virally suppressed in the Deliver Health Study was **significantly higher among those who received home-delivered ART refills** vs. clinic-based care (88% vs. 74%; RR=1.21, 95% CI: 1.02-1.42)
- Home delivery of ART and monitoring is **effective** for achieving viral suppression and retention in HIV care, but what is the environmental impact of scaling up home-delivered ART in high HIV prevalence settings such as rural South Africa? **Could scaling up home delivery support a carbon-neutral approach by reducing overall greenhouse gas emissions from individual commuters to clinics?**

Objective: To analyze carbon dioxide (CO₂) emissions data to compare environmental costs of different ART refill methods for people living with HIV.

Methods:

- Compared cumulative CO₂ emissions for quarterly visits over 12 months for participants receiving home-delivered vs. clinic-based refills
- Six- and 12-month refill scenarios were evaluated
- Per person living with HIV, delivery log data was used to:
 - Calculate total distance travelled (km)
 - Estimate cumulative CO₂ emissions (kgCO₂)

Table 1. Estimated CO₂ emissions per km travelled by transport method to ART refill visits

Refill Method	Transport Method	Estimated CO ₂ Emissions (kgCO ₂ /km)
Clinic	Walking	0
	Bicycling	0
	Driving (personal vehicle)	0.148
	Public Transit/Taxi (Toyota Quantum Sefikile bus)	0.339
Home delivery	Delivery vehicle (2016 Ford Ranger 2.2L Diesel Double Cab 4x4)	0.185

Two outcomes:

- 1) Incremental cost-effectiveness ratio (ICER):

$$\frac{(\text{Avg. total CO}_2 \text{ emissions}_{\text{home delivery group}}) - (\text{Avg. total CO}_2 \text{ emissions}_{\text{clinic group}})}{\left(\frac{n_{\text{virally suppressed in home delivery group}}}{n_{\text{home delivery group}}}\right) - \left(\frac{n_{\text{virally suppressed in clinic group}}}{n_{\text{clinic group}}}\right)}$$

- 2) Comparative cost-effectiveness (CCE):

$$\left(\frac{\text{Avg. total CO}_2 \text{ emissions}_{\text{home delivery group}} * n_{\text{home delivery group}}}{n_{\text{virally suppressed in home delivery group}}}\right) - \left(\frac{\text{Avg. total CO}_2 \text{ emissions}_{\text{clinic group}} * n_{\text{clinic group}}}{n_{\text{virally suppressed in clinic group}}}\right)$$

Results

- Of those who knew their HIV status at baseline (n=103), 3% had taken ART in the past and stopped taking them and 10% took a break from taking ART at some point
 - Reasons for discontinued ART use included long clinic waiting times and clinics being located too far away
- Walking was the most common (64%) mode of transportation to ART refill visits for participants the clinic group, followed by minibuss (33%), then driving (3%)
- **ICER:** Incremental carbon-cost was 50 kg of CO₂ per person virally suppressed through home delivery, equivalent to driving 271 extra km in the delivery vehicle

- If 6- or 12-month refills were provided instead, incremental emissions for home-delivered ART would be 25 and 13 kg of CO₂, equivalent to 136 and 68 km driven, respectively
- **CCE:** Compared to the clinic group, home delivery cost an extra 7.8 kg of CO₂ emissions per person virally suppressed

Table 2. Characteristics of people living with HIV in the Deliver Health Study

Characteristic	Clinic (N=73)	Home Delivery (N=80)	Total (N=153)
	mean (SD) or n (%)		
Age (years)	35.6 (8.5)	38.8 (9.3)	37.3 (9.1)
Men	38 (52.1)	44 (55.0)	82 (53.6)
Marital status			
Single	38 (52.1)	40 (50.0)	78 (51.0)
Member of an unmarried couple	31 (42.5)	29 (36.3)	60 (39.2)
Married	0 (0)	6 (7.5)	6 (3.9)
Divorced or widowed	4 (5.5)	5 (6.3)	9 (5.9)
Educational attainment			
Primary	23 (31.5)	15 (18.8)	38 (24.8)
Secondary	40 (54.8)	57 (71.3)	97 (63.4)
Tertiary and above	10 (13.7)	8 (10.0)	18 (11.8)
Employed	26 (35.6)	33 (41.3)	59 (38.6)
Individuals known to be living with HIV	51 (69.9)	52 (65.0)	103 (67.3)
Past ART use (among individuals known to be living with HIV, n=103)			
Currently on ART	48 (65.8)	50 (62.5)	98 (64.1)
Taken ART in the past	2 (2.7)	1 (1.3)	3 (2.0)
Never taken ART	23 (31.5)	29 (36.3)	52 (34.0)
Virally suppressed (<20 copies per mL) at month 12	54 (74.0)	70 (87.5)	124 (81.0)
Total distance travelled to refill visit per participant (km)	7.3 (13.3)	6.3 (5.1)	6.8 (9.8)
Cumulative CO₂ emissions across all refill visits over study follow-up per participant (kgCO₂)	0.4 (0.6)	7.3 (8.2)	0.4 (0.7)

Conclusions

In rural South Africa, incremental CO₂ emissions were higher for participants receiving home-delivered vs. clinic-based ART refills but could be offset by 6- or 12-month refills and/or changing number of deliveries or vehicle type. Other research studies should consider quantifying environmental impacts of their work alongside the public health impacts.

What's next?

We are collecting data in an ongoing randomized controlled trial, the SMART ART Study, in KwaZulu-Natal, South Africa. We will analyze CO₂ data from the SMART ART Study along with the Deliver Health Study.

Funding

The Deliver Health Study was funded by the U.S. National Institute of Mental Health (NIMH; R21MH115770). AST received support through an administrative diversity supplement from NIMH (R01MH124465-02S1) and a Department of Epidemiology Travel Award. Collaborating authors are supported through the SMART ART Study, a NIMH-funded study (R01MH124465; co-PIs: RVB and AVH). The funder had no role in study design, data collection, data analysis, or writing of this work.

Acknowledgements

Thank you to all Deliver Health Study participants and the communities that supported the work, and to the Deliver Health Study team for all their hard work.

Contact

Ashley S. Tseng, PhD(c), MPH, BSc: atseng23@uw.edu

