



Master's thesis
Astrophysical Sciences

Supermassive black holes and the cosmological formation of massive early-type galaxies (title not final)

Atte Keitaanranta

June 3, 2021

Supervisor(s): Prof. Peter Johansson
M.Sc. Matias Mannerkoski

Censor(s): Prof. Peter Johansson

UNIVERSITY OF HELSINKI
DEPARTMENT OF PHYSICS

PL 64 (Gustaf Hållströmin katu 2)
00014 University of Helsinki

| | | | |
|--|--|--|---|
| Tiedekunta — Fakultet — Faculty | | Koulutusohjelma — Utbildningsprogram — Education programme | |
| Faculty of Science | | Department of Physics | |
| Tekijä — Författare — Author | | | |
| Atte Keitaanranta | | | |
| Työn nimi — Arbetets titel — Title | | | |
| Supermassive black holes and the cosmological formation of massive early-type galaxies (title not final) | | | |
| Opintosuunta — Studieriktning — Study track | | | |
| Astrophysical Sciences | | | |
| Työn laji — Arbetets art — Level | | Aika — Datum — Month and year | Sivumäärä — Sidoantal — Number of pages |
| Master's thesis | | June 3, 2021 | 0 pages |
| Tiivistelmä — Referat — Abstract | | | |
| Abstract goes here. | | | |
| Avainsanat — Nyckelord — Keywords | | | |
| Your keywords here | | | |
| Säilytyspaikka — Förvaringsställe — Where deposited | | | |
| Muita tietoja — övriga uppgifter — Additional information | | | |

Contents

| | | |
|----------|---|----------|
| 1 | Introduction | 1 |
| 1.1 | Information about galaxies, shortly | 1 |
| 1.2 | Aim of the thesis | 1 |
| 2 | Background | 2 |
| 2.1 | Cosmology | 2 |
| 2.1.1 | Hubble parameter, Friedmann equations and so on | 2 |
| 2.1.2 | Cosmological perturbations | 2 |
| 2.2 | Early-type galaxies | 2 |
| 2.2.1 | Types of ellipticals | 2 |
| 2.2.2 | Photometric and kinematic profiles | 2 |
| 2.3 | Feedback processes | 2 |
| 3 | GADGET-3 and KETJU | 3 |
| 3.1 | Overview of GADGET-3 | 3 |
| 3.2 | Smoothed Particle Hydrodynamics | 3 |
| 3.3 | Gas cooling? | 3 |
| 3.4 | Feedback? | 3 |
| 3.5 | KETJU | 3 |
| 4 | Creating initial conditions for the cosmological zoom-in simula- | |

| | |
|---|-----------|
| tions | 4 |
| 4.1 Zoom-in technique | 4 |
| 4.2 MUSIC | 4 |
| 4.2.1 Basic properties of MUSIC | 4 |
| 4.2.2 Algorithms of MUSIC | 4 |
| 4.2.3 Creating an IC-file with a zoom-in box | 4 |
| 4.3 GADGET-3 setup for the zoom-in -simulations | 5 |
| 4.3.1 Cosmological setup | 5 |
| 4.3.2 Low-resolution run | 5 |
| 4.3.3 Choosing the zoom-in regions | 5 |
| 4.3.4 Initial conditions | 5 |
| 5 Cosmological GADGET-3 simulations | 6 |
| 5.1 Computational load of the simulations | 6 |
| 5.2 Locating galaxy centers: the shrinking sphere -method | 6 |
| 5.3 Properties of the galaxies | 6 |
| 5.3.1 Rotation curves | 7 |
| 5.3.2 Star formation history | 9 |
| 5.3.3 Colors and magnitudes | 10 |
| 6 Simulations with KETJU | 14 |
| 7 Conclusions | 15 |

1. Introduction

1.1 Information about galaxies, shortly

1.2 Aim of the thesis

2. Background

2.1 Cosmology

2.1.1 Hubble parameter, Friedmann equations and so on

2.1.2 Cosmological perturbations

2.2 Early-type galaxies

2.2.1 Types of ellipticals

2.2.2 Photometric and kinematic profiles

2.3 Feedback processes

3. GADGET-3 and KETJU

- Haven't really thought about the contents of this chapter yet

3.1 Overview of GADGET-3

3.2 Smoothed Particle Hydrodynamics

3.3 Gas cooling?

3.4 Feedback?

3.5 KETJU

4. Creating initial conditions for the cosmological zoom-in simulations

4.1 Zoom-in technique

4.2 MUSIC

4.2.1 Basic properties of MUSIC

4.2.2 Algorithms of MUSIC

- Short section on how MUSIC generates initial particle positions, velocity fields and so on (basically summary of the first few sections of the MUSIC paper)

4.2.3 Creating an IC-file with a zoom-in box

- Step by step explanation of creating the IC file with a zoom-in region

4.3 GADGET-3 setup for the zoom-in -simulations

4.3.1 Cosmological setup

4.3.2 Low-resolution run

4.3.3 Choosing the zoom-in regions

- FoF -algorithm
- Conditions of the chosen halos
- Figure showing the zoom-in regions from the low res run

4.3.4 Initial conditions

| h_0 | Ω_m | Ω_b | Ω_Λ | σ_8 | ρ_{crit} |
|-------|------------|------------|------------------|------------|---------------------------------------|
| 70.3 | 0.276 | 0.045 | 0.724 | 0.811 | $9.28 \times 10^{-27} \text{ kg/m}^3$ |

Table 4.1: Cosmological parameters used for the simulations. If a simulation doesn't include baryons, the dark matter density parameter Ω_{DM} is equal to the matter density parameter Ω_m . If baryons are included, $\Omega_{\text{DM}} = \Omega_m - \Omega_b$.

- Information from GADGET3 config files

5. Cosmological GADGET-3 simulations

5.1 Computational load of the simulations

- Quick overview: CPUs used, time elapsed, where simulations were run

5.2 Locating galaxy centers: the shrinking sphere -method

5.3 Properties of the galaxies

| Simulation | r_{vir} (kpc) | M_* (M_{\odot}) | M_V (mag) | $M_{*,\text{gal}}/M_{\text{vir}}$ | M_{bh} (M_{\odot}) |
|-------------|------------------------|-----------------------|-------------|-----------------------------------|---------------------------------|
| Med res, A | 517 | 3.98×10^{11} | -22.4 | 0.025 | 5.82×10^9 |
| Med res, B | 574 | 5.16×10^{11} | -22.7 | 0.024 | 6.23×10^9 |
| Med res, C | 400 | 2.56×10^{11} | -22.0 | 0.035 | 3.96×10^9 |
| High res, A | 526 | 5.31×10^{11} | -22.6 | 0.032 | 3.64×10^9 |
| High res, B | 578 | 6.38×10^{11} | -22.8 | 0.029 | 4.54×10^9 |
| High res, C | 400 | 3.46×10^{11} | -22.2 | 0.047 | 2.80×10^9 |

Table 5.1: Properties of the zoomed-in galaxies at redshift $z = 0$.

5.3.1 Rotation curves




Figure 5.1: Rotation curves for each galaxy including baryons, at redshift 0. The continuous lines represent the high resolution simulations and the dashed lines represent the medium resolution simulations.

- Med res galaxy A simulation is performed with single precision, I'm currently doing the simulation in double precision
- Rotation curves do not have the same limits on the y-axis




Figure 5.2: Rotation curves for each galaxy including only dark matter, at redshift 0. The continuous lines represent the high resolution simulations and the dashed lines represent the medium resolution simulations.




Figure 5.3: Rotation curves for each galaxy including baryons, at redshift 1. The continuous lines represent the high resolution simulations and the dashed lines represent the medium resolution simulations.




Figure 5.4: Rotation curves for each galaxy including baryons, at redshift 3. The continuous lines represent the high resolution simulations and the dashed lines represent the medium resolution simulations.

5.3.2 Star formation history

- The stellar mass evolution plot is not yet done for high res simulations.
- Redshifts missing from the stellar mass evolution plot
- SFRs, also histograms?
- Again, med res A results will probably change a bit when double precision run is finished.
- Formation efficiencies, comparing to the cosmological parameter




Figure 5.5: Stellar mass evolution for the medium resolution galaxies. The calculated stellar mass is the stellar mass within $r_{\text{gal}} = r_{\text{vir}}/10$.

5.3.3 Colors and magnitudes




Figure 5.6: Stellar formation rates for each zoomed-in galaxy, plotted as a function of lookback time. The lines are created from histograms having a length of 5 Myr, which are then smoothed. The continuous lines represent the high resolution simulations and the dashed lines represent the medium resolution simulations.




Figure 5.7: Galaxy formation efficiencies for each galaxy, plotted with their stellar masses. The cosmological baryon density Ω_b is set to 0.045 in the simulations. The diamond and circular markers show the results of the high resolution and the medium resolution zoom-in simulations, respectively.




Figure 5.8: B-R colors for each simulated galaxy, plotted with each galaxy's absolute magnitude in the V-band. The diamond and circular markers show the results of the high resolution and the medium resolution zoom-in simulations, respectively.




Figure 5.9: R-K colors for each simulated galaxy, plotted with each galaxy's absolute magnitude in the V-band. The diamond and circular markers show the results of the high resolution and the medium resolution zoom-in simulations, respectively.

6. Simulations with KETJU

7. Conclusions

- recap on what was written/studied
- more own thoughts on results
- future missions
- how could the simulations be more realistic (higher resolution, more feedback stuff?)