

November 15, 2022
Today's Paper

The 2022 Chronicle

20 °C 26 °C 13 °C
Dow +0.1% ↑

World U.S. Politics N.Y. Business Opinion Tech Science Health Sports Arts Books Style Food Travel Magazine T Magazine Real Estate Video

Fotios Ptochos Promoted to Professor!

Congratulations to Fotios Ptochos for his promotion to the rank of Full Professor, effective November 2022. This promotion recognizes Prof. Ptochos' achievements in scholarship, teaching in physics and research in high-energy physics (HEP), and his overall service to the CDF and CMS Collaborations. He is a Harvard University PhD in physics graduate (1998) and has been active in HEP-research, both in detector development and physics analyses since 1987. In particular, from 1987 to 1988 he worked in the development of a technique to monitor the purity of Liquid Argon (LAr) for the first ever prototype of the ICARUS detector, a technique that was subsequently used in the experiment. From 1989 to 1994 he worked in the characterization of various Tetramethyl liquids as part of a research project to find appropriate warm liquids media for the envisioned calorimeter detectors at SSC. He also worked in the construction, installation and calibration of the Central Muon Extension (CMX) system for the CDF detector. From 1994 to 1996 he developed an algorithm to improve electron identification for the CDF end-plug ECAL based on the information from the calorimeter and hits on the silicon tracker detector. The algorithm led to the development and implementation of the PHOENIX tracker system in the CDF-II detector. In the period of 2000–2003, he was the coordinator of the group responsible for the development, installation, maintenance and performance monitoring of the CDF-II Hadronic Calorimeter (HCAL) timing system. For the entire period of the Tevatron Run-II (2001–2011) he served as the coordinator of the CDF central HCAL calibration (CHA and WHA), maintenance and performance group. Since 2004, when he joined the faculty of the UCY Physics Department, he has been involved in the UCY HEP group activities related to the construction and running of the CMS ECAL at CERN. In 2009, he initiated the involvement of the group in the activities related

to the CMS tracking detector. He was also involved in the development of the dual-readout calorimetry concept in a total absorption HCAL for future linear-collider experiments. Professor Ptochos has led numerous physics analyses, spanning from precision measurements on properties of heavy flavour quark production and their use as probes for searching for the SM and SUSY Higgs bosons, to searches for BSM physics including SUSY, extra dimensions and other exotic processes. He has tremendous experience in heavy flavour tagging techniques and algorithms, tau-lepton identification techniques and new physics model building. He was the first ever recipient of the "Fermi National Accelerator Laboratory Fellowship" and has co-coordinated multiple research program funded primarily by the European Commission (EC) via Marie Skłodowska-Curie Actions, the Cyprus Research Promotion Foundation (RPF) through Didaktor or Excellence Hubs programs, the European Regional Development Fund, and UCY. Professor Ptochos is the author and co-author of more than 1700 publications in refereed scientific journals and a member of the editorial group in charge for producing the education material for the entire Cyprus Secondary Education. In addition, he has been the supervisor of the research activities of six postdoctoral fellows, five PhD and eleven MSc students, as well as the theses projects of more than 20 undergraduate students.



Multi-Muons In CDF: The Mystery Continues

We present a phenomenological conjecture of new physics that is suggested by the topology and kinematic properties of the multi-muon events recently reported by the CDF collaboration. We show that the salient features of the data can be accounted for by postulating the pair production of three new states h_1 , h_2 , and h_3 with masses in the range of 15, 7.3, and $3.6 \text{ GeV}/c^2$, respectively. The heavier states cascade-decay into the lighter ones, whereas the lightest state decays into a τ pair with a lifetime of the order of 20 ps.

Physicists Find Elusive Particle Seen as Key to Universe

Results are presented from searches for the standard model Higgs boson in proton-proton collisions at 7 and 8 TeV in the Compact Muon Solenoid experiment at the LHC, using data samples corresponding to integrated luminosities of up to 5.1 fb^{-1} at 7 TeV and 5.3 fb^{-1} at 8 TeV. The search is performed in five decay modes: $\gamma\gamma$, ZZ, $\tau^+\tau^-$, and $b\bar{b}$. An excess of events is observed above the expected background, with a local significance of 5.0 standard deviations, at a mass near 125 GeV, signalling the production of a new



Top Quark, Last Piece of Matter, Appears to Be in Place

We establish the existence of the top quark using a 67 pb^{-1} data sample of pp collisions at $\sqrt{s} = 1.8 \text{ TeV}$ collected with the Collider Detector at Fermilab (CDF). Employing techniques similar to those we previously published, we observe a signal consistent with $t\bar{t}$ decay to WWbb, but inconsistent with the background prediction by 4.8σ . Additional evidence for the top



CDF publishes multi-muons!

We report a study of multi-muon events produced at the Fermilab Tevatron collider and recorded by the CDF II detector. In a data set acquired with a dedicated dimuon trigger and corresponding to an integrated luminosity of 2100 pb^{-1} , we isolate a significant sample of events in which at least one of the muon candidates is produced outside of the beam pipe, in terms of the same understanding of the CDF II detector, trigger, and event reconstruction. Several topological and kinematic properties of these events are

which include heavy flavor production. In contrast, we are presently unable to fully account for the number and properties of the remaining events, in which at least one muon candidate is produced outside of the beam pipe, in terms of the same understanding of the CDF II detector, trigger, and event reconstruction. Several topological and kinematic properties of these events are