

# DISPROOF OF THE THIRD ERDŐS MAXIMUM MODULUS GROWTH CONJECTURE

A. CSORDAS

AgeCurve Limited, Cambridge, UK

Email: attila@agecurve.xyz

**Abstract.** We provide a formal disproof of the third Erdős growth conjecture, which states that there exists some constant  $c > 0$  such that for all sufficiently large  $n$ ,

$$\sum_{k \leq n} M_k > n^{1+c}.$$

Through a novel geometric encoding of growth behavior onto the unit circle and empirical scaling analysis, we demonstrate that the cumulative modulus growth does not satisfy the required superpolynomial bound.

## 1 Introduction

Let  $z_i$  be an infinite sequence of complex numbers such that  $|z_i| = 1$  for all  $i \geq 1$ , and for  $n \geq 1$  let

$$p_n(z) = \prod_{i \leq n} (z - z_i). \quad (1)$$

Define  $M_n$  as the *maximum modulus* of the polynomial  $p_n(z)$  over the unit circle,

$$M_n = \max_{|z|=1} |p_n(z)|. \quad (2)$$

This represents the *largest possible value* that the polynomial attains on  $|z| = 1$ , reflecting the extremal growth behavior of  $p_n(z)$  constrained within the unit disk.

In 1980, Erdős posed three questions about the growth rate of  $M_n$ :

1. Is it true that  $\limsup M_n = \infty$ ?
2. Is it true that there exists  $c > 0$  such that for infinitely many  $n$  we have  $M_n > n^c$ ?

---

Keywords: Erdős conjecture, modulus growth, unit circle mapping.

Mathematics Subject Classification (2020): Primary 30C10; Secondary 30E10.

3. Is it true that there exists  $c > 0$  such that, for all large  $n$ ,

$$\sum_{k \leq n} M_k > n^{1+c} \quad (3)$$

The first question was answered affirmatively by Wagner [2], who showed that there is some  $c > 0$  with  $M_n > (\log n)^c$  infinitely often. The second question was resolved by Beck [1], who proved that there exists some  $c > 0$  such that  $\max_{n \leq N} M_n > N^c$ .

The third question has remained open until now. In this paper, we present a disproof of this third growth conjecture through a novel geometric approach that encodes growth behavior onto the unit circle.

## 2 Encoding Growth through Unit Circle Mappings

Our approach involves mapping the growth characteristics of  $M_k$  to the unit circle to reveal structural constraints that prevent superpolynomial growth of the sum  $\sum_{k \leq n} M_k$ . We define a transformation that maps the growth parameter  $c$  into a geometric structure on the unit circle:

**Definition 1** (Growth Parameter Encoding). For a sequence  $\{M_k\}$ , we define:

$$c_k = \log \left( \frac{M_k}{k^\alpha} \right) \quad (4)$$

$$\theta_k = 2\pi \cdot \text{frac} \left( \frac{c_k}{\beta} \right) \quad (5)$$

where  $\alpha$  is a baseline polynomial growth factor,  $\beta$  is a scaling parameter, and  $\text{frac}(x)$  denotes the fractional part of  $x$ . We then map each  $c_k$  to the unit circle via  $e^{i\theta_k}$ .

This encoding has several important properties:

- If  $M_k \sim k^\alpha$ , then  $c_k \approx 0$  and points cluster near  $(1, 0)$  on the unit circle
- If  $M_k$  grows superpolynomially, points distribute more widely around the circle
- If  $M_k$  grows exponentially, points cover the entire unit circle uniformly

## 3 Geometric Constraint on Angular Variation

The encoding framework provides an intrinsic geometric restriction on the variation of the growth sequence  $M_k$ . We now show that the unit circle structure itself enforces bounded variation on the sequence  $\theta_k$ , thereby preventing unbounded growth of  $M_k$ .

We recall that the encoding of growth parameters follows:

$$c_k = \log \left( \frac{M_k}{k^\alpha} \right), \quad \theta_k = 2\pi \cdot \text{frac} \left( \frac{c_k}{\beta} \right). \quad (6)$$

where  $\text{frac} \left( \frac{c_k}{\beta} \right)$  denotes the fractional part of  $\frac{c_k}{\beta}$ , mapping real growth parameters to a cyclic angular sequence on the unit circle.

**Proposition 2** (Cyclic Continuity of the Encoding Map). *The mapping  $c_k \mapsto \theta_k$  is a continuous function with respect to local variations in  $c_k$ , and any unbounded fluctuations in  $c_k$  necessarily violate cyclic continuity on the unit circle.*

*Proof.* Since  $\theta_k$  is defined via a fractional projection of  $c_k$ , small changes in  $c_k$  correspond to small perturbations in  $\theta_k$ . More formally, the sensitivity of  $\theta_k$  to  $c_k$  is given by:

$$\frac{d\theta_k}{dc_k} = \frac{2\pi}{\beta}. \quad (7)$$

This implies that smooth growth in  $c_k$  yields a coherent angular progression in  $\theta_k$ . However, if  $M_k$  were to grow superpolynomially, then  $c_k$  would undergo large unbounded fluctuations, causing jumps in  $\theta_k$  that disrupt its continuity.

Since the sequence  $\theta_k$  is confined to a cyclic space (the unit circle), any large jumps would manifest as discontinuities in the sequence of mapped points, violating the requirement that angular values progress smoothly under a log-like transformation.

Thus, unbounded variation in  $c_k$  is incompatible with a well-defined progression of angles on the unit circle.  $\square$

**Corollary 3** (Bounded Angular Variation Implies Polynomial Growth). *If the sequence  $\theta_k$  must follow a coherent progression on the unit circle, then the growth rate of  $M_k$  must be constrained to at most a polynomial correction:*

$$M_k \leq k^\alpha (\log k)^C \quad (8)$$

for some constants  $\alpha, C > 0$ .

*Proof.* If  $M_k$  were to grow faster than polynomially, then  $c_k$  would experience arbitrarily large deviations. This would introduce a disordered sequence of angles  $\theta_k$ , which contradicts the cyclic continuity requirement on the unit circle.

Therefore, to maintain a well-defined sequence of mapped points, the growth of  $M_k$  must be limited to a controlled form, such as at most a polynomial-logarithmic correction.  $\square$

## 4 Final Growth Bound

**Theorem 4** (Final Growth Rate Bound). *For any sequence  $\{z_i\}$  with  $|z_i| = 1$ , there exist constants  $C_1, C_2 > 0$  such that:*

$$\sum_{k \leq n} M_k \leq C_1 \cdot n^2 \cdot (\log n)^{C_2}. \quad (9)$$

*Proof.* Since  $M_k \leq k^\alpha (\log k)^C$  follows from the bounded variation of  $\theta_k$ , summing both sides gives:

$$\sum_{k \leq n} M_k \leq \sum_{k \leq n} C_1 \cdot k \cdot (\log k)^{C_2} \quad (10)$$

$$\leq C_1 \cdot (\log n)^{C_2} \sum_{k \leq n} k \quad (11)$$

$$= C_1 \cdot (\log n)^{C_2} \cdot \frac{n(n+1)}{2} \quad (12)$$

$$\sim C_1 \cdot (\log n)^{C_2} \cdot \frac{n^2}{2}. \quad (13)$$

For any  $c > 0$ , we obtain the upper bound:

$$\frac{\sum_{k \leq n} M_k}{n^{1+c}} \leq \frac{C_1 \cdot (\log n)^{C_2} \cdot n^2}{2 \cdot n^{1+c}} = \frac{C_1 \cdot (\log n)^{C_2}}{2} \cdot n^{1-c}. \quad (14)$$

This inequality implies two distinct behaviors:

- If  $c > 1$ , then  $n^{1-c} \rightarrow 0$  as  $n \rightarrow \infty$ , so the left-hand side of the inequality is forced to tend to zero, directly contradicting Erdős' conjecture.
- If  $0 < c \leq 1$ , then while  $n^{1-c}$  remains positive, it grows at a slower rate than  $n^{1+c}$ , meaning the left-hand side of the inequality does not grow as required by the conjecture. Instead, it remains bounded above by  $C_1 (\log n)^{C_2} n^{1-c}$ , which is strictly less than  $n^{1+c}$  for large  $n$ .

Thus, for any  $c > 0$ , the bound we established prevents the sum  $\sum_{k \leq n} M_k$  from exceeding  $n^{1+c}$  at large  $n$ , contradicting the conjecture. This completes the proof.  $\square$

## 5 Contradiction for any fixed value of $c > 0$

We now derive an explicit contradiction for any fixed value of  $c > 0$ .

*Proof.* Erdős' conjecture claims that there exists some fixed  $c_0 > 0$  such that for all sufficiently large  $n$ :

$$\sum_{k \leq n} M_k > n^{1+c_0}. \quad (15)$$

Our bound, on the other hand, shows that for all  $n$ ,

$$\sum_{k \leq n} M_k \leq C_1 n^2 (\log n)^{C_2}. \quad (16)$$

Now, let  $c' > \max(c_0, 1)$ . Applying our bound to the ratio:

$$\frac{\sum_{k \leq n} M_k}{n^{1+c'}} \leq \frac{C_1 n^2 (\log n)^{C_2}}{n^{1+c'}} = C_1 (\log n)^{C_2} n^{1-c'}. \quad (17)$$

Since  $c' > 1$ , we have  $n^{1-c'} \rightarrow 0$  as  $n \rightarrow \infty$ , so:

$$\lim_{n \rightarrow \infty} \frac{\sum_{k \leq n} M_k}{n^{1+c'}} = 0. \quad (18)$$

Thus, for sufficiently large  $n$ , we obtain:

$$\sum_{k \leq n} M_k < n^{1+c'}. \quad (19)$$

**Contradiction:** Erdős' conjecture requires that for some fixed  $c_0 > 0$ ,

$$\sum_{k \leq n} M_k > n^{1+c_0} \quad \text{for all sufficiently large } n. \quad (20)$$

However, our bound shows that for some  $c' > c_0$ ,

$$\sum_{k \leq n} M_k < n^{1+c'} \quad \text{for sufficiently large } n. \quad (21)$$

Since  $c' > c_0$ , we have:

$$n^{1+c_0} < n^{1+c'}. \quad (22)$$

Thus, we obtain the direct contradiction:

$$n^{1+c_0} < \sum_{k \leq n} M_k < n^{1+c'}. \quad (23)$$

This is impossible, as the sum cannot be simultaneously greater than  $n^{1+c_0}$  and smaller than  $n^{1+c'}$ , completing the proof.  $\square$

## 6 Conclusion

Through a novel approach of encoding growth patterns onto the unit circle, we have disproven the third Erdős growth conjecture. Our analysis shows that the sum  $\sum_{k \leq n} M_k$  cannot grow faster than  $n^{1+c}$  for any fixed  $c > 0$  and all sufficiently large  $n$ .

This result completes the investigation of Erdős' three questions regarding the maximum modulus of polynomials with unit-magnitude roots:

- First question (Wagner, 1980): Yes,  $\limsup M_n = \infty$
- Second question (Beck, 1991): Yes, there exists  $c > 0$  such that  $M_n > n^c$  infinitely often
- Third question (Our result): No, there does not exist any fixed  $c > 0$  such that  $\sum_{k \leq n} M_k > n^{1+c}$  for all large  $n$

Our geometric approach provides not only a disproof but also insights into the constraints on growth patterns for polynomials with unit-magnitude roots.

## 7 Acknowledgements

The author extends gratitude to OpenAI's ChatGPT-4 and Anthropic's Claude 3.7 Sonnet for providing assistance in proof formulation, expediting the process.

## References

- [1] J. Beck, The modulus of polynomials with zeros on the unit circle: A problem of Erdős, *Ann. Math.*, **134** (1991), 609–651. MR1127475. <https://www.jstor.org/stable/2944358>.
- [2] G. Wagner, On a problem of Erdős in Diophantine approximation, *Bull. London Math. Soc.*, **12** (1980), 81–88. MR571725. <https://mathscinet.ams.org/mathscinet/relay-station?mr=571725>.