

PHY115: Introduction

Anabela R. Turlione

Digipen

Spring 2022

Introduction

General information of the course

Kinematics

Doing vector calculation using components

Unit Vectors

Review: Functions

General information of the course

Grading Policy

1. HOMEWORKS 40 % (Quiz in Moodle, every 2/3 weeks)
2. MIDTERM 30 % (test week 7)
3. FINAL EXAM 30 % (test week 14)

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)
2. Math basics

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)
2. Math basics
 - ▶ vectors
 - ▶ trigonometry
 - ▶ algebra
 - ▶ functions

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)
2. Math basics
 - ▶ vectors
 - ▶ trigonometry
 - ▶ algebra
 - ▶ functions
3. Mechanics (understand what causes the motion)

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)
2. Math basics
 - ▶ vectors
 - ▶ trigonometry
 - ▶ algebra
 - ▶ functions
3. Mechanics (understand what causes the motion)
4. Rotation of rigid bodies

Upon a successfully completion of this course the students will gain a fundamental understanding of basic physical principles including:

1. Kinematics (describe motion)
2. Math basics
 - ▶ vectors
 - ▶ trigonometry
 - ▶ algebra
 - ▶ functions
3. Mechanics (understand what causes the motion)
4. Rotation of rigid bodies
5. Light

Why to study Physics?

1. you'll need to understand what goes on in real life so you can make decisions about how to convey motion and effects in your work (plausible motion).

Why to study Physics?

1. you'll need to understand what goes on in real life so you can make decisions about how to convey motion and effects in your work (plausible motion).
2. Keep the suspension of Disbelief: The representation of any story succeeds only because of the audience's willingness to ignore the fact that it's just a story.

Why to study Physics?

1. you'll need to understand what goes on in real life so you can make decisions about how to convey motion and effects in your work (plausible motion).
2. Keep the suspension of Disbelief: The representation of any story succeeds only because of the audience's willingness to ignore the fact that it's just a story.
3. Too many errors in the motion will distract viewers and remind them that it's just a story.

Why to study Physics?

1. you'll need to understand what goes on in real life so you can make decisions about how to convey motion and effects in your work (plausible motion).
2. Keep the suspension of Disbelief: The representation of any story succeeds only because of the audience's willingness to ignore the fact that it's just a story.
3. Too many errors in the motion will distract viewers and remind them that it's just a story.
 - ▶ Wrong response of material to light
 - ▶ Objects that move as if they were heavier than they look
 - ▶ Different falling acceleration for different bodies
 - ▶ ...

What is Physics?

What is Physics?

- ▶ Physics describes the nature. It's based in try-error Laws.

What is Physics?

- ▶ Physics describes the nature. It's based in try-error Laws.
- ▶ The language is math.

What is Physics?

- ▶ Physics describes the nature. It's based in try-error Laws.
- ▶ The language is math.
- ▶ kinematics describes the motion path of objects (the timing and spacing)

What is Physics?

- ▶ Physics describes the nature. It's based in try-error Laws.
- ▶ The language is math.
- ▶ kinematics describes the motion path of objects (the timing and spacing)
- ▶ Dynamics describes what causes the motion


Why do we want to do this?

Why do we want to do this?

If we are able to understand the motion of bodies, we can predict their motion and do incredible things. . .

Why do we want to do this?

If we are able to understand the motion of bodies, we can predict their motion and do incredible things. . .




On the other hand...

On the other hand. . .


Knowing Dynamics can help us figure out accurate timing. Even if you plan to exaggerate your motion for effect, it helps to start with reality and work from there.

Kinematics




- Describes the motion path: Position vs. time $x(t)$, $y(t)$ and $z(t) \rightarrow$ timing and spacing.

Kinematics



- ▶ Describes the motion path: Position vs. time $x(t)$, $y(t)$ and $z(t) \rightarrow$ timing and spacing.
- ▶ There are three main quantities that describe the motion: **acceleration**, **velocity** and **position**


Kinematics




- ▶ Describes the motion path: Position vs. time $x(t)$, $y(t)$ and $z(t) \rightarrow$ timing and spacing.
- ▶ There are three main quantities that describe the motion: **acceleration**, **velocity** and **position**
- ▶ The position is related with velocity and the velocity is related with acceleration.

You do this intuitively . . .


You do this intuitively . . .



So our chart will look like this.



You do this intuitively ...




So our chart will look like this.




What is the right separation?

You do this intuitively ...


You do this intuitively ...




So our chart will look like this.



You do this intuitively ...




So our chart will look like this.




What generate that exact separation?

You do this intuitively ...




So our chart will look like this.




What generate that exact separation? **FORCES**

You do this intuitively ...



So our chart will look like this.



What generate that exact separation? **FORCES Dynamics**

We are going to find those laws, but first, we need to review some Math basics:

We are going to find those laws, but first, we need to review some Math basics:

- ▶ Frames of Reference
- ▶ escalar and vectors
- ▶ funciones
- ▶ trigonometry

Frame of reference

When we want to describe the motion of a body, the first thing we need to define is a coordinate system.

Frame of reference


When we want to describe the motion of a body, the first thing we need to define is a coordinate system.

- ▶ The position is determined by the numbers (x, y, z)

Frame of reference

When we want to describe the motion of a body, the first thing we need to define is a coordinate system.



- ▶ The position is determined by the numbers (x, y, z)
- ▶ Defining these 3 numbers plus a time scale we can represent the position vs time.




Frame of reference

When we want to describe the motion of a body, the first thing we need to define is a coordinate system.

- ▶ The position is determined by the numbers (x, y, z)
- ▶ Defining these 3 numbers plus a time scale we can represent the position vs time.



Another example of a reference system . . .



Vectors


- Some physical quantities, such as time, temperature, mass, and density, can be described completely by a single number with a unit.

Vectors

- ▶ Some physical quantities, such as time, temperature, mass, and density, can be described completely by a single number with a unit.
- ▶ But many other important quantities in physics have a direction associated with them and cannot be described by a single number

For example, you use vectors to determine the direction of a curve in vectorial illustration. . .

For example, you use vectors to determine the direction of a curve in vectorial illustration. . .



Another examples of vectors?

Another examples of vectors?

- ▶ velocity
- ▶ acceleration
- ▶ force
- ▶ ...

Another examples of vectors?

- ▶ velocity
- ▶ acceleration
- ▶ force
- ▶ ...

How do we use them?

Vectors

- ▶ When a physical quantity is described by a single number, we call it a **scalar quantity**.


Vectors

- ▶ When a physical quantity is described by a single number, we call it a **scalar quantity**.
- ▶ A **vector quantity** has both a magnitude and a direction in space.


Vectors

- ▶ When a physical quantity is described by a single number, we call it a **scalar quantity**.
- ▶ A **vector quantity** has both a magnitude and a direction in space.
- ▶ We represent a vector quantity by a single letter, italic type with an arrow above it: \vec{v}


- ▶ Vectors are represented by arrows in diagrams.
- ▶ the magnitude is just the length of the arrow.
- ▶ $\vec{A} = \vec{A}'$



- ▶ If two vectors have the same magnitude and the same direction, they are equal, no matter where they are located in space.
- ▶ $\vec{A} = \vec{A}'$



- ▶ We define the negative of a vector as a vector having the same magnitude as the original vector but the opposite direction.
- ▶ $\vec{A} = -\vec{B}$
- ▶ When two vectors are parallel and have opposite directions, we say that they are antiparallel.



We usually represent the magnitude of a vector by the same letter used for the vector and vertical bars on both sides:

We usually represent the magnitude of a vector by the same letter used for the vector and vertical bars on both sides:

$$(\text{magnitude of } \vec{A} = |\vec{A}|) \quad (1)$$

We usually represent the magnitude of a vector by the same letter used for the vector and vertical bars on both sides:

$$(\text{magnitude of } \vec{A} = |\vec{A}|) \quad (1)$$

- The magnitude of a vector is a positive scalar (a positive number).


We usually represent the magnitude of a vector by the same letter used for the vector and vertical bars on both sides:

$$(\text{magnitude of } \vec{A} = |\vec{A}|) \quad (1)$$

- ▶ The magnitude of a vector is a positive scalar (a positive number).
- ▶ A vector can never be equal to a scalar because they are different kind of quantities.


Displacement

- Displacement is simply a change in the position of an object.




Displacement

- Displacement is simply a change in the position of an object.





Displacement

- ▶ Displacement is simply a change in the position of an object.
- ▶ It is a vector because we must state how far the object moves and the direction.



Displacement


- ▶ Displacement is simply a change in the position of an object.
- ▶ It is a vector because we must state how far the object moves and the direction.
- ▶ The displacement only depend on the initial and final positions.



Operations with vector: Addition and subtraction

Operations with vector: Addition and subtraction


Suppose a particle undergoes a displacement \vec{A} followed by a second displacement \vec{B} . The final result is the same as if the particle had started at the same initial point and undergone a single displacement \vec{C} .



Operations with vector: Addition and subtraction

$$\vec{C} = \vec{A} + \vec{B}$$

- ▶ vector addition obeys the commutative law
- ▶ $\vec{A} + \vec{B} = \vec{B} + \vec{A}$
- ▶ note that the magnitude of $|\vec{C}| \leq |\vec{A}| + |\vec{B}|$




Sum of more than two vector:

$$\vec{R} = (\vec{A} + \vec{B}) + \vec{C} = \vec{D} + \vec{C}$$


or

$$\vec{R} = \vec{A} + (\vec{B} + \vec{C}) = \vec{A} + \vec{E}$$



We define the difference of two vectors \vec{A} and \vec{B} to be the vector sum of \vec{A} and $-\vec{B}$:

$$\vec{A} - \vec{B} = \vec{A} + (-\vec{B})$$




A vector can be multiplied by a positive (negative) scalar quantity. The result is a vector in the same direction (opposite) as the vector but with a different magnitude.

The magnitude is $|c||\vec{A}|$


EXAMPLE 1: Adding two vectors that are perpendicular

A cross-country skier skis 1.00 km north and then 2.00 km east on a horizontal snowfield. How far and in what direction is she from the starting point?

Distance from the starting point:




Distance from the starting point:



$$\sqrt{(1.00 \text{ km})^2 + (2.00 \text{ km})^2} = 2.24 \text{ km}$$


Distance from the starting point:



$$\sqrt{(1.00 \text{ km})^2 + (2.00 \text{ km})^2} = 2.24 \text{ km}$$

direction:

Distance from the starting point:




$$\sqrt{(1.00 \text{ km})^2 + (2.00 \text{ km})^2} = 2.24 \text{ km}$$

direction:

$$\tan \phi = \frac{\text{opposite side}}{\text{Adjacent side}} = \frac{2.00 \text{ km}}{1.00 \text{ km}}$$

Distance from the starting point:



$$\sqrt{(1.00 \text{ km})^2 + (2.00 \text{ km})^2} = 2.24 \text{ km}$$

direction:

$$\tan \phi = \frac{\text{opposite side}}{\text{Adjacent side}} = \frac{2.00 \text{ km}}{1.00 \text{ km}}$$

$$\phi = 63.4^\circ$$

EXAMPLE 2: Test Your Understanding

Two displacement vectors, \vec{u} and \vec{v} have magnitudes 3 and 4. Which of the following could be the magnitude of the difference vector (There may be more than one correct answer.)

- ▶ 9 m
- ▶ 7 m
- ▶ 5 m
- ▶ 1 m
- ▶ 0 m
- ▶ -1 m

- Calculations with right triangles work only when the two vectors are perpendicular.

- ▶ Calculations with right triangles work only when the two vectors are perpendicular.
- ▶ Measuring a diagram offers only very limited accuracy.

- ▶ Calculations with right triangles work only when the two vectors are perpendicular.
- ▶ Measuring a diagram offers only very limited accuracy.
- ▶ So we need a simple but general method for adding vectors.

- ▶ Calculations with right triangles work only when the two vectors are perpendicular.
- ▶ Measuring a diagram offers only very limited accuracy.
- ▶ So we need a simple but general method for adding vectors.
- ▶ This is called the method of components.

COMPONENTS OF VECTORS

- ▶ Consider a rectangular (Cartesian) coordinate system of axes

COMPONENTS OF VECTORS

- ▶ Consider a rectangular (Cartesian) coordinate system of axes
- ▶ We then draw the vector with its tail at O

COMPONENTS OF VECTORS

- ▶ Consider a rectangular (Cartesian) coordinate system of axes
- ▶ We then draw the vector with its tail at O
- ▶ We can represent any vector in the xy -plane as the sum of a vector in the x -axis and a vector in the y -axis

COMPONENTS OF VECTORS


- ▶ Consider a rectangular (Cartesian) coordinate system of axes
- ▶ We then draw the vector with its tail at O
- ▶ We can represent any vector in the xy-plane as the sum of a vector in the x-axis and a vector in the y-axis

$$\vec{A} = \vec{A}_x + \vec{A}_y \quad (2)$$

COMPONENTS OF VECTORS

- ▶ Consider a rectangular (Cartesian) coordinate system of axes
- ▶ We then draw the vector with its tail at O
- ▶ We can represent any vector in the xy-plane as the sum of a vector in the x-axis and a vector in the y-axis

$$\vec{A} = \vec{A}_x + \vec{A}_y \quad (2)$$



- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.

- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.
- ▶ When \vec{A}_x points in the positive x-direction, we define the number A_x , $A_x = |\vec{A}_x|$

- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.
- ▶ When \vec{A}_x points in the positive x-direction, we define the number A_x , $A_x = |\vec{A}_x|$
- ▶ When \vec{A}_x points in the negative x-direction, we define the number A_x , $A_x = -|\vec{A}_x|$

- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.
- ▶ When \vec{A}_x points in the positive x-direction, we define the number A_x , $A_x = |\vec{A}_x|$
- ▶ When \vec{A}_x points in the negative x-direction, we define the number A_x , $A_x = -|\vec{A}_x|$
- ▶ We define the number \vec{A}_y in the same way

- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.
- ▶ When \vec{A}_x points in the positive x-direction, we define the number A_x , $A_x = |\vec{A}_x|$
- ▶ When \vec{A}_x points in the negative x-direction, we define the number A_x , $A_x = -|\vec{A}_x|$
- ▶ We define the number \vec{A}_y in the same way

The two numbers A_x and A_y are called the components of \vec{A} .

- ▶ Since each component vector lies along a coordinate-axis direction, we need only a single number to describe each one.
- ▶ When \vec{A}_x points in the positive x-direction, we define the number A_x , $A_x = |\vec{A}_x|$
- ▶ When \vec{A}_x points in the negative x-direction, we define the number A_x , $A_x = -|\vec{A}_x|$
- ▶ We define the number \vec{A}_y in the same way

The two numbers A_x and A_y are called the components of \vec{A} .

We represent \vec{A} using its components as:


$$\vec{A} = (A_x, A_y) \quad (3)$$

- If we know the direction and the magnitude of a vector, we can know the components.

$$\frac{A_x}{A} = \cos\theta, \text{ and } \frac{A_y}{A} = \sin\theta$$

$$A_x = A\cos\theta, \text{ and } A_y = A\sin\theta$$

$$\rightarrow \vec{A} = \vec{A}_x + \vec{A}_y$$



Finding a vector's magnitude and direction from its components.

$$A = \sqrt{A_x^2 + A_y^2} \quad (4)$$

$$\tan\theta = \frac{A_y}{A_x} \quad \text{and} \quad \theta = \arctan\left(\frac{A_y}{A_x}\right) \quad (5)$$

Multiplying a vector by a scalar.

$$\vec{D} = c\vec{A} \quad (6)$$


$$D_x = cA_x \text{ and } D_y = cA_y \quad (7)$$

Using the components to calculate the vector sum

$$\vec{R} = \vec{A} + \vec{B}$$

$$R_x = A_x + B_x$$

$$R_y = A_y + B_y$$




\vec{R} is the vector sum
(resultant) of \vec{A} and \vec{B} .

The components of \vec{R} are the sums
of the components of \vec{A} and \vec{B} :

$$R_y = A_y + B_y \quad R_x = A_x + B_x$$


EXAMPLE:

You are using vector sums to move objects in 3ds Max all the time...




EXAMPLE:

Position vector




EXAMPLE:

Displacement Vector




EXAMPLE:

Displacement Vector



EXAMPLE:


Final Position = Position Vector +
Displacement Vector



EXAMPLE:

A more elegant notation...


$$\vec{r}_f = \vec{r}_i + \Delta\vec{r} \quad (8)$$



EXAMPLE:



so, the displacement vector is...

$$\Delta \vec{r} = \vec{r}_f - \vec{r}_i \quad (9)$$





In general, we use the symbol Δ to represent the change of a magnitude...

- In the previous definitions we measure the angles as in the figure.



EXAMPLE: (a) What are the x and y components of vector in the figure? The magnitude of the vector \vec{D} is $D = 3.00\text{ m}$, and the angle $\alpha = 45^\circ$. (b) What are the x- and y-components of vector \vec{E} in? The magnitude of the vector is $E = 4.50\text{ m}$, and the angle $\beta = 45^\circ$



Adding more than one vector:

$$\vec{R} = \vec{A} + \vec{B} + \vec{C}$$

$$\rightarrow R_x = A_x + B_x + C_x$$

$$\rightarrow R_y = A_y + B_y + C_y$$

$$\rightarrow R_z = A_z + B_z + C_z$$

Magnitude of a vector with 3 components:

$$|\vec{A}| = \sqrt{A_x^2 + A_y^2 + A_z^2}$$

Test your understanding:

Two vectors \vec{A} and \vec{B} both lie in the xy -plane.

- ▶ Is it possible for \vec{A} to have the same magnitude as \vec{B} but different components?
- ▶ Is it possible for \vec{A} to have the same components as \vec{B} but a different magnitude?

- ▶ A unit vector is a vector that has a magnitude of 1 with no units


- ▶ A unit vector is a vector that has a magnitude of 1 with no units
- ▶ Its only purpose is to describe a direction in space

- ▶ A unit vector is a vector that has a magnitude of 1 with no units
- ▶ Its only purpose is to describe a direction in space
- ▶ We represent them using a "hat"

- ▶ A unit vector is a vector that has a magnitude of 1 with no units
- ▶ Its only purpose is to describe a direction in space
- ▶ We represent them using a "hat"
- ▶ the unit vectors that point into the direction of the coordinate axes are:

- ▶ A unit vector is a vector that has a magnitude of 1 with no units
- ▶ Its only purpose is to describe a direction in space
- ▶ We represent them using a "hat"
- ▶ the unit vectors that point into the direction of the coordinate axes are:

\hat{i} , \hat{j} , and \hat{k}



We can represent component vectors like this:

$$\rightarrow \vec{A}_x = A_x \hat{i}$$


$$\rightarrow \vec{A}_y = A_y \hat{j}$$

$$\rightarrow \vec{A}_z = A_z \hat{k}$$

We can represent a vector using this notation:

$$\vec{A} = \vec{A}_x + \vec{A}_y$$

$$\rightarrow \vec{A} = A_x \hat{i} + A_y \hat{j}$$



How can we represent the sum of two vectors using this notation?

Guess!

Test your understanding

Arrange the following vectors in order of their magnitude, with the vector of largest magnitude first.

1. $\vec{A} = (3\hat{i} + 5\hat{j} - 2\hat{k})$
2. $\vec{A} = (-3\hat{i} + 5\hat{j} - 2\hat{k})$
3. $\vec{A} = (3\hat{i} - 5\hat{j} - 2\hat{k})$
4. $\vec{A} = (3\hat{i} + 5\hat{j} + 2\hat{k})$

Functions

Functions


- ▶ A function is a binary relation between two sets that associates every element of the first set to exactly one element of the second set

Functions

- ▶ A function is a binary relation between two sets that associates every element of the first set to exactly one element of the second set
- ▶ some useful functions:
 1. linear functions
 2. quadratic functions
 3. sin, cos
 4. exponential, logarithmic function

Examples ...

Examples ...





Relations



A relation shows a relationship between two values.

A function is a relation where each input has only one output.

Functions



Not Functions



Instead of using diagrams, we can use a law to draw the function in the x-y axes ...

Instead of using diagrams, we can use a law to draw the function in the x-y axes ...

$$y = f(x)$$

Linear Functions:


$$f(x) = Ax + C$$

Linear Functions:

$$f(x) = Ax + C$$

Representation in the xy axes
coordinates:

$$y = Ax + C$$



Cuadratic Functions:


$$f(x) = ax^2 + bx + c$$

Cuadratic Functions:

$$f(x) = ax^2 + bx + c$$


Representation in the xy axes
coordinates:

$$y = ax^2 + bx + c$$




Trygonometric Functions:

$$\sin(\alpha) = \frac{\text{opposite}}{\text{hypotenuse}}$$




Trygonometric Functions:

$$\cos(\alpha) = \frac{\text{adjacent}}{\text{hypotenuse}}$$




Trygonometric Functions:


$$\tan(\alpha) = \frac{\textit{opposite}}{\textit{adjacent}}$$



Trygonometric Functions:



Measuring angles: radians



Radians to degrees

How many rads are 360 deg?

Radians to degrees

How many rads are 360 deg?

$$\text{radians} = \frac{\text{arc length}}{\text{radius}}$$

Radians to degrees

How many rads are 360 deg?

$$\text{radians} = \frac{\text{arc length}}{\text{radius}} = \frac{2\pi}{1}$$

Radians to degrees

How many rads are 360 deg?

$$\text{radians} = \frac{\text{arc length}}{\text{radius}} = \frac{2\pi}{1}$$

$$2\pi \text{ rad} \rightarrow 360\text{deg}$$

$$x \text{ rad} \rightarrow \frac{360}{2\pi} x$$

Radians to degrees

How many rads are 360 deg?

$$\text{radians} = \frac{\text{arc length}}{\text{radius}} = \frac{2\pi}{1}$$

$$2\pi \text{ rad} \rightarrow 360\text{deg}$$

$$x \text{ rad} \rightarrow \frac{360}{2\pi} x = \frac{180}{\pi} x$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

$$\cos(30) = 0.86602540378$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

$$\cos(30) = 0.86602540378 \rightarrow \text{acos}(0.86602540378) = 30$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

$$\cos(30) = 0.86602540378 \rightarrow \text{acos}(0.86602540378) = 30$$

$$\tan(30) = 0.57735026919$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

$$\cos(30) = 0.86602540378 \rightarrow \text{acos}(0.86602540378) = 30$$

$$\tan(30) = 0.57735026919 \rightarrow \text{atan}(0.57735026919) = 30$$

How do I calculate the angle if I know the sin/cos/tan?

$$\sin(30) = 0.5 \rightarrow \text{asin}(0.5) = 30$$

$$\cos(30) = 0.86602540378 \rightarrow \text{acos}(0.86602540378) = 30$$

$$\tan(30) = 0.57735026919 \rightarrow \text{atan}(0.57735026919) = 30$$

You have to check what are the units of the angles. . .