PHY115

Anabela R. Turlione

Digipen

Spring 2022

Light

Introduction

Electric Field

Magnetic Field

Electromagnetic Radiation

Introduction Electric Field Magnetic Field Electromagnetic Radiation

Introduction

What is light?

What is light?

Light is Electromagnetic radiation

lacktriangle Changing Magnetic Field ightarrow Changing Electric Field

- ightharpoonup Changing Magnetic Field ightarrow Changing Electric Field
- lacktriangle Changing Electric Field ightarrow Changing Magnetic Field

Electromagnetic Wave: Wave of Electric and Magnetic Field

- ightharpoonup Changing Magnetic Field ightharpoonup Changing Electric Field
- lacktriangle Changing Electric Field ightarrow Changing Magnetic Field

Electromagnetic Wave: Wave of Electric and Magnetic Field

Changing Electric Field \rightarrow Changing Magnetic Field \rightarrow Changing Electric Field

- ► Changing Magnetic Field → Changing Electric Field
- lacktriangle Changing Electric Field ightarrow Changing Magnetic Field

Electromagnetic Wave: Wave of Electric and Magnetic Field

Changing Electric Field \rightarrow Changing Magnetic Field \rightarrow Changing Electric Field

The light is an electromagnetic wave that can propagate through space.

To describe an electromagnetic wave we have to know first:

To describe an electromagnetic wave we have to know first:

what is an Electric Field?

To describe an electromagnetic wave we have to know first:


- what is an Electric Field?
- what is a Magnetic Field?

Electric Field

Force between two point charged particles:


Electric Field

Force between two point charged particles:




Magnetic Field

Sources of Magnetic Fields: Ampere's Law



Magnetic Field

Sources of Magnetic Fields: Ampere's Law



The magnetic field generated by a straight wire is circular

Example: Magnet



Example: Magnet



In some materials called **Ferromagnetic**, the angular momentum of electrons are aligned so that they create a resultant macroscopic magnetic field.

Example: Magnet



In some materials called **Ferromagnetic**, the angular momentum of electrons are aligned so that they create a resultant macroscopic magnetic field.

Iron - Cobalt - Nickel

When the electric and magnetic fields are static, the are separate entities. But when they are changing, we can not treat them separately

When the electric and magnetic fields are static, the are separate entities. But when they are changing, we can not treat them separately

 $\rightarrow \text{Maxwell Equations}$

When the electric and magnetic fields are static, the are separate entities. But when they are changing, we can not treat them separately

$\rightarrow \text{Maxwell Equations}$


A changing magnetic field generates a changing electric field that generates a changing magnetic field.


When the electric and magnetic fields are static, the are separate entities. But when they are changing, we can not treat them separately

$\rightarrow \text{Maxwell Equations}$


A changing magnetic field generates a changing electric field that generates a changing magnetic field.

\rightarrow Wave traveling in the espace







 $lackbox{ We connect two rod to a battery}
ightarrow$ Electric Field




- $\hbox{$\stackrel{\bullet}{\scriptstyle}$ We connect two rod to a battery} \rightarrow \\ \hbox{$\stackrel{\bullet}{$\scriptstyle$}$ Electric Field}$
- lacktriangle The charge is re-distributed ightarrow Magnetic field







ightharpoonup sinusoidal voltage ightarrow Alternating Current




- ightharpoonup sinusoidal voltage ightarrow Alternating Current
- ightharpoonup variable Electric and Magnetic Fields.







Old Field lines fold back to connect to some of the new lines → closed loops




- Old Field lines fold back to connect to some of the new lines → closed loops
- ► The old field don't suddenly disappear



- Old Field lines fold back to connect to some of the new lines → closed loops
- ► The old field don't suddenly disappear
- Changing EF → changing MF → changing EF ... → so on



- Old Field lines fold back to connect to some of the new lines → closed loops
- The old field don't suddenly disappear
- ► changing EF \rightarrow changing MF \rightarrow changing EF ... \rightarrow so on
- This combination of changing EF and MF moving outward is self supporting, no longer depending on the antenna.



- Old Field lines fold back to connect to some of the new lines → closed loops
- The old field don't suddenly disappear
- ► changing EF \rightarrow changing MF \rightarrow changing EF ... \rightarrow so on
- This combination of changing EF and MF moving outward is self supporting, no longer depending on the antenna.
- They are on their way to distant points.

Characteristics of Electromagnetic Waves

▶ The field not far the antenna (Near Field) is complicated.

Characteristics of Electromagnetic Waves


- ▶ The field not far the antenna (Near Field) is complicated.
- ▶ The field far from the antenna ($Far\ Field$) is quite flat o Plane Waves

- ▶ The field not far the antenna (Near Field) is complicated.
- ▶ The field far from the antenna ($Far\ Field$) is quite flat o Plane Waves
- ► The field also travels in other directions, the strength is greater in the direction perpendicular to the antenna.


- ▶ The field not far the antenna (Near Field) is complicated.
- ▶ The field far from the antenna ($Far\ Field$) is quite flat o Plane Waves
- ► The field also travels in other directions, the strength is greater in the direction perpendicular to the antenna.
- ▶ The magnitude of \vec{E} and \vec{B} decreases with distance as 1/r.

- ▶ The field not far the antenna (Near Field) is complicated.
- ▶ The field far from the antenna ($Far\ Field$) is quite flat o Plane Waves
- ► The field also travels in other directions, the strength is greater in the direction perpendicular to the antenna.
- ▶ The magnitude of \vec{E} and \vec{B} decreases with distance as 1/r.
- $ightharpoonup ec{E} \perp ec{B}$, and they are \perp to the direction of propagation.


- ▶ The field not far the antenna (Near Field) is complicated.
- ▶ The field far from the antenna ($Far\ Field$) is quite flat o Plane Waves
- ► The field also travels in other directions, the strength is greater in the direction perpendicular to the antenna.
- ▶ The magnitude of \vec{E} and \vec{B} decreases with distance as 1/r.
- $ightharpoonup ec{E} \perp ec{B}$, and they are \perp to the direction of propagation.
- ightharpoonup of \vec{E} and \vec{B} are in phase.




► Transverse wave




- Transverse wave
- ► Waves of field, not of matter



- Transverse wave
- Waves of field, not of matter
- ► EM waves can propagate in empty space.




- Transverse wave
- Waves of field, not of matter
- ► EM waves can propagate in empty space.



- Transverse wave
- Waves of field, not of matter
- EM waves can propagate in empty space.

EM waves are produced by electric charges that are oscillating

 \rightarrow are undergoing into acceleration in general.




- Transverse wave
- Waves of field, not of matter
- EM waves can propagate in empty space.

EM waves are produced by electric charges that are oscillating \rightarrow are undergoing into acceleration in general.


Accelerated electric charges give rise to electromagnetic waves

Measuring the speed of light


Michelson's Experiment




$$c = f\lambda$$




$$c = f\lambda$$




$$c = f\lambda$$



$$c = f\lambda$$




$$c = f\lambda$$



Huygens' principle

Every point on a wave front can be considered as a source of tiny wavelets that spread out in the forward direction at the speed of the wave itself The new wave front is the envelope of all the wavelets—that is, the tangent to all of them.



What happens when waves impinge on an obstacle?

→ waves bend in behind an obstacle

- → waves bend in behind an obstacle
- \rightarrow The bending into the "shadow region" is known as diffraction


- → waves bend in behind an obstacle
- \rightarrow The bending into the "shadow region" is known as diffraction
- \rightarrow If the opening is much larger then than the wavelength, diffraction goes unnoticed.


- → waves bend in behind an obstacle
- \rightarrow The bending into the "shadow region" is known as diffraction
- \rightarrow If the opening is much larger then than the wavelength, diffraction goes unnoticed.




What happens when a waves front changes the medium?


What happens when a waves front changes the medium?







$$sin\theta_1 = \frac{v_1 t}{AD}, \quad sin\theta_2 = \frac{v_2 t}{AD}$$



$$sin\theta_1 = rac{v_1 t}{AD}, \quad sin\theta_2 = rac{v_2 t}{AD}$$

$$ightarrow rac{sin\theta_1}{sin\theta_2} = rac{v_1}{v_2}$$

The frequency does not change, but the wavelength does,

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

where,

$$\eta = \frac{c}{v} \tag{1}$$

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

where,

$$n = \frac{c}{v} \tag{1}$$

is the refraction index.

Why the velocity change?

Why the velocity change?

$$v = \frac{1}{\sqrt{\epsilon \mu}} \tag{2}$$

 $\mu \to {\rm The}$ resistance of a material to be penetrated by a magnetic field.

Why the velocity change?

$$v = \frac{1}{\sqrt{\epsilon \mu}} \tag{2}$$

 $\mu \to {\sf The}$ resistance of a material to be penetrated by a magnetic field.

 $\epsilon \to \mathsf{The}$ resistance of a material to be penetrated by an electric field.

Why the velocity change?

Why the velocity change?

$$v = \frac{1}{\sqrt{\epsilon \mu}} \tag{3}$$

 $\mu \to {\rm The}$ resistance of a material to be penetrated by a magnetic field.


Why the velocity change?

$$v = \frac{1}{\sqrt{\epsilon \mu}} \tag{3}$$


 $\mu \to {\sf The}$ resistance of a material to be penetrated by a magnetic field.

 $\epsilon \to \mathsf{The}$ resistance of a material to be penetrated by an electric field.

Example: Mirages



Interference: Youn's double-slit Experiment





(particle theory


prediction)



(c) Viewing screen (actual)

If light consists of tiny particles, we might expect to see two bright lines on a screen placed behind the slits as in (b). But instead a series of bright lines are seen, as in (c). Young was able to explain this result as a wave-interference phenomenon.

Interference in Thin Films



Light

Introduction Electric Field Magnetic Field Electromagnetic Radiation

Colors in a Thin Soap Film

